

# Introduction to Geometric Measure Theory

Leon Simon<sup>1</sup>

© Leon Simon 2017, 2018

<sup>1</sup>The research described here was partially supported by NSF grants DMS-9504456 & DMS-9207704 at Stanford University

# Contents

<b>Chapter 1: Preliminary Measure Theory</b>	<b>1</b>
§ 1 Basic Notions . . . . .	1
§ 2 Hausdorff Measure . . . . .	11
§ 3 Densities . . . . .	14
§ 4 Differentiation Theorems . . . . .	19
§ 5 Radon Measures, Representation Theorem . . . . .	28
<b>Chapter 2: Some Further Preliminaries from Analysis</b>	<b>43</b>
§ 1 Lipschitz Functions . . . . .	43
§ 2 BV Functions . . . . .	52
§ 3 The Area Formula . . . . .	57
§ 4 Submanifolds of $\mathbb{R}^{n+k}$ . . . . .	61
§ 5 First Variation of a Submanifold . . . . .	70
§ 6 Second Variation . . . . .	73
§ 7 Co-Area Formula and $C^1$ Sard Theorem . . . . .	75
<b>Chapter 3: Countably <math>n</math>-Rectifiable Sets</b>	<b>85</b>
§ 1 Basic Notions, Tangent Properties . . . . .	85
§ 2 Gradients, Jacobians, Area, Co-Area . . . . .	92
§ 3 Purely Unrectifiable Sets, Structure Theorem . . . . .	94
§ 4 Sets of Locally Finite Perimeter . . . . .	96
<b>Chapter 4: Rectifiable <math>n</math>-Varifolds</b>	<b>103</b>
§ 1 Basic Definitions and Properties . . . . .	104
§ 2 First Variation . . . . .	105
§ 3 Monotonicity Formulae in the Stationary Case . . . . .	107
§ 4 Monotonicity Formulae for $L^p$ Mean Curvature . . . . .	113
§ 5 Local Conical Approximation . . . . .	116
§ 6 Poincaré and Sobolev Inequalities . . . . .	118
§ 7 Miscellaneous Additional Consequences . . . . .	123
<b>Chapter 5: The Allard Regularity Theorem</b>	<b>127</b>
§ 1 Harmonic Approximation in the Smooth Case . . . . .	128
§ 2 Preliminaries, Lipschitz Approximation . . . . .	130
§ 3 Approximation by Harmonic Functions . . . . .	139
§ 4 The Tilt-Excess Decay Lemma . . . . .	140

§ 5	Main Regularity Theorem.....	145
§ 6	Some Initial Applications of the Allard Theorem .....	150
<b>Chapter 6:</b>	<b>Currents</b>	<b>155</b>
§ 1	Preliminaries: Vectors, Co-vectors, and Forms .....	155
§ 2	General Currents.....	159
§ 3	Integer Multiplicity Rectifiable Currents .....	172
§ 4	Slicing.....	180
§ 5	The Deformation Theorem .....	185
§ 6	Applications of Deformation Theorem.....	195
§ 7	The Flat Metric Topology .....	196
§ 8	The Rectifiability and Compactness Theorems.....	199
<b>Chapter 7:</b>	<b>Area Minimizing Currents</b>	<b>211</b>
§ 1	Basic Concepts .....	211
§ 2	Existence and Compactness Results .....	213
§ 3	Tangent Cones and Densities .....	217
§ 4	Some Regularity Results (Arbitrary Codimension).....	223
§ 5	Codimension 1 Theory .....	226
<b>Chapter 8:</b>	<b>Theory of General Varifolds</b>	<b>235</b>
§ 1	Basics, First Rectifiability Theorem .....	235
§ 2	First Variation .....	238
§ 3	Monotonicity and Consequences .....	240
§ 4	Constancy Theorem.....	243
§ 5	Varifold Tangents and Rectifiability Theorem.....	245
<b>Appendix A:</b>	<b>A General Regularity Theorem</b>	<b>255</b>
<b>Appendix B:</b>	<b>Non-existence of Stable Minimal Cones, <math>n \leq 6</math></b>	<b>259</b>

# Preface to the Tsinghua Lectures 2016

The present text is a revision and updating of the author's 1983 "Lectures on Geometric Measure Theory," and is meant to provide an introduction to the subject at beginning/intermediate graduate level. The present draft is still in rather rough form, with a generous scattering of (hopefully not serious, mainly expository) errors. During the Tsinghua lectures (March–April 2016) the notes will be further revised, with the ultimate aim of providing a useful and accessible introduction to the subject at the appropriate level.

The author would greatly appreciate feedback about errors and other deficiencies.

Leon Simon

Beijing, China, March 1, 2016

lsimon@stanford.edu

# Preface to the NTU Lectures 2018

The text here is basically the same as the text for the Tsinghua 2016 lectures, with some minor additions and corrections. The text will be further revised before the start of the NTU lectures on March 5, 2018. To assist in the assimilation of the material some exercises have been added at the end of each of the first six chapters.

During the NTU lectures (March 5–May 11 2018) the notes will be further revised, with the ultimate aim of providing a useful and accessible introduction to the subject at the appropriate level.

The author would greatly appreciate feedback about errors and other deficiencies.

Leon Simon

Taipei, Taiwan, March 5, 2018

lsimon@stanford.edu

Last updated: Wednesday March 7 at 4:30pm China Standard Time

# Notation

$\mathbb{Z} = \{0, \pm 1, \pm 2, \dots\}$ ,  $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$ .

$\bar{A}$  = closure of  $A$ , assuming  $A$  is a subset of some topological space  $X$

$A \subset\subset U$  means  $\bar{A}$  is a compact subset of  $U$

$B \setminus A = \{x \in B : x \notin A\}$

$\chi_A$  = indicator function of  $A$  ( $= 1$  at points of  $A$  and  $= 0$  at points not in  $A$ )

$\mathbb{1}_A$  = identity map  $A \rightarrow A$

$\mathcal{L}^n$  = Lebesgue outer measure in  $\mathbb{R}^n$

$B_\rho(y)$  = closed ball with center  $y$  radius  $\rho$  (more specifically denoted  $B_\rho^n(y)$  if we wish to emphasize that we are working in  $\mathbb{R}^n$ ). Thus  $B_\rho(y) = \{x \in \mathbb{R}^n : |x - y| \leq \rho\}$ , or, more generally,  $B_\rho(y) = \{x \in X : d(x, y) \leq \rho\}$  in any metric space  $X$ .

$\check{B}_\rho(y)$  = open ball  $= \{x \in X : d(x, y) < \rho\}$ ;

$\omega_k = \frac{\pi^{k/2}}{\int_0^\infty t^{k/2} e^{-t} dt}$  for  $k \geq 0$  (so  $\omega_k = \mathcal{L}^k(\{x \in \mathbb{R}^k : |x| \leq 1\})$  if  $k \in \{1, 2, \dots\}$ ).

$\eta_{y,\lambda} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  (for  $\lambda > 0$ ,  $y \in \mathbb{R}^n$ ) is defined by  $\eta_{y,\lambda}(x) = \lambda^{-1}(x - y)$ ; thus  $\eta_{y,1}$  is translation  $x \mapsto x - y$ , and  $\eta_{0,\lambda}$  is homothety  $x \mapsto \lambda^{-1}x$

$C^k(U, V)$  ( $U, V$  open subsets of Euclidean spaces  $\mathbb{R}^n$  and  $\mathbb{R}^m$  respectively) denotes the space of  $C^k$  maps from  $U$  into  $V$

$C_c^k(U, V) = \{\varphi \in C^k(U, V) : \varphi \text{ has compact support}\}$

$p_L$ , for any linear subspace  $L$  of  $\mathbb{R}^n$ , denotes orthogonal projection of  $\mathbb{R}^n$  onto  $L$

$Df$ , for  $f \in C^1(U, V)$ , is the derivative matrix with entries  $D_i f_j$  in the  $i$ -th row and  $j$ -th column, and  $|Df|^2 = \sum_{i=1}^n \sum_{j=1}^m (D_i f_j)^2$ .

$\nabla f$ , for  $f \in C^1(U, \mathbb{R})$ , denotes the gradient  $(D_1 f, \dots, D_n f)$  of  $f$ .

$\emptyset$  = the empty set.

$\text{spt } \mu$ , for a Borel measure  $\mu$  on a metric space  $X$ , is the support of  $\mu$ , i.e.  $\{x \in X : \mu(B_\rho(x)) > 0 \ \forall \rho > 0\}$  (which is a closed subset of  $X$ ).

$\text{diam } A$ , for any set  $A$  in a metric space  $X$ , denotes the diameter of the set  $A$ , i.e.  $\sup_{x,y \in A} d(x, y)$ , interpreted to be  $\infty$  if  $A$  is not bounded, 0 if  $A = \emptyset$

$W^{1,p}(\Omega)$ , for  $\Omega \subset \mathbb{R}^n$  open, will denote the Sobolev space of functions  $f : \Omega \rightarrow \mathbb{R}$  such that  $f, \nabla f \in L^p(\Omega)$ .

$\delta_{ij}$  = Kronecker delta ( $= 1$  if  $i = j$ , 0 if  $i \neq j$ ).



# Chapter 1

## Preliminary Measure Theory

§ 1	Basic Notions .....	1
§ 2	Hausdorff Measure .....	11
§ 3	Densities .....	14
§ 4	Differentiation Theorems .....	19
§ 5	Radon Measures, Representation Theorem .....	28

In this chapter we briefly review the basic theory of outer measure, which is based on Caratheodory’s definition of measurability. Hausdorff (outer) measure is discussed, including the main results concerning  $n$ -dimensional densities and the way in which they relate more general measures to Hausdorff measures. The final two sections of the chapter give the basic theory of Radon (outer) measures including the Riesz representation theorem and the standard differentiation theory for Radon measures.

For the first section of the chapter  $X$  will denote an abstract space, and later we impose further restrictions on  $X$  as appropriate. For example in the second and third sections  $X$  is a metric space and in the last section of the chapter we shall assume that  $X$  is a locally compact, separable metric space.

### 1 Basic Notions

Recall that an outer measure (sometimes simply called a *measure* if no confusion is likely to arise) on  $X$  is a monotone subadditive function  $\mu : \mathcal{X} \rightarrow [0, \infty]$  ( $\mathcal{X}$  = the collection of all subsets of  $X$ ) with  $\mu(\emptyset) = 0$ . Thus  $\mu(\emptyset) = 0$ , and

1.1 
$$\mu(A) \leq \sum_{j=1}^{\infty} \mu(A_j) \quad \text{whenever} \quad A \subset \bigcup_{j=1}^{\infty} A_j$$

with  $A, A_1, A_2, \dots$  any countable collection of subsets of  $X$ . Of course this in particular implies  $\mu(A) \leq \mu(B)$  whenever  $A \subset B \subset X$ , because we can take  $A_1 = B$  and  $A_j = \emptyset$  for each  $j \geq 2$ .

We adopt Caratheodory's notion of measurability:

**1.2 Definition:** A subset  $A \subset X$  is said to be  $\mu$ -measurable if

$$\mu(S) = \mu(S \setminus A) + \mu(S \cap A)$$

for each subset  $S \subset X$ . (Thus, roughly speaking,  $A$  is  $\mu$ -measurable if it “cuts every other set  $S$  additively with respect to  $\mu$ .”)

Since  $X \setminus (X \setminus A) = A$  we see that  $\mu$ -measurability of  $A$  is equivalent to  $\mu$ -measurability of  $X \setminus A$  for any set  $A \subset X$ .

**1.3 Remark:** Then the set  $A$  is  $\mu$ -measurable if and only if

$$\mu(S) \geq \mu(S \setminus A) + \mu(S \cap A)$$

for each subset  $S \subset X$  with  $\mu(S) < \infty$ , because this is trivially true when  $\mu(S) = \infty$ , and the reverse inequality also holds in both cases  $\mu(S) < \infty$  and  $\mu(S) = \infty$  by the subadditivity 1.1 of  $\mu$ .

Notice that the empty set  $\emptyset$  is  $\mu$ -measurable, as is any set of  $\mu$ -measure zero since in these cases the term  $\mu(S \cap A)$  on the right side of the inequality in Remark 1.3 is zero, and so the inequality in 1.3 holds trivially.

A key lemma, due to Caratheodory, asserts that such  $\mu$ -measurable sets form a  $\sigma$ -algebra, where the terminology is as follows:

**1.4 Definition:** A collection  $\mathcal{S}$  of subsets of  $X$  is a  $\sigma$ -algebra if

- (1)  $\emptyset, X \in \mathcal{S}$
- (2)  $A \in \mathcal{S} \Rightarrow X \setminus A \in \mathcal{S}$
- (3)  $A_1, A_2, \dots \in \mathcal{S} \Rightarrow \bigcup_{j=1}^{\infty} A_j \in \mathcal{S}$ .

**1.5 Remarks:** (1) Observe that then, since  $\bigcap_{j=1}^{\infty} A_j = X \setminus (\bigcup_{j=1}^{\infty} (X \setminus A_j))$ , we also have  $\bigcap_{j=1}^{\infty} A_j \in \mathcal{S}$  whenever  $A_1, A_2, \dots \in \mathcal{S}$ , by (2), (3).

(2) In the context of a fixed space  $X$ , it is easily checked that the intersection of any non-empty family of  $\sigma$ -algebras is again a  $\sigma$ -algebra, so there is always a smallest  $\sigma$ -algebra which contains a given collection of subsets of  $X$ —namely just take the intersection of all  $\sigma$ -algebras which contain the given collection of sets (this collection is non-empty because the collection of all subsets of  $X$  is a  $\sigma$ -algebra).

In view of Remark 1.5(2) above we can make the following definition:

**1.6 Definition:** If  $X$  is a topological space then we define *the Borel subsets of  $X$*  to be the smallest  $\sigma$ -algebra of subsets of  $X$  which contains all the open sets (same as the smallest  $\sigma$ -algebra containing all the closed sets since  $\sigma$ -algebras are closed under complementation).

As mentioned above, we have the following lemma:

**1.7 Lemma.** *The collection  $\mathcal{M}$  of all  $\mu$ -measurable subsets is a  $\sigma$ -algebra which includes all subsets of  $X$  of  $\mu$ -measure zero.*

**1.8 Remark:** In the course of the proof we shall establish the important additional fact that for  $\mu$ -measurable sets  $A_j$ ,  $j = 1, 2, \dots$ ,

$$A_1, A_2, \dots \text{ pairwise disjoint} \Rightarrow \mu(S \cap (\cup_{j=1}^{\infty} A_j)) = \sum_{j=1}^{\infty} \mu(S \cap A_j)$$

for each subset  $S \subset X$ .

**Proof of Lemma 1.7 and Remark 1.8:** We already noted above that Properties 1.4(1) and 1.4(2) are trivially checked direct from the definition of measurability.

Checking 1.4(3) involves several steps:

Step 1:  $A_1, A_2 \in \mathcal{M} \Rightarrow A_1 \cup A_2 \in \mathcal{M}$ . To check this we first use Definition 1.2 with  $A = A_2$  and with  $S \setminus A_1$  in place of  $S$  and  $\mu(S) < \infty$  to give

$$\begin{aligned} \mu(S \setminus (A_1 \cup A_2)) &= \mu((S \setminus A_1) \setminus A_2) \\ &= \mu(S \setminus A_1) - \mu((S \setminus A_1) \cap A_2) \end{aligned}$$

and then use Definition 1.2 with  $A = A_1$  on the right side together with the subadditivity of  $\mu$  to give

$$\begin{aligned} \mu(S \setminus (A_1 \cup A_2)) &= \mu(S) - \mu(S \cap A_1) - \mu((S \setminus A_1) \cap A_2) \\ &\leq \mu(S) - \mu((S \cap A_1) \cup ((S \setminus A_1) \cap A_2)) \\ &= \mu(S) - \mu(S \cap (A_1 \cup A_2)), \end{aligned}$$

so  $A_2 \cup A_1$  is  $\mu$ -measurable by Remark 1.3. Notice that the first line above gives

$$\mu(S) - \mu(S \setminus (A_1 \cup A_2)) = \mu(S \cap A_1) + \mu((S \setminus A_1) \cap A_2)$$

and since we have established  $\mu(S) - \mu(S \setminus (A_1 \cup A_2)) = \mu(S \cap (A_1 \cup A_2))$  we thus conclude the important additional fact that if  $A_1, A_2$  are *disjoint* and  $\mu$ -measurable then

$$(\ddagger) \quad \mu(S \cap (A_1 \cup A_2)) = \mu(S \cap A_1) + \mu(S \cap A_2) \text{ for every } S \subset X.$$

Step 2:  $A_1, A_2 \in \mathcal{M} \Rightarrow A_1 \cap A_2 \in \mathcal{M}$ . This is clear from Step 1 and 1.4(2) because  $A_1 \cap A_2 = X \setminus ((X \setminus A_1) \cup (X \setminus A_2))$ .

Step 3: For each  $N = 1, 2, \dots$ ,  $A_1, A_2, \dots, A_N \in \mathcal{M} \Rightarrow \cup_{j=1}^N A_j \in \mathcal{M}$ , which follows from Step 2 by induction on  $N$ . Using the additional additivity conclusion (§) of Step 1 we also conclude the additivity  $\mu(S \cap (\cup_{j=1}^N A_j)) = \sum_{j=1}^N \mu(S \cap A_j)$  provided  $A_1, A_2, \dots, A_N$  are pairwise disjoint sets in  $\mathcal{M}$ .

Step 4: If  $A_1, A_2, \dots$  are pairwise disjoint sets in  $\mathcal{M}$  then  $\cup_{j=1}^\infty A_j \in \mathcal{M}$  and furthermore  $\mu(S \cap (\cup_{j=1}^\infty A_j)) = \sum_{j=1}^\infty \mu(S \cap A_j)$  for each  $S \subset X$ . To check this we use the conclusions of Step 3 to observe

$$\begin{aligned} \mu(S) &= \mu(S \cap (\cup_{j=1}^N A_j)) + \mu(S \setminus (\cup_{j=1}^N A_j)) \\ &\geq \mu(S \cap (\cup_{j=1}^N A_j)) + \mu(S \setminus (\cup_{j=1}^\infty A_j)) \\ &= \sum_{j=1}^N \mu(S \cap A_j) + \mu(S \setminus (\cup_{j=1}^\infty A_j)). \end{aligned}$$

Since  $\sum_{j=1}^N \mu(S \cap A_j) \rightarrow \sum_{j=1}^\infty \mu(S \cap A_j) \geq \mu(S \cap (\cup_{j=1}^\infty A_j))$ , in view of Remark 1.3 this completes the proof of Step 3, and also establishes the additivity property of Remark 1.8.

Step 5:  $A_1, A_2, \dots \in \mathcal{M} \Rightarrow \cup_{j=1}^\infty A_j \in \mathcal{M}$  (i.e. we do indeed have that  $\mathcal{M}$  has property 1.4(3)). To check this, observe that  $\cup_{j=1}^\infty A_j = \cup_{j=1}^\infty \tilde{A}_j$ , where  $\tilde{A}_j = A_j \setminus (\cup_{i=0}^{j-1} A_i)$ , with  $A_0 = \emptyset$ . Then  $\tilde{A}_j \in \mathcal{M}$  by Step 2, Step 3 and 1.4(2). Since the  $\tilde{A}_j$  are pairwise disjoint we can then apply Step 4 to complete the proof.  $\square$

Observe that by 1.8 we have

$$1.9 \quad A_j \text{ } \mu\text{-measurable, } A_j \subset A_{j+1} \forall j \geq 1 \Rightarrow \lim_{j \rightarrow \infty} \mu(A_j) = \mu(\cup_{j=1}^\infty A_j),$$

because we can write  $\cup_{j=1}^\infty A_j = \cup_{j=1}^\infty (A_j \setminus A_{j-1})$  with  $A_0 = \emptyset$ , and hence, by 1.8,

$$\begin{aligned} \mu(\cup_{j=1}^\infty A_j) &= \sum_{j=1}^\infty \mu(A_j \setminus A_{j-1}) = \lim \sum_{j=1}^n \mu(A_j \setminus A_{j-1}) \\ &= \lim \mu(\cup_{j=1}^n (A_j \setminus A_{j-1})) = \lim \mu(A_n), \end{aligned}$$

where at the last step we used  $\cup_{j=1}^n (A_j \setminus A_{j-1}) = A_n$ .

If  $A_1 \supset A_2 \supset \dots$  then, for each  $i$ ,  $A_i \cap \cap_{j=1}^\infty A_j = \cup_{j=1}^\infty (A_i \setminus A_j)$ , and hence 1.9 implies  $\lim_{j \rightarrow \infty} \mu(A_i \setminus A_j) = \mu(A_i \setminus \cap_{j=1}^\infty A_j)$ , and if  $\mu(A_i) < \infty$  this gives  $\mu(A_i) - \lim_j \mu(A_j) = \mu(A_i) - \mu(\cap_{j=1}^\infty A_j)$ , and hence

$$\begin{aligned} 1.10 \quad A_j \text{ } \mu\text{-measurable and } A_{j+1} \subset A_j \text{ for each } j = 1, 2, \dots \\ \Rightarrow \lim_{j \rightarrow \infty} \mu(A_j) = \mu(\cap_{j=1}^\infty A_j), \text{ provided } \mu(A_i) < \infty \text{ for some } i. \end{aligned}$$

An outer measure  $\mu$  on  $X$  is said to be *regular* if for each subset  $A \subset X$  there is a  $\mu$ -measurable subset  $B \supset A$  with  $\mu(B) = \mu(A)$ .

**1.11 Remark:** If  $A_i \subset A_{i+1} \forall i$  and  $\mu$  is regular, then the identity in 1.9 is valid, i.e.

$$\lim_{i \rightarrow \infty} \mu(A_i) = \mu(\cup_{i=1}^{\infty} A_i),$$

even if the  $A_i$  are not assumed to be  $\mu$ -measurable, because for each  $i$  we can select a  $\mu$ -measurable set  $\tilde{A}_i \supset A_i$  with  $\mu(\tilde{A}_i) = \mu(A_i)$ , and then  $\hat{A}_i = \cap_{j=i}^{\infty} \tilde{A}_j (\supset A_i)$  is increasing with  $\mu(\hat{A}_i) = \mu(A_i)$  and  $\lim \mu(A_i) \leq \mu(\cup_{i=1}^{\infty} A_i) \leq \mu(\cup_{i=1}^{\infty} \hat{A}_i) = \lim \mu(\hat{A}_i)$  (by 1.9)  $= \lim \mu(A_i)$ .

We have the following additional corollary of 1.9.

**1.12 Corollary (Egoroff's Theorem.)** *If  $A \subset X$  is  $\mu$ -measurable with finite measure, if  $f_k : A \rightarrow [-\infty, \infty]$  are  $\mu$ -measurable for each  $k = 1, 2, \dots$ , and if  $\lim f_k(x) = 0$  for each  $x \in A$ , then for each  $\varepsilon > 0$  there is a  $\mu$ -measurable subset  $B \subset A$  with  $f_k \rightarrow 0$  uniformly on  $B$  and  $\mu(A \setminus B) < \varepsilon$ .*

**Proof:** For each  $x \in A$  and each  $j = 1, 2, \dots$  there is an  $\ell$  such that  $|f_k(x)| < 1/j$  for all  $k \geq \ell$ , so  $x \in B_{j,\ell} = \cap_{k=\ell}^{\infty} \{x \in A : |f_k(x)| < 1/j\}$ . Thus  $A = \cup_{\ell=1}^{\infty} B_{j,\ell}$  for each  $j$  and of course  $B_{j,\ell} \subset B_{j,\ell+1}$ , so by 1.9 we have  $\mu(A) = \lim_{\ell \rightarrow \infty} \mu(B_{j,\ell})$  for each  $j$ . In particular for each  $\varepsilon > 0$  and each  $j = 1, 2, \dots$  there is  $\ell_j$  with  $\mu(A \setminus B_{j,\ell_j}) < \varepsilon 2^{-j}$ . Hence, with  $B = \cap_{j=1}^{\infty} B_{j,\ell_j}$ ,  $\mu(A \setminus B) = \mu(\cup_{j=1}^{\infty} (A \setminus B_{j,\ell_j})) \leq \varepsilon \sum_{j=1}^{\infty} 2^{-j} = \varepsilon$  and, for each  $j = 1, 2, \dots$ ,  $|f_k(x)| < 1/j$  for all  $x \in B (\subset B_{j,\ell_j})$  and all  $k \geq \ell_j$ .  $\square$

In case  $X$  is a topological space, an outer measure  $\mu$  on  $X$  is said to be *Borel-regular* if all Borel sets (see Remark 1.5(2)) are  $\mu$ -measurable and if for each subset  $A \subset X$  there is a Borel set  $B \supset A$  such that  $\mu(B) = \mu(A)$ . (Notice that this does *not* imply  $\mu(B \setminus A) = 0$  unless  $A$  is  $\mu$ -measurable and  $\mu(A) < \infty$ .)

**1.13 Remark:** There is a close relationship between Borel regular outer measures on a topological space  $X$  and abstract Borel measures  $\mu_0$  on  $X$ . (Recall that a Borel measure  $\mu_0$  on  $X$  is a map  $\mu_0 : \{\text{all Borel sets}\} \rightarrow [0, \infty]$  such that (i)  $\mu_0(\emptyset) = 0$ , and (ii)  $\mu_0(\cup_{j=1}^{\infty} B_j) = \sum_{j=1}^{\infty} \mu_0(B_j)$  whenever  $B_1, B_2, \dots$  are pairwise disjoint Borel sets in  $X$ .) In fact if  $\mu_0$  is such a Borel measure on  $X$  then

$$\mu(A) = \inf_{B \text{ Borel}, B \supset A} \mu_0(B)$$

defines a Borel regular outer measure on  $X$  which agrees with  $\mu_0$  on the Borel sets; to check  $\mu$ -measurability of any Borel set  $B$  we just check the inequality in 1.3 by first choosing a Borel set  $C \supset S$  with  $\mu(C) = \mu(S)$ . Conversely, if  $\mu$  is a Borel regular outer

measure on  $X$  then the restriction of  $\mu$  to the Borel sets gives us a Borel measure  $\mu_0$  on  $X$ .

Given any subset  $Y \subset X$  and any outer measure  $\mu$  on  $X$ , we can define a new outer measure  $\mu \llcorner Y$  on  $X$  by

$$1.14 \quad (\mu \llcorner Y)(Z) = \mu(Y \cap Z), \quad Z \subset X.$$

One readily checks (by using  $S \cap Y$  in place of  $S$  in Definition 1.2) that all  $\mu$ -measurable subsets are also  $(\mu \llcorner Y)$ -measurable (even if  $Y$  is *not*  $\mu$ -measurable). It is also easy to check that  $\mu \llcorner Y$  is Borel regular whenever  $\mu$  is, provided  $Y$  is  $\mu$ -measurable with  $\mu(Y) < \infty$ . To check this, first use Borel regularity of  $\mu$  to pick a Borel set  $B_1$  with  $B_1 \supset Y$  and  $\mu(B_1 \setminus Y) = 0$  and a Borel set  $B_2 \supset B_1 \setminus Y$  with  $\mu(B_2) = 0$ . Then given an arbitrary set  $A \subset X$  we have

$$\begin{aligned} A &= (A \cap Y) \cup (A \setminus Y) \subset (A \cap Y) \cup (X \setminus Y) \\ &= (A \cap Y) \cup (X \setminus B_1) \cup (B_1 \setminus Y) \subset (A \cap Y) \cup (X \setminus B_1) \cup B_2. \end{aligned}$$

Finally select a Borel set  $B_3 \supset A \cap Y$  with  $\mu(B_3) = \mu(A \cap Y)$  and observe that then  $A \subset (X \setminus B_1) \cup B_2 \cup B_3$  (which is a Borel set) and  $(\mu \llcorner Y)((X \setminus B_1) \cup B_2 \cup B_3) = (\mu \llcorner Y)(A)$ .

The following theorem, due to Caratheodory and applicable in case  $X$  is a metric space with metric  $d$ , is particularly useful. In the statement we use the notation

$$\text{dist}(A, B) = \inf\{d(a, b) : a \in A, b \in B\},$$

interpreted as  $\infty$  if  $A$  or  $B$  is empty.

**1.15 Theorem (Caratheodory's Criterion.)** *If  $X$  is a metric space with metric  $d$  and if  $\mu$  is an outer measure on  $X$  with the property*

$$\mu(A \cup B) = \mu(A) + \mu(B) \text{ for all sets } A, B \subset X \text{ with } \text{dist}(A, B) > 0,$$

*then all Borel sets are  $\mu$ -measurable.*

**Proof:** Since the measurable sets form a  $\sigma$ -algebra, it is enough to prove that all closed sets are  $\mu$ -measurable (because by definition the Borel sets are the smallest  $\sigma$ -algebra containing all the closed sets), so that by Remark 1.3 we have only to check that

$$(\ddagger) \quad \mu(S) \geq \mu(S \setminus C) + \mu(S \cap C)$$

for all sets  $S \subset X$  with  $\mu(S) < \infty$  and for all closed sets  $C \subset X$ .

Let  $C_j = \{x \in X : \text{dist}(x, C) \leq 1/j\}$ . Then  $\text{dist}(S \setminus C_j, S \cap C) > 0$ , hence

$$\mu(S) \geq \mu((S \setminus C_j) \cup (S \cap C)) = \mu(S \setminus C_j) + \mu(S \cap C),$$

and we will have (‡) if we can show  $\lim_{j \rightarrow \infty} \mu(S \setminus C_j) = \mu(S \setminus C)$ . To check this, note that since  $C$  is closed we can write

$$S \setminus C = \{x \in X : \text{dist}(x, C) > 0\} = (S \setminus C_j) \cup (\cup_{k=j}^{\infty} R_k), \quad j = 1, 2, \dots,$$

where  $R_k = \{x \in S : \frac{1}{k+1} < \text{dist}(x, C) \leq \frac{1}{k}\}$ . But then by subadditivity of  $\mu$  we have

$$\mu(S \setminus C_j) \leq \mu(S \setminus C) \leq \mu(S \setminus C_j) + \sum_{k=j}^{\infty} \mu(R_k),$$

and hence we will have  $\lim_{j \rightarrow \infty} \mu(S \setminus C_j) = \mu(S \setminus C)$  as required, provided only that  $\sum_{k=1}^{\infty} \mu(R_k) < \infty$ .

To check this we note that  $\text{dist}(R_i, R_j) > 0$  if  $j \geq i + 2$ , and hence by the hypothesis of the theorem and induction on  $N$  we have, for each  $N \geq 1$ ,

$$\sum_{k=1}^N \mu(R_{2k}) = \mu(\cup_{k=1}^N R_{2k}) \leq \mu(S) < \infty$$

and

$$\sum_{k=1}^N \mu(R_{2k-1}) = \mu(\cup_{k=1}^N R_{2k-1}) \leq \mu(S) < \infty. \quad \square$$

A key example to which the above is applicable is of course Lebesgue measure  $\mathcal{L}^n$  on  $\mathbb{R}^n$ . This is defined (as usual) as follows: If  $\mathcal{K}$  denotes the collection of all  $n$ -dimensional intervals  $I$  of the form  $I = (a_1, b_1) \times (a_2, b_2) \times \dots \times (a_n, b_n)$ , where  $a_i, b_i \in \mathbb{R}$  and  $b_i - a_i > 0$ , and if  $|I| = \text{volume of } I (= (b_1 - a_1) \dots (b_n - a_n))$ , then

$$1.16 \quad \mathcal{L}^n(A) = \inf \sum_j |I_j|$$

where the inf is taken over all countable (or finite) collections  $\{I_1, I_2, \dots\} \subset \mathcal{K}$  with  $A \subset \cup_j I_j$ .

Clearly for any  $I = (a_1, b_1) \times (a_2, b_2) \times \dots \times (a_n, b_n) \in \mathcal{K}$ , by using “slight fattenings” of a sufficiently fine subdivision of each edge  $(a_j, b_j)$ , for each  $\delta, \varepsilon > 0$  we can find open  $J_1, \dots, J_N \in \mathcal{K}$  with  $I \subset \cup_{j=1}^N J_j$ ,  $\sum_{j=1}^N |J_j| < |I| + \varepsilon$  and  $\text{diam } J_j < \delta$  for each  $j = 1, \dots, N$ , so for each  $\delta > 0$  we can alternatively (and equivalently) define

$$1.16' \quad \mathcal{L}^n(A) = \inf \sum_j |I_j|$$

where the inf is taken over all countable (or finite) collections  $\{I_1, I_2, \dots\} \subset \mathcal{K}$  with  $A \subset \cup_j I_j$  and  $\text{diam } I_j < \delta$ .

Evidently  $\mathcal{L}^n$ , so defined, has the additivity property needed to apply Theorem 1.15, so all Borel sets in  $\mathbb{R}^n$  are  $\mathcal{L}^n$ -measurable, and direct from the definition of  $\mathcal{L}^n$  it is also evident that for each  $A \subset \mathbb{R}^n$  there is a sequence of open sets  $U_1, U_2$  with  $A \subset \cap_j U_j$  and  $\mathcal{L}^n(A) = \mathcal{L}^n(\cap_j U_j)$ . So  $\mathcal{L}^n$  is a Borel regular outer measure on  $\mathbb{R}^n$  and

$$1.17 \quad \mathcal{L}^n(A) = \inf_{U \text{ open}, U \supset A} \mathcal{L}^n(U) \quad \forall A \subset \mathbb{R}^n.$$

Note also that if  $U$  is any bounded open subset of  $\mathbb{R}^n$  we can always find closed balls  $B_{\rho_1}(x_1), B_{\rho_2}(x_2), \dots$  with

$$1.18 \quad B_{\rho_i}(x_i) \cap B_{\rho_j}(x_j) = \emptyset \quad \forall i \neq j, \cup_j B_{\rho_j}(x_j) \subset U, \mathcal{L}^n(U \setminus (\cup_{j=1}^{\infty} B_{\rho_j}(x_j))) = 0.$$

(See problem 1.5 of Ch.1 problems.) In view of 1.17, 1.18 it then evidently follows that  $\mathcal{L}^n(A)$  is invariant under application of orthogonal transformations to the set  $A$ . Since  $\mathcal{L}^n(A)$  is also trivially invariant under translation of  $A$  we thus have

$$1.19 \quad \mathcal{L}^n(y + Q(A)) = \mathcal{L}^n(A) \quad \forall y \in \mathbb{R}^n, \forall \text{orthogonal } Q : \mathbb{R}^n \rightarrow \mathbb{R}^n, \forall A \subset \mathbb{R}^n.$$

As a corollary of the above invariance property we establish the classical area formula for linear maps  $\mathbb{R}^n \rightarrow \mathbb{R}^n$ :

**1.20 Corollary.** *Suppose  $\tau : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is linear and  $A \subset \mathbb{R}^n$ . Then*

$$\mathcal{L}^n(\tau(A)) = |\det \tau| \mathcal{L}^n(A), \quad A \subset \mathbb{R}^n.$$

**Proof of 1.20:** If  $\det \tau = 0$  then  $\tau(A)$  is contained in an  $(n-1)$ -dimensional subspace and the theorem trivially holds (with both sides zero) in this case, so we can assume without loss of generality that  $\det \tau \neq 0$ . Then the symmetric transformation  $\tau^* \circ \tau$  has positive eigenvalues  $\lambda_1, \dots, \lambda_n$ , so  $(\det \tau)^2 = \lambda_1 \cdots \lambda_n$  and by the spectral theorem there is an orthogonal transformation  $Q : \mathbb{R}^n \rightarrow \mathbb{R}^n$  with  $Q^* \circ \tau^* \circ \tau \circ Q = \Lambda$ , where  $\Lambda$  is represented by a diagonal matrix with diagonal entries  $\lambda_1, \dots, \lambda_n$ . Hence  $\Gamma \circ Q^* \circ \tau^* \circ \tau \circ Q \circ \Gamma = I$ , where  $\Gamma = \sqrt{\Lambda^{-1}}$  (i.e.  $\Gamma$  is represented by a diagonal matrix with diagonal entries  $\lambda_1^{-1/2}, \dots, \lambda_n^{-1/2}$ ), so  $\tau \circ Q \circ \Gamma$  is an orthogonal transformation  $P$ . Hence  $\tau(A) = P(\Gamma^{-1}(Q^*A))$  and, by the invariance 1.19,  $\mathcal{L}^n(\tau(A)) = \mathcal{L}^n(\Gamma^{-1}(Q^*A))$ , and since  $\Gamma^{-1}J = (\lambda_1^{1/2}(a_1, b_1) \times \cdots \times \lambda_n^{1/2}(a_n, b_n))$  for each  $n$ -dimensional interval  $J = (a_1, b_1) \times \cdots \times (a_n, b_n)$ , Definition 1.16 implies  $\mathcal{L}^n(\Gamma^{-1}(Q^*A)) = |\det \tau| \mathcal{L}^n(Q^*A) = |\det \tau| \mathcal{L}^n(A)$ , where we again used the invariance 1.19.  $\square$

We next prove some important regularity properties for Borel regular measures which have an “open  $\sigma$ -finiteness property” as in the following definition:

**1.21 Definition:** We say a Borel regular measure  $\mu$  on a topological space  $X$  is “open  $\sigma$ -finite” if  $X = \cup_j V_j$  where  $V_j$  is open in  $X$  and  $\mu(V_j) < \infty$  for each  $j = 1, 2, \dots$

Of course  $\mu$  automatically satisfies such a condition if  $\mu(X) < \infty$  (then we just take  $V_j = X$  for each  $j$ ) or if  $X$  is a separable metric space and  $\mu$  is locally finite (i.e.  $x \in X \Rightarrow \mu(B_\rho(x)) < \infty$  for some  $\rho > 0$ ).

The following theorem tells us that open  $\sigma$ -finite Borel regular measures have a property analogous to the property 1.17 of  $\mathcal{L}^n$ , at least in a large class of topological spaces  $X$ , including all metric spaces:

**1.22 Theorem.** Suppose  $X$  is a topological space with the property that every closed subset of  $X$  is the countable intersection of open sets (this trivially holds e.g. if  $X$  is a metric space), and suppose  $\mu$  is an open  $\sigma$ -finite (as in 1.21 above) Borel-regular measure on  $X$ . Then

$$(1) \quad \mu(A) = \inf_{U \text{ open}, U \supset A} \mu(U)$$

for each subset  $A \subset X$ , and

$$(2) \quad \mu(A) = \sup_{C \text{ closed}, C \subset A} \mu(C)$$

for each  $\mu$ -measurable subset  $A \subset X$ .

**1.23 Remark:** In case  $X$  is a Hausdorff space (so compact sets in  $X$  are closed) which is  $\sigma$ -compact (i.e.  $X = \cup_j K_j$  with  $K_j$  compact for each  $j$ ), then the conclusion (2) in the above theorem guarantees that

$$\mu(A) = \sup_{K \text{ compact}, K \subset A} \mu(K)$$

for each  $\mu$ -measurable subset  $A \subset X$  with  $\mu(A) < \infty$ , because under the above conditions on  $X$  any closed set  $C$  can be written as the union of an increasing sequence of compact sets.

**Proof of 1.22:** We assume first that  $\mu(X) < \infty$ . First note that in this case (2) can be proved by applying (1) to the complement  $X \setminus A$ . Also, by Borel regularity of the measure  $\mu$ , it is enough to prove (1) in case  $A$  is a Borel set. Then let

$$\mathcal{A} = \{ \text{Borel sets } A \subset X : (1) \text{ holds} \}.$$

Trivially  $\mathcal{A}$  contains all open sets, and we claim that  $\mathcal{A}$  is closed under both countable unions and intersections, which we check as follows:

If  $A_1, A_2, \dots \in \mathcal{A}$  then for any given  $\varepsilon > 0$  there are open  $U_1, U_2, \dots$  with  $U_j \supset A_j$  and  $\mu(U_j \setminus A_j) \leq 2^{-j} \varepsilon$ . Then  $(\cup_j U_j) \setminus (\cup_k A_k) = \cup_j (U_j \setminus (\cup_k A_k)) \subset \cup_j (U_j \setminus A_j)$ . Also  $(\cap_j U_j) \setminus (\cap_k A_k) = (\cap_j U_j) \cap (\cup_k (X \setminus A_k)) = \cup_k ((\cap_j U_j) \setminus A_k) \subset \cup_k (U_k \setminus A_k)$ . So by subadditivity we have both  $\mu(\cup_{j=1}^{\infty} U_j \setminus (\cup_k A_k)) < \varepsilon$  and  $\lim_{N \rightarrow \infty} \mu(\cap_{j=1}^N U_j \setminus (\cap_k A_k)) = \mu(\cap_{j=1}^{\infty} U_j \setminus (\cap_k A_k)) < \varepsilon$ , so both  $\cup_k A_k$  and  $\cap_k A_k$  are in  $\mathcal{A}$  as claimed.

In particular  $\mathcal{A}$  must also contain the *closed sets*, because we are assuming any closed set in  $X$  can be written as a countable intersection of open sets. Notice however that at this point it is not clear that  $\mathcal{A}$  is a  $\sigma$ -algebra since it is not clear that  $\mathcal{A}$  is closed under complementation. For this reason, we let  $\tilde{\mathcal{A}} = \{A \in \mathcal{A} : X \setminus A \in \mathcal{A}\}$ , which we claim is a  $\sigma$ -algebra, since it clearly has properties 1.4 (1), (2) of  $\sigma$ -algebra, and if  $A_1, A_2, \dots \in \tilde{\mathcal{A}}$  then  $X \setminus A_1, X \setminus A_2, \dots \in \mathcal{A}$  and hence both  $\cup_{j=1}^{\infty} A_j$  and  $X \setminus ((\cup_{j=1}^{\infty} A_j)) (= \cap_{j=1}^{\infty} (X \setminus A_j))$  are in  $\mathcal{A}$  (because  $\mathcal{A}$  is closed under countable unions and intersections); thus  $\cup_{j=1}^{\infty} A_j \in \tilde{\mathcal{A}}$

and indeed  $\tilde{\mathcal{A}}$  is a  $\sigma$ -algebra as claimed. Thus  $\tilde{\mathcal{A}}$  is a  $\sigma$ -algebra containing all the closed sets, and hence  $\tilde{\mathcal{A}}$  contains all the Borel sets. Thus  $\mathcal{A}$  contains all the Borel sets (so actually we conclude that  $\mathcal{A}$  is equal to the collection of all Borel subsets of  $X$ ) and (1),(2) are both proved in case  $\mu(X) < \infty$ .

In the case  $\mu(X) = \infty$  it still suffices to prove (1) when  $A$  is a Borel set. For each  $j = 1, 2, \dots$  apply the previous case  $\mu(X) < \infty$  to the measure  $\mu \llcorner V_j$ ,  $j = 1, 2, \dots$ , with  $V_j$  as in 1.21. Then for each  $\varepsilon > 0$  we can select an open  $U_j \supset A$  such that

$$\mu(U_j \cap V_j \setminus A) < \frac{\varepsilon}{2^j},$$

and hence (summing over  $j$ )

$$\mu(\cup_{j=1}^{\infty} (U_j \cap V_j) \setminus A) < \varepsilon.$$

Since  $\cup_{j=1}^{\infty} (U_j \cap V_j)$  is open and contains  $A$ , this completes the proof of (1).

(2) for the case when  $\mu(X) = \infty$  also follows by applying (2) in the finite measure case to the measure  $\mu \llcorner V_j$ , thus giving, for each  $\varepsilon > 0$  and each  $j = 1, 2, \dots$ , a closed  $C_j \subset A$  with  $\mu(A \cap V_j \setminus C_j) < 2^{-j} \varepsilon$ . Since  $(\cup_{j=1}^{\infty} V_j) \setminus (\cup_{k=1}^{\infty} C_k) = \cup_{j=1}^{\infty} (V_j \setminus (\cup_{k=1}^{\infty} C_k)) \subset \cup_{j=1}^{\infty} (V_j \setminus C_j)$ , this gives, by countable subadditivity of  $\mu$ ,  $\mu(A \setminus (\cup_{k=1}^{\infty} C_k)) < \varepsilon$ . Thus either  $\mu(A) = \infty$  and  $\mu(\cup_{j=1}^N C_j) \rightarrow \infty$  or else  $\mu(A) < \infty$  and  $\mu(A \setminus (\cup_{j=1}^N C_j)) < \varepsilon$  for sufficiently large  $N$ . In either case this completes the proof of (2).  $\square$

Using the above theorem, we can now prove Lusin's Theorem:

**1.24 Theorem (Lusin's Theorem.)** *Let  $\mu$  be a Borel regular outer measure on a topological space  $X$  having the property that every closed subset can be expressed as the countable intersection of open sets (e.g.  $X$  is a metric space), let  $A$  be any  $\mu$ -measurable subset of  $X$  with  $\mu(A) < \infty$ , and let  $f : A \rightarrow \mathbb{R}$  be  $\mu$ -measurable. Then for each  $\varepsilon > 0$  there is a closed set  $C \subset X$  with  $C \subset A$ ,  $\mu(A \setminus C) < \varepsilon$ , and  $f|_C$  continuous.*

**1.25 Remark.** There are various scenarios which make it possible to drop the hypothesis that  $\mu(A) < \infty$ . For example if  $X$  is a metric space with each closed ball compact and of finite  $\mu$ -measure, then if  $A$  is  $\mu$ -measurable we can take any  $x_0 \in X$  and apply the above theorem in annular regions  $\check{B}_j(x_0) \setminus \check{B}_{j-1}(x_0)$  ( $= \check{B}_1(x_0)$  if  $j = 1$ ) to the subsets  $A_j = A \cap \check{B}_j(x_0) \setminus \check{B}_{j-1}(x_0)$ . This gives compact sets  $C_j \subset A_j$  with  $\mu(A_j \setminus C_j) < \varepsilon/2^j$  and  $f|_{C_j}$  is continuous. Then observe that  $C = \cup_{j=1}^{\infty} C_j$  is closed, because for each  $R > 0$  there are only finitely many  $j$  with  $C_j \cap B_R(x_0) \neq \emptyset$ . Also  $f|_C$  is continuous (because  $d(C_i, C_j) > 0 \forall i \neq j$  by the compactness of each  $C_i$ ), and  $\mu(A \setminus C) = \sum_{j=1}^{\infty} \mu(A_j \setminus C) < \sum_{j=1}^{\infty} \varepsilon/2^j = \varepsilon$ .

**Proof of Theorem 1.24:** For each  $i = 1, 2, \dots$  and  $j = 0, \pm 1, \pm 2, \dots$  let

$$A_{ij} = f^{-1}[\frac{j-1}{i}, \frac{j}{i}),$$

so that  $A_{ij}, j = 0, \pm 1, \pm 2, \dots$ , are pairwise disjoint sets in  $A$  and  $\bigcup_{j=-\infty}^{\infty} A_{ij} = A$  for each  $i = 1, 2, \dots$ . By the remarks following 1.14, we know that  $\mu \ll A$  is Borel regular, and since it is finite we can apply Theorem 1.22, so for each given  $\varepsilon > 0$  there are closed sets  $C_{ij}$  in  $X$  with  $C_{ij} \subset A_{ij}$  and  $\mu(A_{ij} \setminus C_{ij}) = (\mu \ll A)(A_{ij} \setminus C_{ij}) < 2^{-i-|j|-2}\varepsilon$ , hence  $\mu(A_{ij} \setminus (\bigcup_{\ell=-\infty}^{\infty} C_{i\ell})) < 2^{-i-|j|-2}\varepsilon$ , hence  $\mu(A \setminus (\bigcup_{j=-\infty}^{\infty} C_{ij})) < 2^{-i}\varepsilon$ . So for each  $i = 1, 2, \dots$  there is an integer  $J(i)$  with  $\mu(A \setminus (\bigcup_{|j| \leq J(i)} C_{ij})) < 2^{-i}\varepsilon$ . Since  $A \setminus (\bigcap_{i=1}^{\infty} (\bigcup_{|j| \leq J(i)} C_{ij})) = \bigcup_{i=1}^{\infty} (A \setminus (\bigcup_{|j| \leq J(i)} C_{ij}))$  this implies  $\mu(A \setminus C) < \varepsilon$ , where  $C = \bigcap_{i=1}^{\infty} (\bigcup_{|j| \leq J(i)} C_{ij})$ , which is a closed set in  $X$ .

Finally, define  $g_i : \bigcup_{|j| \leq J(i)} C_{ij} \rightarrow \mathbb{R}$  by setting  $g_i(x) = \frac{j-1}{i}$  on  $C_{ij}, |j| \leq J(i)$ . Then, since the  $C_{i1}, \dots, C_{iJ(i)}$  are pairwise disjoint closed sets,  $g_i$  is clearly continuous and its restriction to  $C$  is continuous for each  $i$ . Furthermore by construction  $0 \leq f(x) - g_i(x) \leq 1/i$  for each  $x \in C$  and each  $i = 1, 2, \dots$ , so  $g_i|_C$  converges uniformly to  $f|_C$  on  $C$ , and hence  $f|_C$  is continuous.  $\square$

## 2 Hausdorff Measure

In this section we suppose  $X$  is a metric space with metric  $d$ , and we let

$$\omega_m = \frac{\pi^{m/2}}{\int_0^{\infty} t^{m/2} e^{-t} dt}, \quad m \geq 0,$$

so that in particular  $\omega_m$  is the volume (Lebesgue measure) of the unit ball  $B_1^m(0)$  in  $\mathbb{R}^m$  in case  $m$  happens to be a positive integer.

For any  $m \geq 0$  we define the  $m$ -dimensional Hausdorff (outer) measure

$$2.1 \quad \mathcal{H}^m(A) = \lim_{\delta \downarrow 0} \mathcal{H}_\delta^m(A), \quad A \subset X,$$

where, for each  $\delta > 0$ ,  $\mathcal{H}_\delta^m(A)$  (called the “size  $\delta$  approximation to  $\mathcal{H}^m$ ”) is defined by taking  $\mathcal{H}_\delta^m(\emptyset) = 0$  and, for any non-empty  $A \subset X$ ,

$$2.2 \quad \mathcal{H}_\delta^m(A) = \omega_m \inf \sum_{j=1}^{\infty} \left( \frac{\text{diam } C_j}{2} \right)^m,$$

where the inf is taken over all countable collections  $C_1, C_2, \dots$  of subsets of  $X$  such that  $\text{diam } C_j < \delta$  and  $A \subset \bigcup_{j=1}^{\infty} C_j$ ; the right side is to be interpreted as  $\infty$  in case there is no such collection  $C_1, C_2, \dots$ . (Of course in a separable metric space  $X$  there are always such collections  $C_1, C_2, \dots$  for each  $\delta > 0$ .) The limit in 2.1 always exists (although it may be  $+\infty$ ) because  $\mathcal{H}_\delta^m(A)$  is a decreasing function of  $\delta$ ; thus  $\mathcal{H}^m(A) = \sup_{\delta > 0} \mathcal{H}_\delta^m(A)$  for each  $m \geq 0$ . It is left as an exercise to check that  $\mathcal{H}_\delta^m$  and  $\mathcal{H}^m$  are indeed outer measures on  $X$ .

Notice also that  $\mathcal{H}^0$  is just “counting measure”:  $\mathcal{H}^0(\emptyset) = 0$ ,  $\mathcal{H}^0(A) =$  the number of elements in the set  $A$  if  $A$  is finite, and  $\mathcal{H}^0(A) = \infty$  if  $A$  is not finite.

**2.3 Remarks:** (1) Since  $\text{diam } C_j = \text{diam } \bar{C}_j$  we can add the additional requirement in the identity 2.2 that the  $C_j$  be *closed* without changing the value of  $\mathcal{H}^m(A)$ ; indeed since for any  $\varepsilon > 0$  we can find an open set  $U_j \supset C_j$  with  $\text{diam } U_j < \text{diam } C_j + \varepsilon/2^j$ , we could also take the  $C_j$  to be *open*.

(2) Evidently  $\mathcal{H}_\delta^m(A) < \infty \forall m \geq 0, \delta > 0$  in case  $A$  is a totally bounded subset of the metric space  $X$ .

One easily checks from the definition of  $\mathcal{H}_\delta^m$  that

$$\mathcal{H}_\delta^m(A \cup B) = \mathcal{H}_\delta^m(A) + \mathcal{H}_\delta^m(B) \text{ whenever } d(A, B) > \delta,$$

hence

$$2.4 \quad \mathcal{H}^m(A \cup B) = \mathcal{H}^m(A) + \mathcal{H}^m(B) \quad \text{whenever } d(A, B) > 0,$$

where  $d(A, B) = \inf_{x \in A, y \in B} d(x, y)$ , and therefore *all Borel sets are  $\mathcal{H}^m$ -measurable* by the Caratheodory Criterion (Theorem 1.15). It then follows from Remark 2.3(1) (see problem 1.2 of Ch.1 problems) that

$$2.5 \quad \mathcal{H}^m \text{ is Borel-regular for each } m \geq 0.$$

**Note:** It is *not* true in general that the Borel sets are  $\mathcal{H}_\delta^m$ -measurable for  $\delta > 0$ ; for instance if  $n = 2$  then one easily checks that the half-space  $H = \{x = (x^1, x^2) \in \mathbb{R}^n : x^2 > 0\}$  is not  $\mathcal{H}_\delta^1$ -measurable, because for example it does not cut the set  $S_\varepsilon = ([0, 1] \times \{0\}) \cup ([0, 1] \times \{\varepsilon\})$  additively for sufficiently small  $\varepsilon$ . Indeed one can directly use the definition of  $\mathcal{H}_\delta^1$  to check that  $\mathcal{H}_\delta^1(S_\varepsilon) \downarrow 1$  as  $\varepsilon \downarrow 0$  (and in particular  $\mathcal{H}_\delta^1(S_\varepsilon) < 2$  for sufficiently small  $\varepsilon$ ), whereas  $\mathcal{H}_\delta^1(S_\varepsilon \cap H) = \mathcal{H}_\delta^1(S_\varepsilon \setminus H) = 1$  for each  $\varepsilon > 0$ .

We will later show that, for each integer  $n \geq 1$ ,  $\mathcal{H}^n$  agrees with the usual definition of  $n$ -dimensional volume measure on an  $n$ -dimensional  $C^1$  submanifold of  $\mathbb{R}^{n+k}$ ,  $k \geq 0$ . As a first step we want to prove that  $\mathcal{H}^n$  and  $\mathcal{L}^n$  ( $n$ -dimensional Lebesgue measure) agree on  $\mathbb{R}^n$ .

We claim that, on  $\mathbb{R}^n$ , the outer measures  $\mathcal{L}^n, \mathcal{H}^n, \mathcal{H}_\delta^n$  all coincide (for each  $\delta > 0$ ):

## 2.6 Theorem.

$$\mathcal{L}^n(A) = \mathcal{H}^n(A) = \mathcal{H}_\delta^n(A) \text{ for every } A \subset \mathbb{R}^n \text{ and every } \delta > 0.$$

**Proof:** We first show

$$(1) \quad \mathcal{H}_\delta^n(A) \leq \mathcal{L}^n(A) \quad \forall \delta > 0$$

as follows: Let  $I_i$ ,  $i = 1, 2, \dots$ , be any open intervals with  $A \subset \cup_i I_i$ . By problem 1.5 of Ch.1 problems, for each  $\delta > 0$  and each  $i = 1, 2, \dots$  we can choose closed balls  $B_{\rho_j}(x_j)$ ,  $j = 1, 2, \dots$  as in 1.18 with  $U = I_i$  and with  $\rho_j < \delta$  for each  $j$ . Since  $\mathcal{L}^n(Z) = 0 \Rightarrow \mathcal{H}_\delta^n(Z) = 0$  for each subset  $Z \subset X$  (by Definitions 2.2, 1.16) we then have

$$(2) \quad \begin{aligned} \mathcal{H}_\delta^n(I_i) &= \mathcal{H}_\delta^n(\cup_{j=1}^\infty B_{\rho_j}(x_j)) \leq \sum_{j=1}^\infty \omega_n \rho_j^n \\ &= \sum_{j=1}^\infty \mathcal{L}^n(B_{\rho_j}(x_j)) = \mathcal{L}^n(\cup_{j=1}^\infty B_{\rho_j}(x_j)) = \mathcal{L}^n(I_i) = |I_i|, \end{aligned}$$

and hence

$$(3) \quad \mathcal{H}_\delta^n(A) \leq \mathcal{H}_\delta^n(\cup_i I_i) \leq \sum_i \mathcal{H}_\delta^n(I_i) \leq \sum_i |I_i|.$$

The proof of (1) is then completed by taking inf over all such collections  $\{I_i\}$  and using Definition 1.16.

To prove the reverse inequality we first need a geometric result concerning Lebesgue measure, known as *the isodiametric inequality*:

## 2.7 Theorem (Isodiametric Inequality.)

$$\mathcal{L}^n(A) \leq \omega_n \left( \frac{\text{diam } A}{2} \right)^n \text{ for every set } A \subset \mathbb{R}^n.$$

**Remark:** Thus among all sets  $A \subset \mathbb{R}^n$  with a given diameter  $\rho$ , the ball with diameter  $\rho$  has the largest  $\mathcal{L}^n$  measure.

**Proof of 2.7:** Observe that it suffices to prove this for compact sets because  $\mathcal{L}^n(A) \leq \mathcal{L}^n(\bar{A})$ , while  $\bar{A}$  has the same diameter as  $A$  and the isodiametric inequality is trivial if  $\text{diam } A = \infty$ . For a compact set  $A$  we proceed to use *Steiner symmetrization*: The Steiner symmetrization  $S_j(A)$  of the compact set  $A$  with respect to the  $j$ -th coordinate plane  $x^j = 0$  is defined as follows: For  $\xi$  in the coordinate plane  $x^j = 0$  let  $\ell_j(\xi) = \{\xi + te_j : t \in \mathbb{R}\}$  and let  $\pi$  be the projection  $\xi + te_j \mapsto t$  of the line  $\ell_j(\xi)$  onto the real line  $\mathbb{R}$ , and let  $\sigma_j(A, \xi)$  denote the closed line segment  $\{\xi + te_j : |t| \leq \frac{1}{2} \mathcal{L}^1(\pi(A \cap \ell_j(\xi)))\}$ . Then

$$S_j(A) = \cup_{\{\xi: A \cap \ell_j(\xi) \neq \emptyset\}} \sigma_j(A, \xi).$$

(Thus  $S_j(A)$  is obtained by replacing  $A \cap \ell_j(\xi)$  with the segment  $\sigma_j(A, \xi)$  for each  $\xi$  such that  $A \cap \ell_j(\xi) \neq \emptyset$ .) This process gives a new compact set  $S_j(A)$  with diameter not larger than the diameter of the original set  $A$  (see Ch.1 problem 1.1) and, by Fubini, the same Lebesgue measure. Further if  $i \neq j$  and if  $A$  is already invariant under reflection in the  $i$ -th coordinate plane  $x^i = 0$ , then by definition  $S_j(A)$  is invariant under reflection in both the  $i$ -th and the  $j$ -th coordinate planes. Thus by applying Steiner symmetrization successively with respect to coordinate planes  $x^1 = 0, x^2 = 0, \dots, x^n = 0$ , we get a new

compact set  $\tilde{A}$  with diameter  $\leq \text{diam } A$ , having the same Lebesgue measure as  $A$ , and being invariant with respect to the transformation  $x \mapsto -x$ . This means in particular that  $2|x| = |x - (-x)| \leq \text{diam } \tilde{A} \leq \text{diam } A$  for each  $x \in \tilde{A}$ , so  $\tilde{A}$  is contained in the closed ball with radius  $\frac{1}{2} \text{diam } A$  and center 0, whence

$$\mathcal{L}^n(A) = \mathcal{L}^n(\tilde{A}) \leq \omega_n \left(\frac{1}{2} \text{diam } A\right)^n$$

as required.  $\square$

**Completion of the proof of 2.6:** We have to prove

$$(\ddagger) \quad \mathcal{L}^n(A) \leq \mathcal{H}_\delta^n(A) \quad \forall \delta > 0, A \subset \mathbb{R}^n.$$

Suppose  $\delta > 0$ ,  $A \subset \mathbb{R}^n$ , and let  $C_1, C_2, \dots$  be any countable collection with  $A \subset \cup_j C_j$  and  $\text{diam } C_j < \delta$ . Then

$$\begin{aligned} \mathcal{L}^n(A) &\leq \mathcal{L}^n(\cup_j C_j) \leq \sum_j \mathcal{L}^n(C_j) \\ &\leq \sum_j \omega_n \left(\frac{1}{2} \text{diam } C_j\right)^n \text{ by 2.7.} \end{aligned}$$

Taking the inf over all such collections  $\{C_j\}$  we have  $(\ddagger)$  as required.  $\square$

### 3 Densities

Throughout this section  $X$  will denote a metric space with metric  $d$ . We first we want to introduce the notion of  $n$ -dimensional density of a measure  $\mu$  on  $X$ , where  $X$  continues to denote a metric space with metric  $d$ . For any outer measure  $\mu$  on  $X$ , any subset  $A \subset X$ , and any point  $x \in X$ , we define the  $n$ -dimensional upper and lower densities  $\Theta^{*n}(\mu, A, x)$ ,  $\Theta_*^n(\mu, A, x)$  by

$$\begin{aligned} 3.1 \quad \Theta^{*n}(\mu, A, x) &= \limsup_{\rho \downarrow 0} \frac{\mu(A \cap B_\rho(x))}{\omega_n \rho^n} \\ \Theta_*^n(\mu, A, x) &= \liminf_{\rho \downarrow 0} \frac{\mu(A \cap B_\rho(x))}{\omega_n \rho^n}. \end{aligned}$$

In case  $A = X$  we simply write  $\Theta^{*n}(\mu, x)$  and  $\Theta_*^n(\mu, x)$  to denote these quantities so that  $\Theta^{*n}(\mu, A, x) = \Theta^{*n}(\mu \llcorner A, x)$ ,  $\Theta_*^n(\mu, A, x) = \Theta_*^n(\mu \llcorner A, x)$ .

**3.2 Remark:** If all Borel sets are  $\mu$ -measurable and if  $\mu(B_\rho(x))$  is finite on each ball  $B_\rho(x) \subset X$ , then  $\mu(A \cap B_\rho(x)) \geq \limsup_{y \rightarrow x} \mu(A \cap B_\rho(y))$  for each fixed  $\rho > 0$  (i.e.  $\mu(A \cap B_\rho(x))$  is an upper semi-continuous function of  $x$  for each fixed  $\rho > 0$ ). Indeed if  $x_k \rightarrow x$  and  $j \in \{1, 2, \dots\}$  then  $B_{\rho+1/j}(x) \supset B_\rho(x_k)$  for sufficiently large

$k$ , hence  $\mu(B_{\rho+1/j}(x)) \geq \limsup_{y \rightarrow x} \mu(B_\rho(y))$  for each  $j = 1, 2, \dots$ , and  $B_\rho(x) = \bigcap_{j=1}^{\infty} B_{\rho+1/j}(x)$  (because  $B_\rho(x)$  is the *closed* ball of radius  $\rho$ ), hence, by 1.10,  $\mu(B_\rho(x)) = \lim_{j \rightarrow \infty} \mu(B_{\rho+1/j}(x)) \geq \limsup_{k \rightarrow \infty} \mu(B_\rho(x_k))$ , which is the claimed upper semi-continuity of  $\mu(B_\rho(x))$ . Hence  $\inf_{0 < \rho < \delta} (\omega_n \rho^n)^{-1} \mu(A \cap B_\rho(x))$  is also upper semi-continuous and hence Borel measurable (because the inf of a family of upper semi-continuous functions is again upper semi-continuous), and so

$$\begin{aligned} \Theta_*^n(\mu, A, x) &= \lim_{\delta \downarrow 0} \inf_{0 < \rho < \delta} (\omega_n \rho^n)^{-1} \mu(A \cap B_\rho(x)) \\ &= \lim_{j \rightarrow \infty} \inf_{0 < \rho < 1/j} (\omega_n \rho^n)^{-1} \mu(A \cap B_\rho(x)) \end{aligned}$$

is also Borel measurable. Similarly since  $\mu(A \cap \check{B}_\rho(x))$  is lower semi-continuous (where  $\check{B}_\rho(x)$  denotes the open ball of radius  $\rho$  and center  $x$ ) and evidently

$$\sup_{0 < \rho < \delta} (\omega_n \rho^n)^{-1} \mu(A \cap \check{B}_\rho(x)) = \sup_{0 < \rho < \delta} (\omega_n \rho^n)^{-1} \mu(A \cap B_\rho(x))$$

(and hence it makes no difference whether we use open or closed balls in the definition of lower density, nor in the definition of upper density for that matter), we see that  $\Theta^{*n}(\mu, A, x)$  can be written  $\lim_{j \rightarrow \infty} \sup_{0 < \rho < 1/j} (\omega_n \rho^n)^{-1} \mu(A \cap \check{B}_\rho(x))$ , so we also have  $\Theta^{*n}(\mu, A, x)$  is Borel measurable. Notice that  $A$  need not be  $\mu$ -measurable here.

Subsequently we use the notation that if  $\Theta^{*n}(\mu, A, x) = \Theta_*^n(\mu, A, x)$  then the common value will be denoted  $\Theta^n(\mu, A, x)$ .

Appropriate information about the upper density gives connections between  $\mu$  and  $\mathcal{H}^n$ . Specifically, we have the following comparison theorem:

**3.3 Theorem.** *Let  $\mu$  be any outer measure on the metric space  $X$  such that all Borel sets are measurable (e.g.  $\mu$  is Borel regular),  $t \geq 0$ , and  $A_1 \subset A_2 \subset X$ . Then*

$$\Theta^{*n}(\mu, A_2, x) \geq t \quad \forall x \in A_1 \Rightarrow t \mathcal{H}^n(A_1) \leq \mu(A_2),$$

An important special case of this theorem is the case  $A_1 = A_2$ . Notice that we do *not* need to assume  $A_1, A_2$  are  $\mu$ -measurable here.

The proof of 3.3 will make use of the following important “5-times covering lemma,” in which we use the notation that if  $B$  is a ball  $B_\rho(x) \subset X$ , then  $\widehat{B} = B_{5\rho}(x)$ .

**3.4 Lemma (5-times Covering Lemma).** *If  $B$  is any family of closed balls in  $X$  with  $R \equiv \sup\{\text{diam } B : B \in B\} < \infty$ , then there is a pairwise disjoint subcollection  $B' \subset B$  such that*

$$\bigcup_{B \in B} B \subset \bigcup_{B \in B'} \widehat{B};$$

in fact we can arrange the stronger property

$$(\ddagger) \quad B \in \mathcal{B} \Rightarrow \exists B' \in \mathcal{B}' \text{ with } B' \cap B \neq \emptyset \text{ and} \\ \text{diam}(B') \geq \frac{1}{2} \text{diam}(B), \text{ hence } \widehat{B'} \supset B.$$

**Proof:** For  $j = 1, 2, \dots$  let  $\mathcal{B}_j = \{B \in \mathcal{B} : R/2^j < \text{diam } B \leq R/2^{j-1}\}$ , so that  $\mathcal{B} = \bigcup_{j=1}^{\infty} \mathcal{B}_j$ , and this is a pairwise disjoint union. Proceed to define  $\mathcal{B}'_j \subset \mathcal{B}_j$  as follows:

(i) Let  $\mathcal{B}'_1$  be any maximal pairwise disjoint subcollection of  $\mathcal{B}_1$ . Such  $\mathcal{B}'_1$  exist by applying Zorn's lemma to  $\mathcal{P} = \{\mathcal{A} : \mathcal{A} \text{ is a pairwise disjoint subcollection of } \mathcal{B}_1\}$ , which is partially ordered by inclusion; notice for any totally ordered subcollection  $\mathcal{T} \subset \mathcal{P}$  we clearly have  $\mathcal{A} \cup \bigcup_{B \in \mathcal{T}} B \in \mathcal{P}$  for each  $\mathcal{A} \in \mathcal{T}$ , so Zorn's lemma is indeed applicable. Notice also that in a general metric space the collection  $\mathcal{B}'_1$  could be uncountable, but of course in a separable metric space (i.e. a metric space with a countable dense subset) all pairwise disjoint collections of balls must be countable.

(ii) Assuming  $j \geq 2$  and that  $\mathcal{B}'_1 \subset \mathcal{B}_1, \dots, \mathcal{B}'_{j-1} \subset \mathcal{B}_{j-1}$  are defined, let  $\mathcal{B}'_j$  be a maximal pairwise disjoint subcollection of  $\{B \in \mathcal{B}_j : B \cap B' = \emptyset \text{ whenever } B' \in \bigcup_{i=1}^{j-1} \mathcal{B}'_i\}$ . Again, Zorn's lemma guarantees such a maximal collection exists.

Now if  $j \geq 1$  and  $B \in \mathcal{B}_j$  we must have

$$B \cap B' \neq \emptyset \text{ for some } B' \in \bigcup_{i=1}^j \mathcal{B}'_i$$

(otherwise we contradict maximality of  $\mathcal{B}'_j$ ), and for such a pair  $B, B'$  we have  $\text{diam } B \leq R/2^{j-1} = 2R/2^j \leq 2 \text{diam } B'$ , so that  $(\ddagger)$  holds; in particular  $B \subset \widehat{B'}$ .  $\square$

**3.5 Remark:** The factor “5” in the above lemma can be improved; indeed by defining  $\mathcal{B}_k = \{B \in \mathcal{B} : R/(1 + \theta)^k < \text{diam } B \leq R/(1 + \theta)^{k-1}\}$  with  $\theta$  small enough, the same argument as that used in the above proof establishes a “ $(3 + \varepsilon)$ -times covering lemma” for any  $\varepsilon > 0$ . However there is no such “3-times covering lemma,” as one sees by taking  $\mathcal{B} = \{B_\rho(-\rho e_1) : \rho < 1\} \cup \{B_\rho(\rho e_1) : \rho < 1\}$ . Then  $\bigcup_{B \in \mathcal{B}} B = \{0\} \cup \check{B}_1(-e_1) \cup \check{B}_1(e_1)$ , whereas, since all the balls in  $\mathcal{B}$  contain 0, a pairwise disjoint subcollection of  $\mathcal{B}$  must consist just of a single ball  $B = B_\rho(\pm \rho e_1)$  for some  $\rho < 1$ , and  $B_R(\rho e_1) \supset \{0\} \cup \check{B}_1(-e_1) \cup \check{B}_1(e_1)$  only if  $R \geq \rho + 2 = 3\rho + 2(1 - \rho) (> 3\rho)$ .

**3.6 Definition:** In the following corollary of 3.4 we use the terminology that a subset  $A \subset X$  is *covered finely* by a collection  $\mathcal{B}$  of balls, meaning that

$$\inf\{\text{diam } B : x \in B \in \mathcal{B}\} = 0 \quad \forall x \in A.$$

**3.7 Corollary.**  $A \subset X$  is covered finely (as in Definition 3.6) by a collection  $\mathcal{B}$  of closed balls, then there is a pairwise disjoint subcollection  $\mathcal{B}' \subset \mathcal{B}$  such that

$$A \setminus \bigcup_{j=1}^N B_j \subset \bigcup_{B \in \mathcal{B} \setminus \{B_1, \dots, B_N\}} \widehat{B}$$

for each finite subcollection  $\{B_1, \dots, B_N\} \subset \mathcal{B}'$ .

**Proof:** Without loss of generality we can assume  $\text{diam } B \leq 1$  for each  $B$  in  $\mathcal{B}$ , because  $\{B \in \mathcal{B} : \text{diam } B \leq 1\}$  clearly still covers  $A$  finely.

Then we can apply the 5-times covering lemma 3.4 to give a pairwise disjoint collection  $\mathcal{B}' \subset \mathcal{B}$  such that 3.4(‡) holds. For any  $B_1, \dots, B_N \in \mathcal{B}'$  take any  $x \in A \setminus \bigcup_{j=1}^N B_j$ , and, since  $X \setminus \bigcup_{j=1}^N B_j$  is open and  $\mathcal{B}$  covers  $A$  finely, we can then find  $B \in \mathcal{B}$  with  $B \cap (\bigcup_{j=1}^N B_j) = \emptyset$  and  $x \in B$ . By 3.4(‡) there is a  $B' \in \mathcal{B}'$  with  $B' \cap B \neq \emptyset$  and  $\widehat{B'} \supset B$ . Evidently  $B' \neq B_j \ \forall j = 1, \dots, N$ , so  $x \in \bigcup_{B' \in \mathcal{B}' \setminus \{B_1, \dots, B_N\}} \widehat{B'}$ .  $\square$

**Proof of 3.3:** We can assume  $\mu(A_2) < \infty$  and  $t > 0$  otherwise the result is trivial. Take  $\tau \in (0, t)$ , so that then

$$\Theta^{*n}(\mu, A_2, x) > \tau \text{ for } x \in A_1.$$

For  $\delta > 0$ , let  $\mathcal{B}$  (depending on  $\delta$ ) be defined by

$$\mathcal{B} = \{\text{closed balls } B_\rho(x) : x \in A_1, 0 < \rho < \delta/2, \mu(A_2 \cap B_\rho(x)) \geq \tau \omega_n \rho^n\}.$$

Evidently  $\mathcal{B}$  covers  $A_1$  finely and hence there is a pairwise disjoint subcollection  $\mathcal{B}' \subset \mathcal{B}$  so that 3.4(‡) holds. Since  $\mu(A_2 \cap B) > 0$  for each  $B \in \mathcal{B}$  and since  $B_1, \dots, B_N \in \mathcal{B}' \Rightarrow \sum_{j=1}^N \mu(A_2 \cap B_j) = \mu(A_2 \cap (\bigcup_{j=1}^N B_j)) \leq \mu(A_2) < \infty$  it follows that  $\mathcal{B}'$  is a countable collection  $\{B_{\rho_1}(x_1), B_{\rho_2}(x_2), \dots\}$  and hence 3.7 implies

$$A_1 \setminus \bigcup_{j=1}^N B_{\rho_j}(x_j) \subset \bigcup_{j=N+1}^\infty B_{5\rho_j}(x_j) \quad \forall N \geq 1,$$

and also, by definition of  $\mathcal{B}$ ,

$$\tau \sum_{j=1}^\infty \omega_n \rho_j^n \leq \sum_{j=1}^\infty \mu(A_2 \cap B_{\rho_j}(x_j)) = \mu(A_2 \cap (\bigcup_{j=1}^\infty B_{\rho_j}(x_j))) \leq \mu(A_2) < \infty.$$

Since  $A_1 \subset (\bigcup_{j=1}^N B_{\rho_j}(x_j)) \cup (\bigcup_{j=N+1}^\infty B_{5\rho_j}(x_j))$ , we have

$$\mathcal{H}_{5\delta}^n(A_1) \leq \sum_{j=1}^N \omega_n \rho_j^n + 5^n \sum_{j=N+1}^\infty \omega_n \rho_j^n$$

by Definition 2.2, and hence letting  $N \rightarrow \infty$  we deduce

$$\tau \mathcal{H}_{5\delta}^n(A_1) \leq \mu(A_2).$$

The required result now follows by letting  $\delta \downarrow 0$  and  $\tau \uparrow t$ .  $\square$

As a corollary to 3.3 we can prove the following “Upper Density Theorem.”

**3.8 Theorem (Upper Density Theorem.)** *If  $\mu$  is an outer measure on  $X$  such that all Borel sets are measurable (e.g.  $\mu$  is Borel regular) and if  $A$  is a  $\mu$ -measurable subset of  $X$  with  $\mu(A) < \infty$ , then*

$$\Theta^{*n}(\mu, A, x) = 0 \text{ for } \mathcal{H}^n\text{-a.e. } x \in X \setminus A.$$

**3.9 Remarks:** (1) Of course  $\mu = \mathcal{H}^n$  is an important special case.

(2) If  $\mu$  is open  $\sigma$ -finite (i.e.  $X = \bigcup_{j=1}^{\infty} V_j$  with  $V_j$  open and  $\mu(V_j) < \infty$  for each  $j = 1, 2, \dots$ ), then one can drop the hypothesis that  $\mu(A) < \infty$ , because we can apply the theorem with  $\mu \llcorner V_j$  in place of  $\mu$  to conclude that

$$\Theta^{*n}(\mu, A, x) = \Theta^{*n}(\mu, A \cap V_j, x) = 0 \text{ for } \mathcal{H}^n\text{-a.e. } x \in V_j \setminus A, \quad j = 1, 2, \dots,$$

and hence  $\Theta^{*n}(\mu, A, x) = 0$  for  $\mathcal{H}^n$ -a.e.  $x \in X \setminus A$ .

**Proof of 3.8:** Let  $C$  be any closed subset of  $A$ ,  $t > 0$  and  $S_t = \{x \in X \setminus A : \Theta^{*n}(\mu, A, x) \geq t\}$ . Since  $X \setminus C$  is open and  $S_t \subset X \setminus A \subset X \setminus C$  we have  $\Theta^{*n}(\mu, A \cap (X \setminus C), x) = \Theta^{*n}(\mu, A, x) \geq t$  for  $x \in S_t$ . Thus we can apply 3.3 with  $\mu \llcorner A$ ,  $S_t$ ,  $X \setminus C$  in place of  $\mu$ ,  $A_1$ ,  $A_2$  to give  $t\mathcal{H}^n(S_t) \leq \mu(A \setminus C)$  for each closed set  $C \subset A$ . But  $\inf_{C \text{ closed}, C \subset A} \mu(A \setminus C) = 0$  by 1.22(2), so  $\mathcal{H}^n(S_t) = 0$ . Taking  $t = 1/i$ ,  $i = 1, 2, \dots$ , we thus conclude  $\mathcal{H}^n(\{x \in X \setminus A : \Theta^{*n}(\mu, A, x) > 0\}) = 0$ .  $\square$

Notice that we have the following important corollary to the above theorem:

**3.10 Corollary.** *If  $A \subset \mathbb{R}^n$  is  $\mathcal{L}^n$ -measurable then the density  $\Theta^n(\mathcal{L}^n, A, x)$  exists  $\mathcal{L}^n$ -a.e.  $x \in \mathbb{R}^n$ , and  $\Theta^n(\mathcal{L}^n, A, x) = 0$   $\mathcal{L}^n$ -a.e.  $x \in \mathbb{R}^n \setminus A$  and  $= 1$   $\mathcal{L}^n$ -a.e.  $x \in A$ .*

**Proof:** Indeed  $(\omega_n \rho^n)^{-1} \mathcal{L}^n(A \cap B_\rho(x)) + (\omega_n \rho^n)^{-1} \mathcal{L}^n(B_\rho(x) \setminus A) = 1$  for each  $\rho > 0$ , and, by the Upper Density Theorem 3.8, the first term on the left  $\rightarrow 0$  as  $\rho \downarrow 0$  for  $\mathcal{L}^n$ -a.e.  $x \in \mathbb{R}^n \setminus A$  while the second term on the left  $\rightarrow 0$  as  $\rho \downarrow 0$  for  $\mathcal{L}^n$ -a.e.  $x \in A$ .  $\square$

We conclude this section with two important bounds for densities of Hausdorff measure.

**3.11 Theorem.** *For any  $\mathcal{H}^n$ -measurable subset of  $A$  of a metric space  $X$ :*

- (1) *If  $\mathcal{H}^n(A) < \infty$ , then  $\Theta^{*n}(\mathcal{H}^n, A, x) \leq 1$  for  $\mathcal{H}^n$ -a.e.  $x \in A$ .*
- (2) *If  $\mathcal{H}_\delta^n(A) < \infty$  for each  $\delta > 0$  (note this is automatic if  $A$  is a totally bounded subset of  $X$ ), then  $\Theta^{*n}(\mathcal{H}_\infty^n, A, x) \geq 2^{-n}$  for  $\mathcal{H}^n$ -a.e.  $x \in A$ .*

**3.12 Remark:** Since  $\mathcal{H}^n \geq \mathcal{H}_\delta^n \geq \mathcal{H}_\infty^n$  (by Definitions 2.1, 2.2) this theorem implies

$$2^{-n} \leq \Theta^{*n}(\mathcal{H}^n, A, x) \leq 1 \text{ for } \mathcal{H}^n\text{-a.e. } x \in A,$$

provided  $A$  is  $\mathcal{H}^n$ -measurable and  $\mathcal{H}^n(A) < \infty$ .

**Proof of 3.11:** To prove (1), let  $\varepsilon, t > 0$ , let  $A_t = \{x \in A : \Theta^{*n}(\mathcal{H}^n, A, x) \geq t\}$  and (using 1.22(1) with  $\mu = \mathcal{H}^n \llcorner A$ ), choose an open set  $U \supset A_t$  such that

$$\mathcal{H}^n(U \cap A) < \mathcal{H}^n(A_t) + \varepsilon.$$

Since  $U$  is open and since  $A_t \subset U$  we have  $\Theta^{*n}(\mathcal{H}^n, A \cap U, x) \geq t$  for each  $x \in A_t$ . Hence 3.3 (with  $\mathcal{H}^n \llcorner A, A_t, A \cap U$  in place of  $\mu, A_1, A_2$ ) implies that

$$t\mathcal{H}^n(A_t) \leq \mathcal{H}^n(A \cap U) \leq \mathcal{H}^n(A_t) + \varepsilon.$$

We thus have  $\mathcal{H}^n(A_t) = 0$  for each  $t > 1$ . Since  $\{x : \Theta^{*n}(\mathcal{H}^n, A, x) > 1\} = \bigcup_{j=1}^{\infty} A_{t_j}$  for any decreasing sequence  $\{t_j\}$  with  $\lim t_j = 1$ , we thus have  $\mathcal{H}^n\{x : \Theta^{*n}(\mathcal{H}^n, A, x) > 1\} = 0$ , as required.

To prove (2), suppose for contradiction that  $\Theta^{*n}(\mathcal{H}^n_{\infty} \llcorner A, x) < 2^{-n}$  for each  $x$  in a set  $B_0 \subset A$  with  $\mathcal{H}^n(B_0) > 0$ . Then for each  $x \in B_0$  select  $\delta_x \in (0, 1)$  such that

$$\mathcal{H}^n_{\infty}(A \cap B_{\rho}(x)) \leq \frac{1 - \delta_x}{2^n} \omega_n \rho^n, \quad 0 < \rho < \delta_x.$$

Therefore, since  $B_0 = \bigcup_{j=1}^{\infty} \{x \in B_0 : \delta_x > 1/j\}$  and since  $\mathcal{H}^n_{\delta}(A \cap B_{\rho}(x)) \equiv \mathcal{H}^n_{\infty}(A \cap B_{\rho}(x))$  for any  $\rho < \delta/2$  (by Definition 2.2), we can select  $\delta > 0$  and  $B \subset B_0$  with  $\mathcal{H}^n(B) > 0$  and

$$(1) \quad \mathcal{H}^n_{\delta}(A \cap B_{\rho}(x)) \leq \frac{1 - \delta}{2^n} \omega_n \rho^n, \quad 0 < \rho < \delta/2, \quad x \in B.$$

Now using Definition 2.2 again, we can choose sets  $C_1, C_2, \dots$  with  $B \subset \bigcup_{j=1}^{\infty} C_j$ ,  $C_j \cap B \neq \emptyset$ ,  $\text{diam } C_j < \delta \forall j$ , and

$$(2) \quad \sum_j \omega_n \rho_j^n < \frac{1}{1 - \delta} \mathcal{H}^n_{\delta}(B), \quad \rho_j = \text{diam } C_j / 2$$

Now take  $x_j \in C_j \cap B$ , so that  $B \subset A \cap (\bigcup_{j=1}^{\infty} B_{2\rho_j}(x_j))$ , and we conclude from (1), (2) that  $\mathcal{H}^n_{\delta}(B) = 0$ , hence  $\mathcal{H}^n(B) = 0$ , contradicting our choice of  $B$ .  $\square$

## 4 Differentiation Theorems

We begin with discussion of the possibility of extending the Comparison and Upper Density Theorems 3.3, 3.8 to the situation when, in a metric space  $X$ , we consider the upper density of a Borel regular measure  $\mu$  with respect to another Borel regular measure  $\mu_0$ . In this case we always assume  $\mu_0$  is locally finite, so that

$$4.1 \quad \forall x \in X \text{ there is } \rho > 0 \text{ with } \mu_0(B_{\rho}(x)) < \infty.$$

We note that this is automatic if  $\mu_0$  is open  $\sigma$ -finite as in 1.21.

The *upper density*  $\Theta^{*\mu_0}(\mu, x)$  of  $\mu$  with respect to  $\mu_0$  at the point  $x \in X$  is defined by

$$4.2 \quad \Theta^{*\mu_0}(\mu, x) = \begin{cases} \limsup_{\rho \downarrow 0} \frac{\mu(B_\rho(x))}{\mu_0(B_\rho(x))} & \text{for } x \in X \setminus (U_0 \cup V_0) \\ \infty & \text{for } x \in U_0 \setminus V_0 \\ 0 & \text{for } x \in V_0, \end{cases}$$

where  $U_0$  is the open set consisting of all points  $x \in X$  with  $\mu_0(B_\rho(x)) = 0$  for some  $\rho > 0$  and  $V_0$  is the open set consisting of all points  $x \in X$  with  $\mu(B_\rho(x)) = 0$  for some  $\rho > 0$ . Notice  $\Theta^{*\mu_0}(\mu, x) = \Theta^{*\mu}(\mu, x)$  in the special case when  $X = \mathbb{R}^n$  and  $\mu_0 = \mathcal{L}^n$ .

To prove a useful analogue to the Upper Density Theorem 3.8 in this situation we need to assume that  $\mu_0$  has the “Symmetric Vitali” property according to the following definition:

**4.3 Definition (Symmetric Vitali Property):** An outer measure  $\mu_0$  on a metric space  $X$  is said to have the Symmetric Vitali Property if given any  $A \subset X$  with  $\mu_0(A) < \infty$  and any collection  $\mathcal{B}$  of closed balls with centers in  $A$  which cover  $A$  finely (i.e.  $\inf\{\rho : B_\rho(x) \in \mathcal{B}\} = 0$  for each  $x \in A$ ),  $\exists$  a countable pairwise disjoint collection  $\mathcal{B}' = \{B_{\rho_j}(x_j) : j = 1, 2, \dots\} \subset \mathcal{B}$  with  $\mu_0(A \setminus (\cup_{j=1}^\infty B_{\rho_j}(x_j))) = 0$ .

Before proceeding, we make some important notes concerning the open set  $U_0$  in 4.2 and the Symmetric Vitali Property:

**4.4 Remarks:** (1) First note that there are various scenarios which guarantee that the open set  $U_0$  in the definition 4.2 of the density  $\Theta^{*\mu_0}(\mu, x)$  has  $\mu_0$ -measure zero. For example if  $X$  is separable (i.e.  $X$  has a countable dense subset) then  $U_0$  can be written as a countable union of balls  $B_\rho(x) \subset U_0$  with  $\mu(B_\rho(x)) = 0$ , and hence  $U_0$  certainly has  $\mu_0$ -measure zero in this case. Also, if  $\mu_0$  is  $\sigma$ -finite and has the Symmetric Vitali Property, then, because  $U_0$  is trivially covered finely by the collection  $\mathcal{B}$  of balls  $B_\rho(x) \subset U_0$  with  $\mu_0(B_\rho(x)) = 0$ , there is a countable subcollection of  $\mathcal{B}$  covering  $\mu_0$ -almost all of  $U_0$ , so again  $\mu_0(U_0) = 0$ .

(2) Observe also that in case  $X$  is a separable this Symmetric Vitali Property is satisfied by any Borel regular measure  $\mu_0$  with  $\mu_0(X) < \infty$  which has the “doubling property” that there is a fixed constant  $C$  such that

$$(\ddagger) \quad \mu_0(B_{2\rho}(x)) \leq C\mu_0(B_\rho(x)) \quad \forall \text{ closed ball } B_\rho(x) \subset X.$$

Indeed in this case, given  $A \subset X$  with  $\mu(A) < \infty$  and a collection  $\mathcal{B}$  of closed balls which cover  $A$  finely, by the Corollary 3.7 of the 5-times Covering Lemma we can select a pairwise disjoint subcollection  $\mathcal{B}'$  (which is countable by the separability of  $X$ , hence

expressible  $\mathcal{B}' = \{B_{\rho_j}(x_j) : j = 1, 2, \dots\}$  with

$$A \setminus \{B_{\rho_1}(x_1), \dots, B_{\rho_N}(x_N)\} \subset \bigcup_{j=N+1}^{\infty} B_{5\rho_j}(x_j)$$

and hence, since  $\mu_0(B_{5\rho}(x)) \leq \mu_0(B_{8\rho}(x)) \leq C^3 \mu_0(B_{\rho}(x))$  by  $(\ddagger)$ ,

$$\mu_0(A \setminus \{B_{\rho_1}(x_1), \dots, B_{\rho_N}(x_N)\}) \leq C^3 \sum_{j=N+1}^{\infty} \mu_0(B_{\rho_j}(x_j)) \rightarrow 0 \text{ as } N \rightarrow \infty$$

because  $\sum_j \mu_0(B_{\rho_j}(x_j)) = \mu_0(\bigcup_j B_{\rho_j}(x_j)) < \infty$ . Thus

$$\mu_0(A \setminus (\bigcup_{j=1}^{\infty} B_{\rho_j}(x_j))) = 0,$$

as claimed.

(3) A very important fact is that *any* Borel regular measure  $\mu_0$  on  $\mathbb{R}^n$  which is finite on each compact subset automatically has the Symmetric Vitali Property. In order to check this we'll need the following famous covering lemma due to Besicovitch:

**4.5 Lemma (Besicovitch Covering Lemma.)** *Suppose  $\mathcal{B}$  is a collection of closed balls in  $\mathbb{R}^n$ , let  $A$  be the set of centers, and suppose the set of all radii of balls in  $\mathcal{B}$  is a bounded set. Then there are sub-collections  $\mathcal{B}_1, \dots, \mathcal{B}_N \subset \mathcal{B}$  ( $N = N(n)$ ) such that each  $\mathcal{B}_j$  is a pairwise disjoint (or empty) collection, and  $\bigcup_{j=1}^N \mathcal{B}_j$  still covers  $A$ —i.e.  $A \subset \bigcup_{j=1}^N (\bigcup_{B \in \mathcal{B}_j} B)$ .*

We emphasize that  $N$  is a certain fixed constant depending only on  $n$ . For the proof of this lemma we refer for example to [EG92] or [Fed69].

**Proof of Remark 4.4(3):** Let  $\mu$  be a Radon measure on  $\mathbb{R}^n$ ,  $A \subset \mathbb{R}^n$  with  $\mu(A) < \infty$ ,  $\mathcal{B}$  a collection of closed balls with centers in  $A$  covering  $A$  finely. By the Besicovitch lemma we can choose  $\mathcal{B}_1, \dots, \mathcal{B}_N \subset \{B \in \mathcal{B} : \text{diam } B \leq 1\}$  such that  $\bigcup_{j=1}^N \mathcal{B}_j$  covers  $A$ . Then for at least one  $j \in \{1, \dots, N\}$  we get

$$\mu(A \setminus \bigcup_{B \in \mathcal{B}_j} B) \leq (1 - \frac{1}{N})\mu(A)$$

and hence taking a suitable finite subcollection  $\{B_1, \dots, B_Q\} \subset \mathcal{B}_j$ ,

$$\mu(A \setminus \bigcup_{k=1}^Q B_k) \leq (1 - \frac{1}{2N})\mu(A).$$

Since  $\mathcal{B}$  covers  $A$  finely, and since  $\bigcup_{k=1}^Q B_k$  is closed, the collection  $\tilde{\mathcal{B}} = \{B \in \mathcal{B} : B \cap (\bigcup_{k=1}^Q B_k) = \emptyset\}$  covers  $A \setminus \bigcup_{k=1}^Q B_k$  finely, so with  $A \setminus \bigcup_{k=1}^Q B_k$  in place of  $A$  the same argument says that we can select new balls  $B_{Q+1}, \dots, B_P \in \tilde{\mathcal{B}}$  such that

$$\begin{aligned} (1) \quad \mu(A \setminus \bigcup_{k=1}^P B_k) &\leq (1 - \frac{1}{2N})\mu(A \setminus \bigcup_{k=1}^Q B_k) \\ &\leq (1 - \frac{1}{2N})^2 \mu(A). \end{aligned}$$

Continuing (inductively) in this way, we conclude that there is a pairwise disjoint se-

quence  $B_1, B_2, \dots$  of balls in  $\mathcal{B}$  such that

$$\mu(A \setminus \bigcup_{k=1}^{\infty} B_k) = 0.$$

Thus Remark 4.4(3) is established.  $\square$

We now want to prove an analogue of the Comparison Theorem 3.3 in case we use  $\Theta^{*\mu_0}(\mu, x)$  of 4.2 in place of the upper density  $\Theta^{*n}(\mu, x)$ .

**4.6 Theorem.** *Suppose  $\mu, \mu_0$  are open  $\sigma$ -finite (as in 1.21) Borel regular measures on a metric space  $X$ ,  $\mu_0$  has the Symmetric Vitali Property, and  $A \subset X, t \geq 0$ . Then*

$$\Theta^{*\mu_0}(\mu, x) \geq t \text{ for all } x \in A \Rightarrow \mu(A) \geq t\mu_0(A).$$

**Note:**  $A$  is not assumed to be measurable.

**Proof:** The proof is similar to the proof of Theorem 3.3, except that we use the Symmetric Vitali Property for  $\mu_0$  in place of the 5 times Covering Lemma: First let  $U_0$  be the open set of  $\mu_0$  measure zero as in the Definition 4.2. As observed in Remark 4.4(1) we have  $\mu_0(U_0) = 0$ . We can assume without loss of generality that  $t > 0$ . Let  $U \supset A$  be open,  $\tau \in (0, t)$ , and consider the collection  $\mathcal{B}$  of all closed balls  $B_\rho(x) \subset U$  with  $x \in A \cap X \setminus U_0$  such that  $\mu(B_\rho(x)) > \tau\mu_0(B_\rho(x))$ . Evidently  $\mathcal{B}$  covers  $A \cap (X \setminus U_0)$  finely, so by the Symmetric Vitali Property for  $\mu_0$  there is a countable pairwise disjoint subcollection  $B_{\rho_j}(x_j), j = 1, 2, \dots$ , of  $\mathcal{B}$  with  $\mu_0(A \setminus (\bigcup_j B_{\rho_j}(x_j))) = 0$  and  $\mu(B_{\rho_j}(x_j)) \geq \tau\mu_0(B_{\rho_j}(x_j))$  for each  $j$ , and hence by summing we obtain

$$\tau\mu_0(A) \leq \mu(\bigcup_j B_{\rho_j}(x_j)) \leq \mu(U).$$

Since  $\mu(A) = \inf_{U \text{ open}, U \supset A} \mu(U)$  by Theorem 1.22, we thus have the stated result by letting  $\tau \uparrow t$ .  $\square$

Observe that in particular the above comparison lemma gives

**4.7 Corollary.** *If  $\mu, \mu_0$  are as in Theorem 4.6 above then  $\Theta^{*\mu_0}(\mu, x) < \infty$  for  $\mu_0$ -a.e.  $x \in X$ .*

**Proof:** We are given open  $V_j$  with  $X = \bigcup_j V_j$  and  $\mu_0(V_j) < \infty$  for each  $j$ . Let  $\mu_j = \mu \llcorner V_j, j = 1, 2, \dots$ . Theorem 4.6, with  $A_t = \{x \in V_j \setminus U_0 : \Theta^{*\mu_0}(\mu, x) \geq t\}$  in place of  $A$  and  $\mu_j$  in place of  $\mu$ , implies

$$t\mu_0(A_t) \leq \mu_j(A_t) \leq \mu(V_j) \quad \forall t > 0,$$

so  $\mu_0(\{x \in V_j : \Theta^{*\mu_0}(\mu, x) = \infty\}) \leq t^{-1}\mu(V_j)$  for each  $t > 0$ , hence  $\mu_0(\{x \in V_j : \Theta^{*\mu_0}(\mu, x) = \infty\}) = 0$  for each  $j$   $\square$

As a second corollary we state the following general Upper Density Theorem:

**4.8 Theorem (General Upper Density Th.)** *If  $\mu, \mu_0$  are Borel regular measures on a metric space  $X$ , if  $\mu_0$  open  $\sigma$ -finite (as in 1.21) and has the Symmetric Vitali Property, and if  $A$  is a  $\mu$ -measurable subset of  $X$  with  $\mu(A) < \infty$ , then*

$$\Theta^{*\mu_0}(\mu \llcorner A, x) = 0 \text{ for } \mu_0\text{-a.e. } x \in X \setminus A.$$

**Proof:** The proof is essentially the same as the proof of Theorem 3.8, except that we use the general comparison theorem 4.6 in place of 3.3. So let  $C$  be an arbitrary closed subset of  $A$ ,  $t > 0$ , and  $S_t = \{x \in X \setminus A : \Theta^{*\mu_0}(\mu \llcorner A, x) \geq t\}$ . Since  $X \setminus C$  is an open set containing  $X \setminus A$  we have  $S_t = \{x \in X \setminus A : \Theta^{*\mu_0}(\mu \llcorner A \cap (X \setminus C), x) \geq t\}$ , and hence, by the comparison theorem 4.6 with  $S_t$  in place of  $A$ ,

$$t\mu_0(S_t) \leq \mu(S_t \cap (A \setminus C)) \leq \mu(A \setminus C).$$

However by the regularity property 1.22(2) we have  $\inf_{C \text{ closed}, C \subset A} \mu(A \setminus C) = 0$ , so  $\mu_0(S_t) = 0$  for each  $t > 0$ .  $\square$

Using the above theorem we can now prove the general density theorem:

**4.9 Theorem.** *If  $\mu$  is open  $\sigma$ -finite (as in 1.21) Borel regular measure on a metric space  $X$ , if  $\mu$  has the Symmetric Vitali Property, and if  $A$  is a  $\mu$ -measurable subset of  $X$ , then*

$$\lim_{\rho \downarrow 0} \frac{\mu(A \cap B_\rho(x))}{\mu(B_\rho(x))} = \begin{cases} 1 & \mu\text{-a.e. } x \in A \\ 0 & \mu\text{-a.e. } x \in X \setminus A. \end{cases}$$

**Proof:** Since  $X = \cup_j V_j$  with  $V_j$  open and  $\mu(V_j) < \infty$  for each  $j$ , we can assume without loss of generality that  $\mu(X) < \infty$ . As in Remark 4.4(1) we see that the set of  $x \in X$  such that  $\mu(B_\rho(x)) = 0$  for some  $\rho > 0$  is an open set  $U_0$  with  $\mu(U_0) = 0$ . For  $x \in X \setminus U_0$  we have

$$\frac{\mu(A \cap B_\rho(x))}{\mu(B_\rho(x))} + \frac{\mu(B_\rho(x) \setminus A)}{\mu(B_\rho(x))} = 1 \text{ for each } \rho > 0,$$

and the first term on the left  $\rightarrow 0$  for  $\mu$ -a.e.  $x \in X \setminus A$  by the Upper Density Theorem 4.8 with  $\mu_0 = \mu$ , whereas the second term on the left  $\rightarrow 0$  for  $\mu$ -a.e.  $x \in A$  by the same theorem with  $\mu_0 = \mu$  and  $X \setminus A$  in place of  $A$ .  $\square$

The following Lebesgue differentiation theorem is an easy corollary:

**4.10 Corollary.** *If  $X, \mu$  are as in Theorem 4.9 and if  $f : X \rightarrow \mathbb{R}$  is locally  $\mu$ -integrable on  $X$  (i.e.  $f$  is  $\mu$ -measurable and  $x \in X \Rightarrow \int_{B_\rho(x)} |f| d\mu < \infty$  for some  $\rho > 0$ ), then*

$$\lim_{\rho \downarrow 0} (\mu(B_\rho(x)))^{-1} \int_{B_\rho(x)} f d\mu = f(x) \text{ for } \mu\text{-a.e. } x \in X$$

**Proof:** Since  $f = \max\{f, 0\} - \max\{-f, 0\}$  we can assume without loss of generality that  $f \geq 0$ . Also, we can without loss of generality assume  $\mu(X) < \infty$  because it suffices to prove the theorem for  $\mu$ -a.e.  $x \in V_j$ , where  $V_j$  are the open sets as in 1.21.

According to Lusin's Theorem 1.24 there are closed sets  $A_j$ ,  $j = 1, 2, \dots$ , with  $\mu(X \setminus (\cup_j A_j)) = 0$  and  $f|_{A_j}$  continuous for each  $j$ . Then, for any  $x \in X$ ,  $j \in \{1, 2, \dots\}$  and  $\rho > 0$ ,

$$\begin{aligned} & (\mu(B_\rho(x)))^{-1} \int_{B_\rho(x)} f \, d\mu = \\ & (\mu(B_\rho(x)))^{-1} \left( \int_{A_j \cap B_\rho(x)} (f(y) - f(x)) \, d\mu + f(x) \mu(A_j \cap B_\rho(x)) + \nu(B_\rho(x) \setminus A_j) \right) \end{aligned}$$

where  $\nu$  is the Borel regular outer measure on  $X$  corresponding, in the sense described in Remark 1.13, to the Borel measure  $\nu_0$  defined by  $\nu_0(A) = \int_A f \, d\mu$ . By continuity of  $f|_{A_j}$ , Theorem 4.9, and the Upper Density Theorem 4.8 (with  $\nu, \mu$  in place of  $\mu, \mu_0$  respectively), we then have, for  $\mu$ -a.e.  $x \in A_j$ ,  $(\mu(B_\rho(x)))^{-1} \int_{B_\rho(x)} f \, d\mu \rightarrow 0 + f(x) + 0 = f(x)$  as  $\rho \downarrow 0$ .  $\square$

Of course we can also take the lower density  $\Theta_*^{\mu_0}(\mu, x)$  of  $\mu$  with respect to  $\mu_0$  which we define, analogous to the definition of upper density in 4.2, by

$$4.11 \quad \Theta_*^{\mu_0}(\mu, x) = \begin{cases} \liminf_{\rho \downarrow 0} \frac{\mu(B_\rho(x))}{\mu_0(B_\rho(x))} & \text{for } x \in X \setminus (U_0 \cup V_0) \\ \infty & \text{for } x \in U_0 \setminus V_0 \\ 0 & \text{for } x \in V_0, \end{cases}$$

with  $U_0, V_0$  as in 4.2. Then there is an analogue of the Comparison Theorem 4.6 for the lower density. Preparatory to that we need the following lemma:

**4.12 Lemma.** *If  $\mu, \mu_0$  is any pair of Borel regular measures on a metric space  $X$  with  $\mu$   $\sigma$ -finite, then there is a Borel set  $B \subset X$  with  $\mu_0(B) = 0$  and  $\mu \perp (X \setminus B)$  absolutely continuous with respect to  $\mu_0$  (i.e.  $\mu_0(S) = 0 \Rightarrow \mu(S \setminus B) = 0$ ).*

**Proof:** In case  $\mu(X) < \infty$  we let  $\mathcal{A} = \{\text{Borel sets } A \subset X \text{ with } \mu_0(A) = 0\}$  and  $\alpha = \sup\{\mu(A) : A \in \mathcal{A}\}$ . Choose a sequence  $A_j \in \mathcal{A}$  with  $\lim \mu(A_j) = \alpha$ . Then  $B = \cup_j A_j \in \mathcal{A}$  with  $\mu(B) = \alpha$ . By Borel regularity of  $\mu_0$ , if  $S \subset X$  with  $\mu_0(S) = 0$  we can select  $A \in \mathcal{A}$  with  $S \subset A$ , and hence  $\mu(S \setminus B) \leq \mu(A \setminus B) = \mu(B \cup (A \setminus B)) - \mu(B) \leq \alpha - \alpha = 0$ , so  $B$  has the required property.

In the general case we select Borel sets  $A_j$  with  $X = \cup_j A_j$  and  $\mu(A_j) < \infty \, \forall j$ , and, applying Case 1 to  $\mu \perp A_j$ , we obtain Borel sets  $B_j$  with  $\mu_0(B_j) = 0$  and  $\mu \perp (A_j \setminus B_j)$  absolutely continuous with respect to  $\mu_0$ . So, with  $B = \cup_j B_j$ ,  $\mu_0(S) = 0 \Rightarrow \mu(S \setminus B) \leq \sum_j \mu(S \cap A_j \setminus B) \leq \sum_j \mu(S \cap A_j \setminus B_j) = 0$ .  $\square$

**4.13 Remark:** The set  $B$  is evidently unique up to a set of  $\mu$ -measure zero, so the Borel regular measure  $\mu \llcorner (X \setminus B)$  is uniquely determined; it is called the absolutely continuous part of  $\mu$  relative to  $\mu_0$ .

We can now prove an analogue of the Comparison Theorem 4.6 for the lower density:

**4.14 Theorem.** Suppose  $\mu, \mu_0$  are open  $\sigma$ -finite (as in 1.21) Borel regular measures on the metric space  $X$ ,  $t > 0$ , and  $A \subset X$  with  $\Theta_*^{\mu_0}(\mu, x) \leq t$  for all  $x \in A$ .

(i) If  $\mu$  has the Symmetric Vitali Property then  $\mu(A) \leq t\mu_0(A)$ .

(ii) If  $\mu_0$  has the Symmetric Vitali Property then  $\mu(A \setminus B) \leq t\mu_0(A)$ , where  $B$  (with  $\mu_0(B) = 0$ ) is as in 4.12.

**Proof:** The proof is similar to the proof of Theorem 4.6. In view of the open  $\sigma$ -finiteness property we can suppose without loss of generality that both  $\mu(X) < \infty$  and  $\mu_0(X) < \infty$ .

Proof of (i): First observe that  $A \subset X \setminus U_0$  (because, by Definition 4.11,  $\Theta_*^{\mu_0}(\mu, x) = \infty$  on  $U_0$ ). Let  $\tau > t$ . By Theorem 1.22(1) we can select an open  $U \supset A$  with  $\mu_0(U) < \mu_0(A) + \tau - t$ .

Define

$$\mathcal{B} = \{B_\rho(x) \subset U : x \in A \text{ and } \mu(B_\rho(x)) < \tau\mu_0(B_\rho(x))\}.$$

$\mathcal{B}$  evidently covers  $A$  finely, so by the Symmetric Vitali Property for  $\mu$  there is a pairwise disjoint collection  $B_{\rho_j}(x_j)$  with  $\mu(A \setminus (\cup_j B_{\rho_j}(x_j))) = 0$  and  $\mu(B_{\rho_j}(x_j)) \leq \tau\mu_0(B_{\rho_j}(x_j))$  for each  $j$ . By summing on  $j$  we then have  $\mu(A) \leq \tau\mu_0(U) \leq \tau(\mu_0(A) + \tau - t)$ , so letting  $\tau \downarrow t$  gives the required result.

Proof of (ii): With  $B$  be as in Lemma 4.12,  $\tilde{\mu} = \mu \llcorner (X \setminus B)$  is absolutely continuous with respect to  $\mu_0$ , hence the Symmetric Vitali Property for  $\mu_0$  implies the Symmetric Vitali Property for  $\tilde{\mu}$ , so we can apply part (i) with  $A \setminus B$  in place of  $A$  and  $\tilde{\mu}$  in place of  $\mu$ . This gives the required result.  $\square$

We define the density  $\Theta^{\mu_0}(\mu, x)$  to be the common value of  $\Theta^*\mu_0(\mu, x)$  and  $\Theta_*^{\mu_0}(\mu, x)$  at points where these quantities are equal. Thus if  $U_0, V_0$  are the open sets in 4.2 and 4.11, then

$$4.15 \quad \Theta^{\mu_0}(\mu, x) = \begin{cases} \lim_{\rho \downarrow 0} \frac{\mu(B_\rho(x))}{\mu_0(B_\rho(x))} & \text{if } x \in X \setminus (U_0 \cup V_0) \text{ and this limit exists} \\ \infty & \text{at points } x \in U_0 \setminus V_0 \\ 0 & \text{at points } x \in V_0, \end{cases}$$

and  $\Theta^{\mu_0}(\mu, x)$  is undefined at points where  $\Theta_*^{\mu_0}(\mu, x) < \Theta^*\mu_0(\mu, x)$ .

**4.16 Theorem (Differentiation Theorem.)** *Suppose  $\mu, \mu_0$  are open  $\sigma$ -finite (as in 1.21) Borel regular measures on the metric space  $X$ .*

(i) *If  $\mu$  has the Symmetric Vitali Property, then there is a Borel set  $S$  of  $\mu$ -measure zero such that  $\Theta^{\mu_0}(\mu, x)$  (as in 4.15) exists for all  $x \in X \setminus S$ .*

(ii) *If  $\mu_0$  has the Symmetric Vitali Property, then there is a Borel set  $S$  of  $\mu_0$ -measure zero such that  $\Theta^{\mu_0}(\mu, x)$  exists and is finite for all  $x \in X \setminus S$ .*

*In either case  $\Theta^{\mu_0}(\mu, x)$  is a Borel measurable function of  $x \in X \setminus S$ .*

**Proof:** First assume  $\mu_0(X), \mu(X) < \infty$  and let  $A \subset X$  be any Borel set.

To prove (i) first note that by the Comparison Theorems 4.6 and 4.14(i), for any given  $a, b > 0$ ,

$$(1) \quad \Theta_*^{\mu_0}(\mu, x) < a \text{ and } \Theta^{*\mu_0}(\mu, x) > b \text{ for all } x \in A \\ \Rightarrow \mu(A) \leq a\mu_0(A) \text{ and } b\mu_0(A) \leq \mu(A).$$

In particular if  $0 < a < b$  and

$$E_{a,b} = \{x \in X \setminus U_0 : \Theta_*^{\mu_0}(\mu, x) < a < b < \Theta^{*\mu_0}(\mu, x)\}.$$

then  $a^{-1}\mu(E_{a,b}) \leq \mu_0(E_{a,b}) \leq b^{-1}\mu(E_{a,b})$ , which implies that

$$(2) \quad \mu_0(E_{a,b}) = \mu(E_{a,b}) = 0.$$

Since  $\{x : \Theta_*^{\mu_0}(\mu, x) < \Theta^{*\mu_0}(\mu, x)\} = \cup_{a,b \text{ rational}, 0 < a < b} E_{a,b}$  we deduce from (2) that  $\Theta_*^{\mu_0}(\mu, x) = \Theta^{*\mu_0}(\mu, x)$  for  $\mu_0$ -a.e.  $x \in X \setminus U_0$ , so indeed  $\Theta^{\mu_0}(\mu, x)$  exists (in  $[0, \infty]$ ) for  $\mu$ -a.e.  $x \in X \setminus U_0$ .  $\Theta^{\mu_0}(\mu, x)$  is also defined in  $U_0$  by Definition 4.15. Thus  $\Theta^{\mu_0}(\mu, x)$  is well-defined  $\mu$ -a.e., so by Borel regularity of  $\mu$  there is a Borel set  $S$  with  $\mu(S) = 0$  such that  $\Theta^{\mu_0}(\mu, x)$  is well-defined for all  $x \in X \setminus S$ .

The measurability of  $\Theta^{\mu_0}(\mu, x)$  as a function of  $x \in X \setminus S$  is proved as follows: For each fixed  $\rho > 0$ ,  $\mu(B_\rho(x))$  and  $\mu_0(B_\rho(x))$  are positive upper semi-continuous functions of  $x \in X \setminus (S \cup V_0 \cup U_0)$ , hence are Borel measurable functions on  $X \setminus (S \cup V_0 \cup U_0)$ , and hence so is the quotient  $\mu(B_\rho(x))/\mu_0(B_\rho(x))$ . Hence  $\Theta^{\mu_0}(\mu, x) = \lim_{i \rightarrow \infty} \mu(B_{1/i}(x))/\mu_0(B_{1/i}(x))$  is Borel measurable on  $X \setminus (S \cup V_0 \cup U_0)$ . Finally, by Definition 4.15,  $\Theta^{\mu_0}(\mu, x) = \infty$  on  $U_0 \setminus V_0$  and  $\Theta^{\mu_0}(\mu, x) = 0$  on  $V_0$ . Since  $U_0, V_0$  are open we then conclude that indeed  $\Theta^{\mu_0}(\mu, x)$  is Borel measurable in case  $\mu, \mu_0$  are finite measures. In the general open  $\sigma$ -finite case, when there are open sets  $V_j$  with  $\cup_j V_j = X$  and  $\mu(V_j), \mu_0(V_j) < \infty$ , we apply the above with  $\mu \llcorner V_j, \mu_0 \llcorner V_j$  in place of  $\mu, \mu_0$  respectively.

To prove (ii), note first that by Corollary 4.7 we have

$$(3) \quad \Theta^{*\mu_0}(\mu, x) < \infty \text{ for } \mu_0\text{-a.e. } x \in X.$$

As in 4.12, let  $B$  be a Borel set of  $\mu_0$ -measure zero such that  $\tilde{\mu} = \mu \llcorner (X \setminus B)$  is absolutely continuous with respect to  $\mu_0$ . Then  $\tilde{\mu}$  has the Symmetric Vitali Property, and hence the argument of (i) above applies with  $\tilde{\mu}$  in place of  $\mu$  to give

$$(4) \quad \mu_0(E_{a,b}) = \mu(E_{a,b} \setminus B) = 0,$$

in place of (2). Hence  $\Theta^{\mu_0}(\mu, x)$  exists for  $\mu_0$ -a.e.  $x \in X$ , and by (3) it is also finite for  $\mu_0$ -a.e.  $x \in X$ , hence there is a Borel set  $S$  with  $\mu_0(S) = 0$  such that  $\Theta^{\mu_0}(\mu, x)$  exists and is finite for all  $x \in X \setminus S$ .

The measurability of  $\Theta^{\mu_0}(\mu, x)$  follows similarly to case (i) above.  $\square$

Next, recall the abstract Radon-Nikodym theorem, which says that if  $\mu, \mu_0$  are abstract  $\sigma$ -finite measures on a  $\sigma$ -algebra  $\mathcal{A}$  of subsets of an abstract space  $X$ , and if  $\mu$  is absolutely continuous with respect to  $\mu_0$  (i.e.  $A \in \mathcal{A}$  with  $\mu_0(A) = 0 \Rightarrow \mu(A) = 0$ ), then there is a non-negative  $\mathcal{A}$ -measurable function  $\Theta$  on  $X$  such that

$$4.17 \quad \mu(A) = \int_A \Theta d\mu_0, \quad A \in \mathcal{A}.$$

In these circumstances the function  $\Theta$  is called “the Radon-Nikodym derivative” of  $\mu$  with respect to  $\mu_0$ , denoted  $\frac{d\mu}{d\mu_0}$  or  $D_{\mu_0}\mu$ .

We show here that in case  $\mu, \mu_0$  are Borel regular open  $\sigma$ -finite (as in 1.21) on the metric space  $X$  with  $\mu_0$  having the Symmetric Vitali Property, then the Radon-Nikodym derivative  $D_{\mu_0}\mu(x)$  is just the density  $\Theta^{\mu_0}(\mu, x) = \lim_{\rho \downarrow 0} \frac{\mu(B_\rho(x))}{\mu_0(B_\rho(x))}$ :

**4.18 Theorem (Radon-Nikodym.)** *Suppose  $\mu, \mu_0$  are open  $\sigma$ -finite (as in 1.21) Borel regular measures on  $X$ , and  $\mu_0$  has the Symmetric Vitali Property.*

(i) *If  $\mu$  is absolutely continuous with respect to  $\mu_0$  (i.e.  $E \subset X$  with  $\mu_0(E) = 0 \Rightarrow \mu(E) = 0$  and hence  $\mu$  also has the Symmetric Vitali Property), then*

$$(*) \quad \mu(A) = \int_A \Theta^{\mu_0}(\mu, x) d\mu_0(x) \text{ for every Borel set } A \subset X.$$

(ii) *If we drop the condition that  $\mu$  is absolutely continuous with respect to  $\mu_0$ , then in place of (\*) we can still conclude that there is a Borel set  $Z$  with  $\mu_0(Z) = 0$  and*

$$(\ddagger) \quad \mu(A) = \int_A \Theta^{\mu_0}(\mu, x) d\mu_0(x) + (\mu \llcorner Z)(A)$$

*for each Borel set  $A \subset X$ .*

(iii) *Finally, if  $\mu$  also has the Symmetric Vitali Property, then we get (\ddagger) with*

$$Z = \{x \in X : \Theta^{\mu_0}(\mu, x) = \infty\}$$

*(which is a set of  $\mu_0$ -measure zero by 4.16(ii)).*

**4.19 Remarks:** (1) By Remark 4.4(3) we always have the conclusion of 4.16(iii) if  $X = \mathbb{R}^n$ .

(2)  $\mu \perp Z$  is called the singular part of  $\mu$  with respect to  $\mu_0$ .

**Proof of Theorem 4.18:** Since  $\mu, \mu_0$  are open  $\sigma$ -finite, we can assume  $\mu(X) < \infty, \mu_0(X) < \infty$ . Let  $S$  be a Borel set of  $\mu_0$ -measure zero as in Theorem 4.16. For any Borel set  $A \subset X \setminus S$  let

$$v(A) = \int_A \Theta^{\mu_0}(\mu, x) d\mu_0(x)$$

and for any subset  $A \subset X \setminus S$  let  $v(A) = \inf_{B \supset A, B \text{ Borel}} v(B)$ . By Remark 1.13,  $v$  is a Radon measure and, with  $0 < a < b$ ,  $A_{a,b} = \{x \in A : a < \Theta^{\mu_0}(\mu, x) < b\}$  and  $A$  any Borel set, we have

$$a\mu_0(A_{a,b}) \leq v(A_{a,b}) \leq b\mu_0(A_{a,b}).$$

On the other hand the Comparison Theorems 4.6, 4.14(i) imply

$$a\mu_0(A_{a,b}) \leq \mu(A_{a,b}) \leq b\mu_0(A_{a,b}),$$

and so

$$\frac{a}{b}\mu(A_{a,b}) \leq v(A_{a,b}) \leq \frac{b}{a}\mu(A_{a,b})$$

and it follows that  $v(A) = \mu(A)$ . Thus  $(*)$  is proved.

In the general case (when we allow the possibility that there are sets  $A$  with  $\mu_0(A) = 0$  and  $\mu(A) > 0$ ), we can apply the previous argument to the Borel regular measure  $\tilde{\mu} = \mu \perp (X \setminus B)$ , where  $B$  is the set of  $\mu_0$ -measure zero of Lemma 4.12. This gives

$$\mu(A \setminus B) = \int_A \Theta^{\mu_0}(\mu, x) d\mu_0 \quad \forall \text{ Borel set } A \subset X.$$

Thus 4.16( $\ddagger$ ) holds with  $Z = B$ .

Finally, in case  $\mu$  also has the Symmetric Vitali Property, Theorem 4.16(i) establishes that  $\Theta^{\mu_0}(\mu, x)$  exists  $\mu$ -almost everywhere (as well as  $\mu_0$ -almost everywhere) in  $X$ . On the other hand if  $\tilde{X} = X \setminus U_0$  and  $A \subset \{x \in \tilde{X} : \Theta^{\mu_0}(\mu, x) < \infty\} (= \cup_{n=1}^{\infty} \{x \in \tilde{X} : \Theta^{\mu_0}(\mu, x) < n\})$  then by Theorem 4.14(i)

$$\mu_0(A) = 0 \Rightarrow \mu(A) = 0,$$

and we can therefore apply  $(*)$  with  $\mu \perp (X \setminus Z)$ ,  $Z = \{x : \Theta^{\mu_0}(\mu, x) = \infty\}$ , in place of  $\mu$ . Hence (iii) is proved.  $\square$

## 5 Radon Measures, Representation Theorem

In this section we work mainly in locally compact Hausdorff spaces, and for the reader's convenience we recall some basic definitions and preliminary topological results for such spaces.

Recall that a topological space is said to be Hausdorff if it has the property that for every pair of distinct points  $x, y \in X$  there are open sets  $U, V$  with  $x \in U$ ,  $y \in V$  and  $U \cap V = \emptyset$ . In such a space *all compact sets are automatically closed*, the proof of which is as follows: observe that if  $x \notin K$  then for each  $y \in K$  we can (by definition of Hausdorff space) pick open  $U_y, V_y$  with  $x \in U_y$ ,  $y \in V_y$  and  $U_y \cap V_y = \emptyset$ . By compactness of  $K$  there is a finite set  $y_1, \dots, y_N \in K$  with  $K \subset \bigcup_{j=1}^N V_{y_j}$ . But then  $\bigcap_{j=1}^N U_{y_j}$  is an open set containing  $x$  which is disjoint from  $\bigcup_j V_{y_j}$  and hence disjoint from  $K$ , so that  $K$  is closed as claimed. In fact we proved a bit more: that for each  $x \notin K$  there are disjoint open sets  $U, V$  with  $x \in U$  and  $K \subset V$ . Then if  $L$  is another compact set disjoint from  $K$  we can repeat this for each  $x \in L$  thus obtaining disjoint open  $U_x, V_x$  with  $x \in U_x$  and  $K \subset V_x$ , and then compactness of  $L$  implies  $\exists x_1, \dots, x_M \in L$  such that  $L \subset \bigcup_{j=1}^M U_{x_j}$  and then  $\bigcup_{j=1}^M U_{x_j}$  and  $\bigcap_{j=1}^M V_{x_j}$  are disjoint open sets containing  $L$  and  $K$  respectively. By a simple inductive argument (left as an exercise) we can extend this to finite pairwise disjoint unions of compact subsets:

**5.1 Lemma.** *Let  $X$  be a Hausdorff space and  $K_1, \dots, K_N$  be pairwise disjoint compact subsets of  $X$ . Then there are pairwise disjoint open subsets  $U_1, \dots, U_N$  with  $K_j \subset U_j$  for each  $j = 1, \dots, N$ .*

Notice in particular that we have the following corollary of Lemma 5.1:

**5.2 Corollary.** *A compact Hausdorff space is normal: i.e. given closed disjoint subsets  $K_1, K_2$  of a compact Hausdorff space, we can find disjoint open  $U_1, U_2$  with  $K_j \subset U_j$  for  $j = 1, 2$ .*

Most of the rest of the discussion here takes place in locally compact Hausdorff space: A space  $X$  is said to be *locally compact* if for each  $x \in X$  there is a neighborhood  $U_x$  of  $x$  such that the closure  $\overline{U}_x$  of  $U_x$  is compact.

An important preliminary lemma in such spaces is:

**5.3 Lemma.** *If  $X$  is a locally compact Hausdorff space and  $V$  is a neighborhood of a point  $x$ , then there is a neighborhood  $U_x$  of  $x$  such that  $\overline{U}_x$  is a compact subset of  $V$ .*

**Proof:** First pick a neighborhood  $W_0$  of  $x$  such that  $\overline{W}_0$  is compact and define  $W = W_0 \cap V$ . Then  $\overline{W}$  is compact and hence, with the subspace topology, is normal by Corollary 1 above. Hence since  $\overline{W} \setminus W$  and  $\{x\}$  are disjoint closed sets in this space, and since open

sets in the subspace  $\overline{W}$  can by definition be expressed as the intersection of open sets from  $X$  with the subset  $\overline{W}$ , we can find open  $U_1, U_2$  in the space  $X$  with  $x \in U_1$ ,  $\overline{W} \setminus W \subset U_2$  and  $U_1 \cap U_2 \cap \overline{W} = \emptyset$ . The last identity says  $U_1 \cap \overline{W} \subset \overline{W} \setminus U_2$ , whence  $x \in U_1 \cap W \subset \overline{W} \setminus U_2 \subset W \subset V$ , and since  $\overline{W} \setminus U_2$  is a closed set, we then have  $x \in U_1 \cap W \subset \overline{U_1 \cap W} \subset \overline{W} \setminus U_2 \subset V$ , so the lemma is proved with  $U_x = U_1 \cap W$ .  $\square$

**Remark:** In locally compact Hausdorff space, using Lemmas 5.1 and 5.3 it is easy to check that we can select the  $U_j$  in Lemma 5.1 above to have compact pairwise disjoint closures.

The following lemma is a version of the Urysohn lemma valid in locally compact Hausdorff space:

**5.4 Lemma.** *Let  $X$  be a locally compact Hausdorff space,  $K \subset X$  compact, and  $K \subset V$ ,  $V$  open. Then there is an open  $U \supset K$  with  $\overline{U} \subset V$ ,  $\overline{U}$  compact, and an  $f : X \rightarrow [0, 1]$  with  $f \equiv 1$  in a neighborhood of  $K$  and  $f \equiv 0$  on  $X \setminus U$ .*

**Proof:** By Lemma 5.3 each  $x \in K$  has a neighborhood  $U_x$  with  $\overline{U}_x \subset V$ . Then by compactness of  $K$  we have  $K \subset U \equiv \bigcup_{j=1}^N U_{x_j}$  for some finite collection  $x_1, \dots, x_N \in K$  and  $\overline{U} = \bigcup_{j=1}^N \overline{U}_{x_j} \subset V$ . Now  $\overline{U}$  is compact, so by Corollary 1 it is a normal space and the Urysohn lemma can be applied to give  $f_0 : \overline{U} \rightarrow [0, 1]$  with  $f_0 \equiv 1$  on  $K$  and  $f_0 \equiv 0$  on  $\overline{U} \setminus U$ . Then of course the function  $f_1$  defined by  $f_1 \equiv f_0$  on  $\overline{U}$  and  $f_1 \equiv 0$  on  $X \setminus \overline{U}$  is continuous (check!) because  $f|_{\overline{U}}$  is continuous and  $f$  is identically zero (the value of  $f|_{X \setminus \overline{U}}$ ) on the overlap set  $\overline{U} \setminus U \equiv \overline{U} \cap (X \setminus U)$ . Finally we let  $f \equiv 2 \min\{f_1, \frac{1}{2}\}$  and observe that  $f$  is then identically 1 in the set where  $f_1 > \frac{1}{2}$ , which is an open set containing  $K$ , and  $f$  evidently has all the remaining stated properties.  $\square$

The following corollary of Lemma 5.4 is important:

**5.5 Corollary (Partition of Unity.)** *If  $X$  is a locally compact Hausdorff space,  $K \subset X$  is compact, and if  $U_1, \dots, U_N$  is any open cover for  $K$ , then there exist continuous  $\varphi_j : X \rightarrow [0, 1]$  such that support  $\varphi_j$  is a compact subset of  $U_j$  for each  $j$ , and  $\sum_{j=1}^N \varphi_j \equiv 1$  in a neighborhood of  $K$ .*

**Proof:** By Lemma 5.3, for each  $x \in K$  there is a  $j \in \{1, \dots, N\}$  and a neighborhood  $U_x$  of  $x$  such that  $\overline{U}_x$  is a compact subset of this  $U_j$ . By compactness of  $K$  we have finitely many of these neighborhoods, say  $U_{x_1}, \dots, U_{x_N}$ , with  $K \subset \bigcup_{i=1}^N U_{x_i}$ . Then for each  $j = 1, \dots, N$  we define  $V_j$  to be the union of all  $U_{x_i}$  such that  $\overline{U}_{x_i} \subset U_j$ . Then the  $\overline{V}_j$  is a compact subset of  $U_j$  for each  $j$ , and the  $V_j$  cover  $K$ . So by Lemma 5.4 for each  $j = 1, \dots, N$  we can select  $\psi_j : X \rightarrow [0, 1]$  with  $\psi_j \equiv 1$  on  $\overline{V}_j$  and  $\psi_j \equiv 0$  on  $X \setminus W_j$  for some open  $W_j$  with  $\overline{W}_j$  a compact subset of  $U_j$  and  $W_j \supset \overline{V}_j$ . We can also use Lemma 5.4 to select  $f_0 : X \rightarrow [0, 1]$  with  $f_0 \equiv 1$  in the neighborhood  $\bigcup_{j=1}^N V_j$  of  $K$  and

$f_0 \equiv 0$  on  $\{x : \sum_{j=1}^N \psi_j(x) = 0\}$ . (This latter set is closed and has (open) complement which is a neighborhood of the compact set  $\cup_{j=1}^N \overline{V}_j$  and so we can indeed construct such  $f_0$  by Lemma 5.4.) Then set  $\psi_0 = 1 - f_0$  and observe that by construction  $\sum_{i=0}^N \psi_i > 0$  everywhere on  $X$ , so we can define continuous functions  $\varphi_j$  by

$$\varphi_j = \frac{\psi_j}{\sum_{i=0}^N \psi_i}, \quad j = 1, \dots, N.$$

Evidently these functions have the required properties.  $\square$

We now give the definition of Radon measure. Radon measures are typically used only in locally compact Hausdorff space, but the definition and the first two lemmas following it are valid in arbitrary Hausdorff space:

**5.6 Definition:** Given a Hausdorff space  $X$ , a “Radon measure” on  $X$  is an outer measure  $\mu$  on  $X$  having the 3 properties:

$$\mu \text{ is Borel regular and } \mu(K) < \infty \quad \forall \text{ compact } K \subset X \quad (\text{R1})$$

$$\mu(A) = \inf_{U \text{ open, } U \supset A} \mu(U) \text{ for each subset } A \subset X \quad (\text{R2})$$

$$\mu(U) = \sup_{K \text{ compact, } K \subset U} \mu(K) \text{ for each open } U \subset X. \quad (\text{R3})$$

Such measures automatically have a property like (R3) with an arbitrary  $\mu$ -measurable subset of finite measure:

**5.7 Lemma.** *Let  $X$  be a Hausdorff space and  $\mu$  a Radon measure on  $X$ . Then  $\mu$  automatically has the property*

$$\mu(A) = \sup_{K \subset A, K \text{ compact}} \mu(K)$$

*for every  $\mu$ -measurable set  $A \subset X$  with  $\mu(A) < \infty$ .*

**Proof:** Let  $\varepsilon > 0$ . By definition of Radon measure we can choose an open  $U$  containing  $A$  with  $\mu(U \setminus A) < \varepsilon$ , and then a compact  $K \subset U$  with  $\mu(U \setminus K) < \varepsilon$  and finally an open  $W$  containing  $U \setminus A$  with  $\mu(W \setminus (U \setminus A)) < \varepsilon$  (so that  $\mu(W) \leq \varepsilon + \mu(U \setminus A) < 2\varepsilon$ ). Then we have that  $K \setminus W$  is a compact subset of  $U \setminus W$ , which is a subset of  $A$ , and also

$$\mu(A \setminus (K \setminus W)) \leq \mu(U \setminus (K \setminus W)) \leq \mu(U \setminus K) + \mu(W) \leq 3\varepsilon,$$

which completes the proof.  $\square$

The following lemma asserts that the defining property (R1) of Radon measures follows automatically from the remaining two properties ((R2) and (R3)) in case  $\mu$  is finite and additive on finite disjoint unions of compact sets.

**5.8 Lemma.** *Let  $X$  be a Hausdorff space and assume that  $\mu$  is an outer measure on  $X$  satisfying the properties (R2), (R3) above, and in addition assume that*

$$\mu(K_1 \cup K_2) = \mu(K_1) + \mu(K_2) < \infty \text{ whenever } K_1, K_2 \text{ are compact and disjoint.}$$

*Then (R1) holds and hence  $\mu$  is a Radon measure.*

**Proof:** Note that (R2) implies that for every set  $A \subset X$  we can find open sets  $U_j$  such that  $A \subset \bigcap_j U_j$  and  $\mu(A) = \mu(\bigcap_j U_j)$ . So to complete the proof of (R1) we just have to check that all Borel sets are  $\mu$ -measurable; since the  $\mu$ -measurable sets form a  $\sigma$ -algebra and the Borel sets form the smallest  $\sigma$ -algebra of subsets of  $X$  which contains all the open sets, we thus need only to check that all open sets are  $\mu$ -measurable.

Let  $\varepsilon > 0$  be arbitrary,  $Y$  an arbitrary subset of  $X$  with  $\mu(Y) < \infty$  and let  $U$  be an arbitrary open subset of  $X$ . By (R2) we can pick an open set  $V \supset Y$  with  $\mu(V) < \mu(Y) + \varepsilon$  and by (R3) we can pick a compact set  $K_1 \subset V \cap U$  with  $\mu(V \cap U) \leq \mu(K_1) + \varepsilon$ , and then a compact set  $K_2 \subset V \setminus K_1$  with  $\mu(V \setminus K_1) \leq \mu(K_2) + \varepsilon$ . Then

$$\begin{aligned} \mu(V \setminus U) + \mu(V \cap U) &\leq \mu(V \setminus K_1) + \mu(K_1) + \varepsilon \\ &\leq \mu(K_2) + \mu(K_1) + 2\varepsilon \\ &= \mu(K_2 \cup K_1) + 2\varepsilon \text{ (by (i))} \\ &\leq \mu((V \setminus K_1) \cup K_1) + 2\varepsilon = \mu(V) + 2\varepsilon \leq \mu(Y) + 3\varepsilon, \end{aligned}$$

hence  $\mu(Y \setminus U) + \mu(Y \cap U) \leq \mu(V \setminus U) + \mu(V \cap U) \leq \mu(Y) + 3\varepsilon$  which by arbitrariness of  $\varepsilon$  gives  $\mu(Y \setminus U) + \mu(Y \cap U) \leq \mu(Y)$ , which establishes the  $\mu$ -measurability of  $U$ . Thus all open sets are  $\mu$ -measurable, and hence all Borel sets are  $\mu$ -measurable, and so (R1) is established.  $\square$

The following lemma guarantees the convenient fact that, in a locally compact space such that all open subsets are  $\sigma$ -compact, all locally finite Borel regular outer measures are in fact Radon measures.

**5.9 Lemma.** *Let  $X$  be a locally compact Hausdorff space and suppose that each open set is the countable union of compact subsets. Then any Borel regular outer measure on  $X$  which is finite on each compact set is automatically a Radon measure.*

**Proof:** First observe that in a Hausdorff space  $X$  the statement “each open set is the countable union of compact subsets” is equivalent to the statement “ $X$  is  $\sigma$ -compact (i.e. the countable union of compact sets) and every closed set is the countable intersection of open sets” as one readily checks by using De Morgan’s laws and the fact that a set is open if and only if its complement is closed. Thus we have at our disposal the facts that

$X$  is  $\sigma$ -compact and every closed set is a countable intersection of open sets. The latter fact enables us to apply the Theorem 1.22 on Borel regular outer measures, and we can therefore assert that

- (1)  $\mu(A) = \inf_{U \text{ open}, A \subset U} \mu(U)$  whenever  $A \subset X$  has the property  
 $\exists \text{ open } V_j \text{ with } A \subset \cup_j V_j \text{ and } \mu(V_j) < \infty \forall j$   
 and
- (2)  $\mu(A) = \sup_{C \text{ closed}, C \subset A} \mu(C)$ , provided  $A = \cup_j A_j$  with  
 $A_j$  is  $\mu$ -measurable and  $\mu(A_j) < \infty \forall j$ .

Now observe that, in a locally compact Hausdorff space  $X$ , for each compact  $K \subset X$  there is an open set  $V \supset K$  such that  $\overline{V}$  (the closure of  $V$ ) is compact. If  $X = \cup_{j=1}^{\infty} K_j$ , where each  $K_j$  is compact, that we can apply this with  $K_j$  in place of  $K$ , and we deduce that there are open sets  $V_j$  in  $X$  such that  $\cup_j V_j = X$  and  $\mu(V_j) < \infty$  for each  $j$ , and so in this case (when  $X$  is  $\sigma$ -compact) the identity in (1) holds for every subset  $A \subset X$ ; that is

$$\mu(A) = \inf_{U \text{ open}, A \subset U} \mu(U) \text{ for every } A \subset X,$$

which is the property (R2). Next we note that if  $A \subset X$  is  $\mu$ -measurable, then we can write  $A = \cup_j A_j$ , where  $A_j = A \cap K_j$  (because  $X = \cup_j K_j$ ) and  $\mu(A_j) \leq \mu(K_j) < \infty$  for each  $j$ , so (2) actually holds for every  $\mu$ -measurable  $A$  in case  $X$  is  $\sigma$ -compact (i.e. in case  $X = \cup_{j=1}^{\infty} K_j$  with  $K_j$  compact), and for any closed set  $C$  we can write  $C = \cup_j C_j$  where  $C_j$  is the increasing sequence of compact sets given by  $C_j = C \cap (\cup_{i=1}^j K_i)$  and so  $\mu(C) = \lim_j \mu(C_j)$  and hence  $\mu(C) = \sup_{K \subset C, K \text{ compact}} \mu(K)$ . Thus in the  $\sigma$ -compact case (2) actually tells us that  $\mu(A) = \sup_{K \subset A, K \text{ compact}} \mu(K)$  for any  $\mu$ -measurable set  $A$ . This in particular holds for  $A =$  an open set, which is the remaining property (R3) we needed.  $\square$

Next we have the following important density result:

**5.10 Theorem.** *Let  $X$  be a locally compact Hausdorff space,  $\mu$  a Radon measure on  $X$  and  $1 \leq p < \infty$ . Then  $C_c(X)$  is dense in  $L^p(\mu)$ ; that is, for each  $\varepsilon > 0$  and each  $f \in L^p$  there is a  $g \in C_c(X)$  such that  $\|g - f\|_p < \varepsilon$ .*

In view of Remark 1.13 and Lemma 5.9 we see that Theorem 5.10 directly implies the following:

**5.11 Corollary.** *If  $X$  is a locally compact Hausdorff space such that every open set in  $X$  is the countable union of compact sets, and if  $\mu$  is any Borel regular outer measure on  $X$  which is finite on each compact set, then the space  $C_c(X)$  is dense in  $L^1(\mu)$ .*

**Proof of Theorem 5.10:** Let  $f : X \rightarrow \mathbb{R}$  be  $\mu$ -measurable with  $\|f\|_p < \infty$  and let  $\varepsilon > 0$ . Observe that the simple functions are dense in  $L^p(\mu)$  (which one can check using the dominated convergence theorem and the fact that both  $f_+$  and  $f_-$  can be expressed as the pointwise limits of increasing sequences of non-negative simple functions), so we can pick a simple function  $\varphi = \sum_{j=1}^N a_j \chi_{A_j}$ , where the  $a_j$  are distinct non-zero reals and  $A_j$  are pairwise disjoint  $\mu$ -measurable subsets of  $X$ , such that  $\|f - \varphi\|_p < \varepsilon$ . Since  $\|\varphi\|_p \leq \|\varphi - f\|_p + \|f\|_p < \infty$  we must then have  $\mu(A_j) < \infty$  for each  $j$ . Pick  $M > \max\{|a_1|, \dots, |a_N|\}$  and use Lemma 5.7 to select compact  $K_j \subset A_j$  with  $\mu(A_j \setminus K_j) < \varepsilon^p / (2^{p+1} M^p N)$ . Also, using the definition of Radon measure, we can find open  $U_j \supset K_j$  with  $\mu(U_j \setminus K_j) < \varepsilon^p / (2^{p+1} M^p N)$  and by Lemma 5.7 we can assume without loss of generality that these open sets  $U_1, \dots, U_N$  are pairwise disjoint (otherwise replace  $U_j$  by  $U_j \cap U_j^0$ , where  $U_1^0, \dots, U_N^0$  are pairwise disjoint open sets with  $K_j \subset U_j^0$ ). By Lemma 5.4 we have  $g_j \in C_c(X)$  with  $g_j \equiv a_j$  on  $K_j$ ,  $\{x : g_j(x) \neq 0\}$  contained in a compact subset of  $U_j$ , and  $\sup |g_j| \leq |a_j|$ , and hence by the pairwise disjointness of the  $U_j$  we have that  $g \equiv \sum_{j=1}^N g_j$  agrees with  $\varphi$  on each  $K_j$  and  $\sup |g| = \sup |\varphi| < M$ . Then  $\varphi - g$  vanishes off the set  $\cup_j ((U_j \setminus K_j) \cup (A_j \setminus K_j))$  and we have  $\int_X |\varphi - g|^p d\mu \leq \sum_j \int_{(U_j \setminus K_j) \cup (A_j \setminus K_j)} |\varphi - g|^p d\mu \leq (2M)^p \sum_j (\mu(A_j \setminus K_j) + \mu(U_j \setminus K_j)) \leq \varepsilon^p$ , and hence  $\|f - g\|_p \leq \|f - \varphi\|_p + \|\varphi - g\|_p \leq 2\varepsilon$ , as required.  $\square$

We now state the Riesz representation theorem for non-negative functionals on the space  $\mathcal{K}_+$ , where, here and subsequently,  $\mathcal{K}_+$  denotes the set of non-negative  $C_c(X, \mathbb{R})$  functions, i.e. the set of continuous functions  $f : X \rightarrow [0, \infty)$  with compact support.

**5.12 Theorem (Riesz for non-negative functionals.)** *Suppose  $X$  is a locally compact Hausdorff space,  $\lambda : \mathcal{K}_+ \rightarrow [0, \infty)$  with  $\lambda(cf) = c\lambda(f)$ ,  $\lambda(f + g) = \lambda(f) + \lambda(g)$  whenever  $c \geq 0$  and  $f, g \in \mathcal{K}_+$ , where  $\mathcal{K}_+$  is the set of all non-negative continuous functions  $f$  on  $X$  with compact support. Then there is a Radon measure  $\mu$  on  $X$  such that  $\lambda(f) = \int_X f d\mu$  for all  $f \in \mathcal{K}_+$ .*

Before we begin the proof of 5.12 we observe the following 2 facts about the functional  $\lambda$ :

**5.13 Remarks (1):** Observe that if  $f, g \in \mathcal{K}_+$  with  $f \leq g$  then  $g - f \in \mathcal{K}_+$  and hence  $\lambda(g) = \lambda(f + (g - f)) = \lambda(f) + \lambda(g - f) \geq \lambda(f)$ .

**(2)** If  $K$  is compact, if support  $f \subset K$  and if  $g \in \mathcal{K}_+$  with  $g \equiv 1$  on  $K$ , then we have  $f \leq (\sup f)g$  and  $f g = f$ , so by Remark (1) above we have

$$(*) \quad \lambda(f) \leq (\sup f) \lambda(g), \quad f \in \mathcal{K}_+, \text{ support } f \subset K.$$

Notice in particular that if  $U$  is an arbitrary neighborhood of  $K$  then we can by Lemma 5.4 select neighborhood  $W$  of  $K$  with  $\overline{W}$  a compact subset of  $U$  and a  $g \in \mathcal{K}_+$  with  $g \equiv 1$

in a neighborhood of  $\overline{W}$ ,  $g \leq 1$  everywhere, and support  $g \subset U$ , whence the above inequality with  $\overline{W}$  in place of  $K$  implies

$$(**) \quad \sup_{f \in \mathcal{K}_+, f \leq 1, \text{ support } f \subset W} \lambda(f) \leq \inf_{g \in \mathcal{K}_+, g \leq 1, g \equiv 1 \text{ in a nhd. of } \overline{W}, \text{ support } g \subset U} \lambda(g).$$

**Proof of Theorem 5.12:** For  $U \subset X$  open, we define

$$(1) \quad \mu(U) = \sup_{f \in \mathcal{K}_+, f \leq 1, \text{ support } f \subset U} \lambda(f),$$

and for arbitrary  $A \subset X$  we define

$$(2) \quad \mu(A) = \inf_{U \text{ open}, U \supset A} \mu(U).$$

Notice that these definitions are consistent when  $A$  is itself open. Notice also that by  $(**)$  we have  $\mu(K) < \infty$  for each compact  $K$ ; indeed  $(**)$  and the definitions (1), (2) evidently imply

$$(3) \quad \mu(K) = \inf_{g \in \mathcal{K}_+, g \leq 1, g \equiv 1 \text{ in a nhd. of } K} \lambda(g) \text{ for each compact } K \subset X,$$

Next we prove that  $\mu$  is an outer measure. To see this, first let  $U_1, U_2, \dots$  be open and  $U = \cup_j U_j$ , then for any  $f \in \mathcal{K}_+$  with  $\sup f \leq 1$  and support  $f \subset U$  we have, by compactness of support  $f$ , that support  $f \subset \cup_{j=1}^N U_j$  for some integer  $N$ , and by using a partition of unity  $\varphi_1, \dots, \varphi_N$  for support  $f$  subordinate to  $U_1, \dots, U_N$  (see the Corollary to Lemma 5.4 above), we have  $\lambda(f) = \sum_{j=1}^N \lambda(\varphi_j f) \leq \sum_{j=1}^N \mu(U_j)$ . Taking sup over all such  $f$  we then have  $\mu(U) \leq \sum_j \mu(U_j)$ . It then easily follows that  $\mu(\cup_j A_j) \leq \sum_j \mu(A_j)$  for each  $j$ . Since we trivially also have  $\mu(\emptyset) = 0$  and  $A \subset B \Rightarrow \mu(A) \leq \mu(B)$  we thus have that  $\mu$  is an outer measure on  $X$ .

Finally we want to show that  $\mu$  is a Radon measure. For this we are going to use Lemma 5.8, so we have to check (R2), (R3) and the additivity property  $\mu(K_1 \cup K_2) = \mu(K_1) + \mu(K_2)$  whenever  $K_1, K_2$  are disjoint compact sets. But hypothesis (R2), (R3) are true by the definitions (1), (2), so we only have to check the additivity on disjoint compact sets. In fact if  $K_1$  and  $K_2$  are disjoint compact subsets then for  $\varepsilon > 0$  we can use (3) to find  $g \in \mathcal{K}_+$  with  $g \leq 1$ ,  $g \equiv 1$  in a neighborhood  $W$  of  $K_1 \cup K_2$ , and with  $\lambda(g) \leq \mu(K_1 \cup K_2) + \varepsilon$ . By Lemma 5.1 we can then select disjoint open  $U_1, U_2$  with  $K_1 \subset U_1$  and  $K_2 \subset U_2$ , and by Lemma 5.4 we can select  $f_1, f_2 \in \mathcal{K}_+$  with  $f_j \equiv 1$  in a neighborhood of  $K_j$  such that support  $f_j$  is a compact subset of  $U_j$  and  $f_j \leq 1$  everywhere,  $j = 1, 2$ . Then by (3)  $\mu(K_1) + \mu(K_2) \leq \lambda(f_1 \cdot g) + \lambda(f_2 \cdot g) = \lambda((f_1 + f_2) \cdot g) \leq \lambda(g) \leq \mu(K_1 \cup K_2) + \varepsilon$ . Thus  $\mu(K_1) + \mu(K_2) \leq \mu(K_1 \cup K_2)$ ,

and of course the reverse inequality holds by subadditivity of  $\mu$ , hence the hypotheses of Lemma 5.8 are all established and  $\mu$  is a Radon measure.

Next observe that by  $(*)$  we have  $\lambda(h) \leq \mu(\text{support } h) \sup h$ ,  $h \in \mathcal{K}_+$ , and hence (observing that  $h$  is the uniform limit of  $\max\{h - 1/n, 0\}$  in  $X$ ) we have

$$(4) \quad \lambda(h) \leq \mu(\{x : h(x) > 0\}) \sup h, \quad h \in \mathcal{K}_+.$$

For  $f \in \mathcal{K}_+$  and  $\varepsilon > 0$ , we can select points  $0 = t_0 < t_1 < t_2 < \dots < t_{N-1} < \sup f < t_N$  with  $t_j - t_{j-1} < \varepsilon$  for each  $j = 1, \dots, N$  and with  $\mu(\{f^{-1}\{t_j\}\}) = 0$  for each  $j = 1, \dots, N$ . Notice that the latter requirement is no problem because  $\mu(\{f^{-1}\{t\}\}) = 0$  for all but a countable set of  $t > 0$ , by virtue of the fact that  $\mu\{x \in X : f(x) > 0\} < \infty$ .

Now let  $U_j = f^{-1}\{(t_{j-1}, t_j)\}$ ,  $j = 1, \dots, N$ . (Notice that then the  $U_j$  are pairwise disjoint and each  $U_j \subset K$ , where  $K$ , compact, is the support of  $f$ .) Now by the definition (1) we can find  $g_j \in \mathcal{K}_+$  such that  $g_j \leq 1$ ,  $\text{support } g_j \subset U_j$ , and  $\lambda(g_j) \geq \mu(U_j) - \varepsilon/N$ . Also for any compact  $K_j \subset U_j$  we can construct a function  $h_j \in \mathcal{K}_+$  with  $h_j \equiv 1$  in a neighborhood of  $K_j \cup \text{support } g_j$ ,  $\text{support } h_j \subset U_j$ , and  $h_j \leq 1$  everywhere. Then  $h_j \geq g_j$ ,  $h_j \leq 1$  everywhere and  $\text{support } h_j$  is a compact subset of  $U_j$  and so

$$(5) \quad \mu(U_j) - \varepsilon/N \leq \lambda(g_j) \leq \lambda(h_j) \leq \mu(U_j), \quad j = 1, \dots, N.$$

Since  $\mu$  is a Radon measure, we can in fact choose the compact  $K_j \subset U_j$  such that  $\mu(U_j \setminus K_j) < \varepsilon/N$ . Then, because  $\{x : (f - f \sum_{j=1}^N h_j)(x) > 0\} \subset \cup(U_j \setminus K_j)$ , by (4) we have

$$(6) \quad \lambda(f - f \sum_{j=1}^N h_j) \leq \varepsilon \sup f.$$

Then by using (5), (6) and the linearity of  $\lambda$  (together with the fact  $t_{j-1}h_j \leq fh_j \leq t_jh_j$ ) for each  $j = 1, \dots, N$ ), we see that

$$\begin{aligned} \sum_{j=1}^N t_{j-1} \mu(U_j) - \varepsilon \sup f &\leq \lambda(f \sum_{j=1}^N h_j) \leq \lambda(f) \leq \lambda(f \sum_{j=1}^N h_j) + \varepsilon \sup f \\ &\leq \sum_{j=1}^N t_j \mu(U_j) + \varepsilon \sup f. \end{aligned}$$

Since trivially

$$\sum_{j=1}^N t_{j-1} \mu(U_j) \leq \int_X f d\mu \leq \sum_{j=1}^N t_j \mu(U_j),$$

we then have

$$\begin{aligned} -\varepsilon(\mu(K) + \sup f) &\leq -\sum_{j=1}^N (t_j - t_{j-1}) \mu(U_j) - \varepsilon \sup f \\ &\leq \int_X f d\mu - \lambda(f) \\ &\leq \sum_{j=1}^N (t_j - t_{j-1}) \mu(U_j) + \varepsilon \sup f \leq \varepsilon(\mu(K) + \sup f), \end{aligned}$$

where  $K = \text{support } f$ . This completes the proof of 5.12.  $\square$

We can now state the Riesz Representation Theorem. In the statement,  $C_c(X, H)$  will denote the set of vector functions  $f : X \rightarrow H$  which are continuous and which have compact support, where  $H$  is a given finite dimensional real Hilbert space with inner product  $\langle \cdot, \cdot \rangle$  and inner product norm  $|| \cdot ||$ .

**5.14 Theorem (Riesz Representation Theorem.)** *Suppose  $X$  is a locally compact Hausdorff space, and  $L : C_c(X, H) \rightarrow \mathbb{R}$  is linear with*

$$\sup_{f \in C_c(X, H), |f| \leq 1, \text{support } f \subset K} L(f) < \infty \text{ whenever } K \subset X \text{ is compact.}$$

*Then there is a Radon measure  $\mu$  on  $X$  and Borel measurable  $v : X \rightarrow H$  with  $|v| = 1$   $\mu$ -a.e. on  $X$ , and*

$$L(f) = \int_X \langle f, v \rangle d\mu \text{ for any } f \in C_c(X, H).$$

**Proof:** By using an orthonormal basis for  $H$ , it suffices to prove the theorem with  $H = \mathbb{R}^n$ . We first define

$$\lambda(f) = \sup_{\omega \in C_c(X, \mathbb{R}^n), |\omega| \leq f} L(\omega)$$

for any  $f \in \mathcal{K}_+$ . We claim that  $\lambda$  has the linearity properties of the lemma. Indeed it is clear that  $\lambda(cf) = c\lambda(f)$  for any constant  $c \geq 0$  and any  $f \in \mathcal{K}_+$ . Now let  $f, g \in \mathcal{K}_+$ , and notice that if  $\omega_1, \omega_2 \in C_c(X, \mathbb{R}^n)$  with  $|\omega_1| \leq f$  and  $|\omega_2| \leq g$ , then  $|\omega_1 + \omega_2| \leq f + g$  and hence  $\lambda(f + g) \geq L(\omega_1) + L(\omega_2)$ . Taking sup over all such  $\omega_1, \omega_2$  we then have  $\lambda(f + g) \geq \lambda(f) + \lambda(g)$ . To prove the reverse inequality we let  $\omega \in C_c(X, \mathbb{R}^n)$  with  $|\omega| \leq f + g$ , and define

$$\omega_1 = \begin{cases} \frac{f}{f+g} \omega & \text{if } f + g > 0 \\ 0 & \text{if } f + g = 0, \end{cases} \quad \omega_2 = \begin{cases} \frac{g}{f+g} \omega & \text{if } f + g > 0 \\ 0 & \text{if } f + g = 0. \end{cases}$$

Then  $\omega_1 + \omega_2 = \omega$ ,  $|\omega_1| \leq f$ ,  $|\omega_2| \leq g$  and it is readily checked that  $\omega_1, \omega_2 \in C_c(X, \mathbb{R}^n)$ . Then  $L(\omega) = L(\omega_1) + L(\omega_2) \leq \lambda(f) + \lambda(g)$ , and hence taking sup over all such  $\omega$  we have  $\lambda(f + g) \leq \lambda(f) + \lambda(g)$ . Therefore we have  $\lambda(f + g) = \lambda(f) + \lambda(g)$  as claimed. Thus  $\lambda$  satisfies the conditions of the lemma, hence there is a Radon measure  $\mu$  on  $X$  such that

$$\lambda(f) = \int_X f d\mu, \quad f \in \mathcal{K}_+, \quad j = 1, \dots, n.$$

That is, we have

$$(\ddagger) \quad \sup_{\omega \in C_c(X, \mathbb{R}^n), |\omega| \leq f} L(\omega) = \int_X f d\mu, \quad f \in \mathcal{K}_+.$$

Thus if  $j \in \{1, \dots, n\}$  we have in particular (since  $|fe_j| = |f| \in \mathcal{K}_+$  for any  $f \in C_c(X, \mathbb{R})$ ) that

$$|L(fe_j)| \leq \int_X |f| d\mu \equiv \|f\|_{L^1(\mu)} \quad \forall f \in C_c(X, \mathbb{R}).$$

Thus  $L_j(f) \equiv L(fe_j)$  extends to a bounded linear functional on  $L^1(\mu)$ , and hence by the Riesz representation theorem for  $L^1(\mu)$  we know that there is a bounded  $\mu$ -measurable function  $v_j$  such that

$$L(fe_j) = \int_X f v_j d\mu, \quad f \in C_c(X, \mathbb{R}).$$

Since any  $f = (f_1, \dots, f_n)$  can be expressed as  $f = \sum_{j=1}^n f_j e_j$ , we thus deduce

$$(*) \quad L(f) = \int_X f \cdot v d\mu, \quad f \in C_c(X, \mathbb{R}^n),$$

where  $v = (v_1, \dots, v_n)$ . Then it only remains to check that  $|v| = 1$   $\mu$ -a.e. To see this, first note that by using the Cauchy-Schwarz inequality in the integral on the right of  $(*)$  we have for any  $f \in \mathcal{K}_+$  that

$$(i) \quad \sup_{|g| \leq f, g \in C_c(X, \mathbb{R}^n)} |L(g)| \leq \int_X f |v| d\mu.$$

On the other hand, we know (since  $C_c(X, \mathbb{R}^n)$  is dense in  $L^1(\mu)$ ), we can find a sequence  $g_k \in C_c(X, \mathbb{R}^n)$  such that  $\lim \int_X |g_k - \widehat{v}| = 0$ , where  $\widehat{v}$  is  $|v|^{-1}v$  at points where  $v \neq 0$  and  $\widehat{v} = 0$  at all other points. Then of course  $\lim \int_X |\widehat{g}_k - \widehat{v}| = 0$  with  $|\widehat{g}_k| \leq 1$ , provided we define  $\widehat{g}_k = R(g_k)$ , with  $R(y) = |y|^{-1}y$  if  $|y| > 1$  and  $R(y) = y$  if  $|y| \leq 1$ , because  $|R(y) - v| \leq |y - v|$  for any  $y, v \in \mathbb{R}^n$  with  $|v| = 1$ . Thus we deduce that actually equality holds in (i). On the other hand by  $(\ddagger)$  for any  $f \in \mathcal{K}_+$  we have that the left side of (i) is  $\int_X f d\mu$ . Thus finally  $\int_X f d\mu = \int_X f |v| d\mu$ , and this evidently implies  $|v| = 1$   $\mu$ -a.e., again using the density of  $C_c(X, \mathbb{R})$  in  $L^1(\mu)$ .  $\square$

Using the Riesz Theorem 5.12 we can deduce the following compactness theorem for Radon measures:

**5.15 Theorem (Compactness Theorem for Radon Measures.)** *Suppose  $\{\mu_k\}$  is a sequence of Radon measures on the locally compact,  $\sigma$ -compact Hausdorff space  $X$  with the property  $\sup_k \mu_k(K) < \infty$  for each compact  $K \subset X$ . Then there is a subsequence  $\{\mu_{k'}\}$  which converges to a Radon measure  $\mu$  on  $X$  in the sense that*

$$\lim \mu_{k'}(f) = \mu(f) \text{ for each } f \in \mathcal{K}(X),$$

where  $\mathcal{K}(X)$  denotes the set of continuous functions  $f : X \rightarrow \mathbb{R}$  with compact support on  $X$  and where we use the notation

$$\mu(f) = \int_X f \, d\mu, \quad f \in \mathcal{K}(X).$$

**Proof:** Let  $K_1, K_2, \dots$  be an increasing sequence of compact sets with  $X = \cup_j K_j$  and let  $F_{j,k} : C(K_j) \rightarrow \mathbb{R}$  be defined by  $F_{j,k}(f) = \int_{K_j} f \, d\mu_k$ ,  $k = 1, 2, \dots$ . By the Banach-Alaoglu theorem (which guarantees weak\* compactness of the closed unit ball in the Banach space of bounded linear functionals on  $C(K_j)$ ) there is a subsequence  $F_{j,k'}$  and a non-negative bounded functional  $F_j : C(K_j) \rightarrow \mathbb{R}$  with  $F_{j,k'}(f) \rightarrow F_j(f)$  for each  $f \in C(K_j)$ . By choosing the subsequences successively and taking a diagonal sequence, we then get a subsequence  $\mu_{k'}$  and a non-negative linear  $F : \mathcal{K}(X) \rightarrow \mathbb{R}$  with  $\int_X f \, d\mu_{k'} \rightarrow F(f)$  for each  $f \in \mathcal{K}(X)$ , where  $F(f) = F_j(f|_{K_j})$  whenever  $\text{spt } f \subset K_j$ . (Notice that this is unambiguous because if  $\text{spt } f \subset K_j$  and  $\ell > j$  then  $F_\ell(f|_{K_\ell}) = F_j(f|_{K_j})$  by construction.) Then by applying Theorem 5.12 we have a Radon measure  $\mu$  on  $X$  such that  $F(f) = \int_X f \, d\mu$  for each  $f \in \mathcal{K}(X)$ , and so  $\int_X f \, d\mu_{k'} \rightarrow \int_X f \, d\mu$  for each  $f \in \mathcal{K}(X)$ .  $\square$

## CHAPTER 1 PROBLEMS

1.1 (i) If  $A_1, A_2$  are non-empty compact subsets of  $\mathbb{R}$ , prove  $\exists a_j \in A_j, j = 1, 2$ , such that  $|a_2 - a_1| \geq \frac{1}{2}(\mathcal{L}^1(A_1) + \mathcal{L}^1(A_2))$ . ( $\mathcal{L}^1$  denotes Lebesgue measure on  $\mathbb{R}$ .)

(ii) Let the notation be as in the proof of Theorem 2.7 and  $j \in \{1, \dots, n\}$ . By applying the result of (i) above to the sets  $A_1 = \pi(\ell_j(\xi_1) \cap A)$ ,  $A_2 = \pi(\ell_j(\xi_2) \cap A)$ , prove that if  $A \subset \mathbb{R}^n$  is compact then  $\text{diam}(\mathcal{S}_j(A)) \leq \text{diam}(A)$ .

(iii) If  $A$  is compact, prove that  $\mathcal{L}^1(\pi(A \cap \ell_j(\xi)))$  (where  $\ell_j(\xi), \pi$  are as in the proof of Theorem 2.7) is an upper semi-continuous function of  $\xi$  if  $\xi$  is restricted to lie in the  $j$ -th coordinate hyperplane  $x^j = 0$ .

Hint: For  $\varepsilon > 0$  we can select open  $U \subset \mathbb{R}$  with  $\pi(A \cap \ell_j(\xi)) \subset U$  and  $\mathcal{L}^1(U) \leq \mathcal{L}^1(\pi(A \cap \ell_j(\xi))) + \varepsilon$ .

(iv) Using the result of (iii) prove that  $A$  compact  $\Rightarrow \mathcal{S}_j(A)$  is compact (where  $\mathcal{S}_j(A)$  is the Steiner symmetrization of  $A$  as in the proof of Theorem 2.7).

1.2 (Borel regularity of Hausdorff measure.) Let  $\mathcal{H}^m$  be  $m$ -dimensional Hausdorff (outer) measure on a metric space  $X, d$ . Prove that  $\mathcal{H}^m$  is Borel regular.

Note: As mentioned in lecture,  $\mathcal{H}^m$  evidently has the property that  $\mathcal{H}^m(A \cup B) = \mathcal{H}^m(A) + \mathcal{H}^m(B)$  whenever  $d(A, B) > 0$ , so all Borel sets are  $\mathcal{H}^m$ -measurable by the Caratheodory theorem 1.15 of the text; thus for this question you merely need to check (directly from the definition of  $\mathcal{H}^m$ ) that for every set  $A \subset X$  there is a Borel set  $B \supset A$  with  $\mathcal{H}^m(B) = \mathcal{H}^m(A)$ .

1.3 Let  $X, d$  be a metric space and let  $\mu$  be a Borel-regular outer measure on  $X$  which is finite on each ball  $B_\rho(x) \subset X$ . In §3 we proved that the lower density  $\Theta_*^n(\mu, x) (= \liminf_{\rho \downarrow 0} \frac{\mu(B_\rho(x))}{\omega_n \rho^n})$  is Borel measurable on  $X$ .

With a similar argument, prove that  $\Theta^{*n}(\mu, x) = \limsup_{\rho \downarrow 0} \frac{\mu(B_\rho(x))}{\omega_n \rho^n}$  is also Borel measurable.

Hint: Start by proving that  $\limsup_{\rho \downarrow 0} \frac{\mu(B_\rho(x))}{\omega_n \rho^n} = \limsup_{\rho \downarrow 0} \frac{\mu(\check{B}_\rho(x))}{\omega_n \rho^n}$ , where  $\check{B}_\rho(x)$  denotes the open ball of radius  $\rho$  and center  $x$ .

1.4 Suppose  $X$  is any metric space and  $\mu$  is an open  $\sigma$ -finite (as in 1.21) Borel regular outer measure with the Symmetric Vitali property. (For example this is true by Corollary 3.7 if  $X = \mathbb{R}^n$  and  $\mu = \mathcal{L}^n$ .)  $f$  is said to be approximately continuous at  $x \in X$  with respect to  $\mu$  if  $\mu(B_\rho(x)) > 0$  for each  $\rho > 0$  and

$$\lim_{\rho \downarrow 0} (\mu(B_\rho(x)))^{-1} \mu(\{y \in B_\rho(x) : |f(y) - f(x)| \geq \varepsilon\}) = 0 \quad \forall \varepsilon > 0.$$

Prove that if  $f$  is  $\mu$ -measurable on  $X$  then  $f$  is approximately continuous at  $\mu$ -a.e.  $x \in X$ .

Hint: Use Lusin's Theorem 1.24 and the Upper Density Theorem 4.8.

1.5 If  $U$  is any bounded open set in  $\mathbb{R}^n$  and  $\delta > 0$ , prove there is closed balls  $B_{\rho_j}(x_j) \subset U$  with  $\rho_j < \delta \forall j$ ,  $B_{\rho_i}(x_i) \cap B_{\rho_j}(x_j) = \emptyset \forall i \neq j$ , and  $\mathcal{L}^n(U \setminus (\cup_{j=1}^\infty B_{\rho_j}(x_j))) = 0$ .

Hint: Using the cubes  $2^{-i}([j_1, j_1 + 1] \times \cdots \times [j_n, j_n + 1])$ ,  $j_1, \dots, j_n \in \mathbb{Z}$ ,  $i \in \mathbb{Z}_+$ , decompose  $U$  as a union  $\cup_{j=1}^{\infty} C_j$  of closed cubes  $C_j$  of diameter  $< \delta$  and with pairwise disjoint interiors, and for each  $j$  select a ball  $B_j \subset \text{interior } C_j$  with  $\text{diam } B_j > \text{edge-length of } C_j / 2$ . Then  $\mathcal{L}^n(C_j \setminus B_j) < (1 - \theta_n)\mathcal{L}^n(C_j)$ ,  $\theta_n = \omega_n/4^n$ , and hence  $\mathcal{L}^n(U \setminus (\cup_{j=1}^{\infty} B_j)) < (1 - \theta_n)\mathcal{L}^n(U)$ , so  $\mathcal{L}^n(U \setminus (\cup_{j=1}^N B_j)) < (1 - \theta_n)\mathcal{L}^n(U)$  for suitably large  $N$ . Since  $U \setminus (\cup_{j=1}^N B_j)$  is open, we can repeat this process, starting with  $U \setminus (\cup_{j=1}^N B_j)$  in place of  $U$ .



# Chapter 2

## Some Further Preliminaries from Analysis

§ 1	Lipschitz Functions .....	43
§ 2	BV Functions .....	52
§ 3	The Area Formula .....	57
§ 4	Submanifolds of $\mathbb{R}^{n+k}$ .....	61
§ 5	First Variation of a Submanifold .....	70
§ 6	Second Variation .....	73
§ 7	Co-Area Formula and $C^1$ Sard Theorem .....	75

Here we develop the necessary further analytical background material needed for later developments. In particular we prove some basic results about Lipschitz and BV functions, and we also present the basic facts concerning  $C^k$  submanifolds of Euclidean space. We also discuss the area and co-area formulae and first and second variation formulae for  $C^2$  submanifolds of Euclidean space. These latter topics will be discussed in a much more general context later.

### 1 Lipschitz Functions

If  $X$  is a metric space with metric  $d$ , recall that a function  $f : X \rightarrow \mathbb{R}$  is said to be *Lipschitz* if there is  $L < \infty$  such that

1.1 
$$|f(x) - f(y)| \leq L d(x, y) \quad \forall x, y \in X.$$

$\text{Lip } f$  denotes the least such constant  $L$ .

First we have the following basic extension theorem.

**1.2 Theorem.** *If  $A$  is a non-empty subset of  $X$  and  $f : A \rightarrow \mathbb{R}$  is Lipschitz, then  $\exists \bar{f} : X \rightarrow \mathbb{R}$  with  $\text{Lip } \bar{f} = \text{Lip } f$ , and  $\bar{f} = f|_A$ . Also,  $\bar{f}$  can be chosen so that  $\sup_X |\bar{f}| = \sup_A |f|$ .*

**Proof:** With  $L = \text{Lip } f$ , we claim that

$$\bar{f}(x) = \inf_{z \in A} (f(z) + Ld(x, z)), \quad x \in X,$$

has the required properties, except possibly the requirement  $\sup_X |\bar{f}| = \sup_A |f|$ . To check this, first note that  $\bar{f}(x) > -\infty$  for each  $x \in X$ , because if  $x_0 \in A$  then  $f(z) + Ld(x, z) = f(x_0) + f(z) - f(x_0) + Ld(x, z) \geq f(x_0) + L(d(x, z) - d(x_0, z)) \geq f(x_0) - Ld(x, x_0)$  by the triangle inequality. Also, if  $x \in A$  then  $\bar{f}(x) - f(x) = \inf_{z \in A} (f(z) - f(x) + Ld(x, z)) \geq \inf_{z \in A} (-Ld(x, z) + Ld(x, z)) = 0$ , so  $\bar{f}(x) \geq f(x)$ , and of course the reverse inequality holds trivially. Hence  $\bar{f}(x) = f(x)$  for  $x \in A$ . So  $\bar{f}$  is well-defined as a map  $X \rightarrow \mathbb{R}$  and it agrees with  $f$  on  $A$ .

For any  $x_1, x_2 \in X$

$$\begin{aligned} \bar{f}(x_1) - \bar{f}(x_2) &= \sup_{z_2 \in A} \inf_{z_1 \in A} (f(z_1) + Ld(x_1, z_1) - f(z_2) - Ld(x_2, z_2)) \\ &\leq \sup_{z_2 \in A} (Ld(x_1, z_2) - Ld(x_2, z_2)) \leq Ld(x_1, x_2) \end{aligned}$$

and the reverse inequality holds by interchanging  $x_1, x_2$ .

Finally, observe that we can replace  $\bar{f}$  by its truncation  $\gamma(\bar{f})$ , where

$$\gamma(t) = \max\{\min\{t, \kappa\}, -\kappa\}, \quad \kappa = \sup_A |f|. \quad \square$$

**1.3 Remark:** Observe that the above proof has a geometric interpretation: the graph of the extension  $\bar{f}$  is obtained by taking the “lower envelope” (inf) of all the half-cones  $C_z = \{(x, y) \in X \times \mathbb{R} : y = f(z) + Ld(x, z)\}$ ; notice that  $C_z$  is a half-cone of slope  $L$  with vertex on the graph of the original function  $f$ .

Next we need the theorem of Rademacher concerning differentiability of Lipschitz functions on  $\mathbb{R}^n$ . (The proof given here is due to C.B. Morrey.)

**1.4 Theorem (Rademacher’s theorem.)** *If  $f$  is Lipschitz on  $\mathbb{R}^n$ , then  $f$  is differentiable  $\mathcal{L}^n$ -almost everywhere; that is, the gradient  $\nabla f(x) = (D_1 f(x), \dots, D_n f(x))$  exists and*

$$(*) \quad \lim_{y \rightarrow x} \frac{f(y) - f(x) - \nabla f(x) \cdot (y - x)}{|y - x|} = 0$$

for  $\mathcal{L}^n$ -a.e.  $x \in \mathbb{R}^n$ .

**Proof:** Let  $v \in \mathbb{S}^{n-1}$ , and whenever it exists let  $D_v f(x)$  denote the directional derivative  $\frac{d}{dt} f(x + tv)|_{t=0}$ . Since  $|\frac{f(y)-f(x)}{|y-x|}| \leq \text{Lip } f$  for  $y \neq x$  (so  $|D_v f| \leq \text{Lip } f$  whenever it exists) and we see that  $D_v f(x)$  exists precisely when the bounded functions

$$\limsup_{t \rightarrow 0} \frac{f(x + tv) - f(x)}{t}, \quad \liminf_{t \rightarrow 0} \frac{f(x + tv) - f(x)}{t}$$

coincide. Now  $\limsup_{t \rightarrow 0} \frac{f(x + tv) - f(x)}{t} = \lim_{j \rightarrow \infty} \sup_{0 < |t| < j^{-1}} \frac{f(x + tv) - f(x)}{t}$  which is Borel measurable because  $\sup_{0 < |t| < j^{-1}} \frac{f(x + tv) - f(x)}{t}$  is lower semi-continuous, and hence Borel measurable, for each  $j$ . Similarly  $\liminf_{t \rightarrow 0} \frac{f(x + tv) - f(x)}{t}$  is Borel measurable, so the set  $A_v = \{x \in \mathbb{R}^n : D_v f \text{ does not exist}\}$  is Borel measurable and hence  $\mathcal{L}^n$ -measurable. However  $\varphi(t) = f(x + tv)$  is an absolutely continuous function of  $t \in \mathbb{R}$  for any fixed  $x$  and  $v$ , and hence is differentiable for almost all  $t$ . Thus  $A_v$  intersects every line  $L$  which is parallel to  $v$  in a set of  $\mathcal{H}^1$  measure zero and hence by Fubini's theorem the Borel set  $A_v$  has  $\mathcal{L}^n$ -measure zero for each  $v$ . That is, for each  $v \in \mathbb{S}^{n-1}$ ,

$$(1) \quad D_v f(x) \text{ exists } \mathcal{L}^n\text{-a.e. } x \in \mathbb{R}^n.$$

Now take any  $C_c^\infty(\mathbb{R}^n)$  function  $\zeta$  and note that for any  $h > 0$

$$(2) \quad \int_{\mathbb{R}^n} \frac{f(x + hv) - f(x)}{h} \zeta(x) d\mathcal{L}^n(x) = - \int_{\mathbb{R}^n} \frac{\zeta(x) - \zeta(x - hv)}{h} f(x) d\mathcal{L}^n(x)$$

(by the change of variable  $z = x + hv$  in the first part of the integral on the left). Using the dominated convergence theorem and (1) we then have

$$(3) \quad \begin{aligned} \int D_v f \zeta &= - \int f D_v \zeta = - \int f v \cdot \nabla \zeta \\ &= - \sum_{j=1}^n v^j \int f D_j \zeta = + \sum_{j=1}^n v^j \int \zeta D_j f = \int \zeta v \cdot \nabla f, \end{aligned}$$

where  $\nabla f$  is the gradient of  $f$  (i.e.  $\nabla f = (D_1 f, \dots, D_n f)$ ), all integrals are with respect to Lebesgue measure on  $\mathbb{R}^n$ , and we have used Fubini's theorem and the absolute continuity of  $f$  on lines to justify the integration by parts. Since  $\zeta$  is arbitrary in (3) we then have, for each  $v \in \mathbb{S}^{n-1}$ ,

$$(4) \quad \nabla f(x), D_v f(x) \text{ exist and } D_v f(x) = v \cdot \nabla f(x) \text{ for } \mathcal{L}^n\text{-a.e. } x \in \mathbb{R}^n.$$

Of course at such points  $x$  we also have

$$(5) \quad |\nabla f(x)| \leq L.$$

Now let  $v_1, v_2, \dots$  be a countable dense subset of  $\mathbb{S}^{n-1}$ , and let

$$A_k = \{x : \nabla f(x), D_{v_k} f(x) \text{ exist and } D_{v_k} f(x) = v_k \cdot \nabla f(x)\}.$$

Then  $A = \bigcap_{k=1}^{\infty} A_k$  we have by (4) that

$$(6) \quad \mathcal{L}^n(\mathbb{R}^n \setminus A) = 0, \quad D_{v_k} f(x) = v_k \cdot \nabla f(x) \quad \forall x \in A, \quad k = 1, 2, \dots$$

Using this, we are now going to prove that  $f$  is differentiable at each point  $x$  of  $A$ . To see this, for any  $x \in A$ ,  $v \in \mathbb{S}^{n-1}$  and  $h > 0$  define

$$(7) \quad Q(x, v, h) = \frac{f(x + hv) - f(x)}{h} - v \cdot \nabla f(x),$$

so by (6)

$$(8) \quad \lim_{h \rightarrow 0} Q(x, v_j, h) = 0, \quad x \in A, \quad j = 1, 2, \dots$$

Now for any given  $\varepsilon > 0$ , select  $P$  large enough so that

$$(9) \quad S^{n-1} \subset \bigcup_{i=1}^P B_{\varepsilon}(v_i),$$

and for each  $i = 1, \dots, P$  use (8) to choose  $\delta_i > 0$  so that

$$(10) \quad 0 < |h| < \delta_i \Rightarrow |Q(x, v_i, h)| < \varepsilon.$$

By (9), for any  $v \in S^{n-1}$  we can select  $i \in \{1, \dots, P\}$  with  $|v - v_i| < \varepsilon$ , and hence by (10)

$$(11) \quad \begin{aligned} |Q(x, h, v)| &\leq |Q(x, v, h) - Q(x, v_i, h)| + |Q(x, v_i, h)| \\ &\leq |h|^{-1} |f(x + hv) - f(x + hv_i)| + |v - v_i| |\nabla f(x)| + |Q(x, v_i, h)| \\ &< (2L + 1)\varepsilon \text{ for all } 0 < |h| < \delta = \min\{\delta_1, \dots, \delta_P\} \end{aligned}$$

by (5). Thus  $v \in S^{n-1}$  and  $0 < |h| < \delta \Rightarrow |Q(x, h, v)| < (2L + 1)\varepsilon$ , hence  $f$  is differentiable at  $x$ .  $\square$

We shall need the following  $C^1$  approximation theorem for Lipschitz functions in our discussion of rectifiable sets in the next chapter.

**1.5 Theorem. ( $C^1$  Approximation Theorem.)** *Suppose  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is Lipschitz. Then for each  $\varepsilon > 0$  there is a  $C^1(\mathbb{R}^n)$  function  $g$  with*

$$\mathcal{L}^n(\{x : f(x) \neq g(x)\} \cup \{x : \nabla f(x) \neq \nabla g(x)\}) < \varepsilon.$$

Before we begin the proof of 1.5 we need to recall Whitney's extension theorem for  $C^1$  functions:

**1.6 Theorem (Whitney Extension Theorem.)** *If  $C \subset \mathbb{R}^n$  is closed and if  $h : C \rightarrow \mathbb{R}$  and  $v : C \rightarrow \mathbb{R}^n$  are continuous, and if for each compact  $K \subset C$*

$$(\ddagger) \quad \lim_{y \rightarrow x, y \in K} R(x, y) = 0 \text{ uniformly for } x \in K,$$

where

$$R(x, y) = \frac{h(y) - h(x) - v(x) \cdot (y - x)}{|x - y|},$$

then there is a  $C^1$  function  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  such that  $g = h$  and  $\nabla g = v$  on  $C$ .

(For the proof see for example [EG92] or [Fed69]; for the case  $n = 1$ , see Remark 1.7(2) below.)

**1.7 Remarks: (1)** The hypothesis 1.6 (‡) above cannot be weakened to the requirement that

$$\lim_{y \rightarrow x, y \in C} R(x, y) = 0, \quad x \in C.$$

For instance we have the example (for  $n = 1$ ) when  $C = \{0\} \cup (\cup_{k=1}^{\infty} \{1/k\})$  and  $h(0) = 0$ ,  $h(1/k) = (-1)^k / k^{3/2}$ ,  $v \equiv 0$ . Evidently in this case we do have  $\lim_{y \rightarrow x, y \in C} R(x, y) = 0 \forall x \in C$ , but there is no  $C^1$  extension because

$$\frac{|h(1/k) - h(1/(k+1))|}{(1/k - 1/(k+1))} \rightarrow \infty \text{ as } k \rightarrow \infty.$$

In fact the condition 1.6 (‡) is *equivalent* to the existence of a  $C^1$  extension  $g$  of  $h$  with  $\nabla g = v$  on  $C$ . Indeed if  $g$  is such an extension and if  $K \subset C$  is compact then for  $x, y \in K$  we have

$$\begin{aligned} R(x, y) &= |y - x|^{-1} (h(y) - h(x) - v(x) \cdot (y - x) = g(y) - g(x) - \nabla g(x) \cdot (y - x)) \\ &= \int_0^1 \frac{d}{dt} g(x + t(y - x)) dt - \nabla g(x) \cdot (y - x) / |y - x| \\ &= \int_0^1 (\nabla g(x + t(y - x)) - \nabla g(x)) \cdot (y - x) / |y - x| dt \end{aligned}$$

and, since  $\nabla g$  is uniformly continuous on the convex hull of  $K$ , we do indeed have 1.6 (‡).

(2) In the case  $n = 1$ , the Whitney Extension Theorem 1.6 above has a simple direct proof. Namely in this case define

$$R(x, y) = \frac{h(y) - h(x)}{y - x} - v(x)$$

and note that the hypothesis 1.6 (‡) guarantees that for each compact subset  $K$  of  $C$  we have a function  $\varepsilon_K$  with  $\varepsilon_K(t) \downarrow 0$  as  $t \downarrow 0$ , and

$$|R(x, y)| \leq \varepsilon_K(|x - y|) \quad \forall x, y \in K,$$

and of course since  $v$  is uniformly continuous on  $K$  we can suppose that  $\varepsilon_K$  is chosen so that

$$(‡) \quad |v(x) - v(y)| \leq \varepsilon_K(|x - y|) \quad \forall x, y \in K.$$

Also  $\mathbb{R} \setminus C$  is a countable disjoint union of open intervals  $I_1, I_2, \dots$ . If  $I_j = (a, b)$ , we then select  $g_j \in C^1([a, b])$  as follows

$$g_j(a) = h(a), \quad g_j(b) = h(b), \quad g'_j(a) = v(a), \quad g'_j(b) = v(b)$$

and

$$\sup_{x \in I_j} |g'_j(x) - v(a)| \leq 2\varepsilon_K(b-a), \quad K = [a-1, b+1] \cap C.$$

This is possible by  $(\ddagger)$ , with  $(x, y) = (a, b)$ . One now defines  $g(x) = g_j(x) \quad \forall x \in I_j$ ,  $j = 1, 2, \dots$ , and  $g(x) = h(x) \quad \forall x \in C$ . It is then easy to check  $g \in C^1(\mathbb{R})$  and  $g' = v$  on  $C$ .

**Proof of Theorem 1.5:** By Rademacher's Theorem  $\nabla f$  exists and  $f$  is differentiable  $\mathcal{L}^n$ -a.e. on  $\mathbb{R}^n$ . Thus with  $R(x, y) = |y-x|^{-1}|f(y) - f(x) - \nabla f(x) \cdot (y-x)|$  we have

$$\eta_k(x) = \sup_{0 < |y-x| < 1/k} |R(x, y)| \downarrow 0 \text{ for } \mathcal{L}^n\text{-a.e. } x \in \mathbb{R}^n.$$

Hence by Egoroff's Theorem (1.12 of Ch.1), applied to the finite measure annular regions  $B_j(0) \setminus B_{j-1}(0)$ ,  $j = 1, 2, \dots$ , there is an  $\mathcal{L}^n$ -measurable set  $E_j \subset B_j(0) \setminus B_{j-1}(0)$  such that  $\mathcal{L}^n((B_j(0) \setminus B_{j-1}(0)) \setminus E_j) < \varepsilon/2^{j+1}$  and  $\eta_k$  converges uniformly to zero  $E_j$ .

By Lusin's Theorem 1.24 of Ch.1 there is a compact set  $C_j \subset E_j$  such that  $\nabla f|_{C_j}$  is continuous and  $\mathcal{L}^n(E_j \setminus C_j) < \varepsilon/2^{j+1}$ . Thus with  $C = \bigcup_{j=1}^{\infty} C_j$  we have  $C$  closed,  $\nabla f|_C$  continuous,  $\mathcal{L}^n(\mathbb{R}^n \setminus C) < 2 \sum_{j=1}^{\infty} 2^{-j-1} \varepsilon = \varepsilon$ , and  $\eta_k$  converges uniformly to zero on each bounded subset of  $C$ . Hence we can apply Whitney's Theorem 1.6 with  $h = f$  and  $v = \nabla f(x)$  in order to give the required  $C^1$  function  $g$ .  $\square$

Next we establish some basic facts about Hausdorff measure of Lipschitz images. In this direction we first observe that if  $X, Y$  are metric spaces, if  $A \subset X$  and if  $f : A \rightarrow Y$  is Lipschitz then, for each  $m \geq 0$  ( $m$  need not be an integer),

$$1.8 \quad \mathcal{H}^m(f(A)) \leq (\text{Lip } f)^m \mathcal{H}^m(A).$$

Of course this is trivial if  $m = 0$ , while if  $\delta, m > 0$  and if  $C_1, C_2, \dots$  are chosen with  $A \subset \bigcup_j C_j$  and  $\text{diam } C_j < \delta$  for each  $j$ , then  $f(A) \subset \bigcup_j f(C_j)$  and  $\text{diam}(f(C_j)) \leq (\text{Lip } f)\delta < (1 + \text{Lip } f)\delta$ . Hence

$$\mathcal{H}_{(1+\text{Lip } f)\delta}^m(f(A)) \leq \sum_j \omega_m(\text{diam } f(C_j)/2)^m \leq (\text{Lip } f)^m \sum_j \omega_m(\text{diam } C_j/2)^m,$$

and taking inf over all such collections  $\{C_j\}$  and then letting  $\delta \downarrow 0$  we obtain 1.8 as claimed.

The following theorem refines 1.8 in case  $\mathcal{H}^m(A) < \infty$  and  $X$  is  $\sigma$ -compact (i.e. in case there are compact  $K_1, K_2, \dots$  with  $X = \bigcup_j K_j$ ).

**1.9 Theorem.** Suppose  $X, Y$  are metric spaces,  $X$  is  $\sigma$ -compact,  $A \subset X$  is  $\mathcal{H}^m$ -measurable,  $\mathcal{H}^m(A) < \infty$  and  $f : A \rightarrow Y$  is Lipschitz, and let  $\mathcal{N}(f, y) = \mathcal{H}^0(f^{-1}y)$  (i.e.  $\mathcal{N}(f, y)$  is the multiplicity function, counting the number of points, possibly  $\infty$ , in the preimage  $f^{-1}y$ ). Then

(i)  $f(A)$  is  $\mathcal{H}^m$ -measurable.

(ii)  $\mathcal{N}(f, y)$  is an  $\mathcal{H}^m$ -measurable function of  $y \in Y$  with

$$\int_Y \mathcal{N}(f, y) d\mathcal{H}^m \leq (\text{Lip } f)^m \mathcal{H}^m(A).$$

**Proof:** Since  $A$  is  $\mathcal{H}^m$ -measurable and  $\mathcal{H}^m(A) < \infty$  we can use the regularity property 1.22(2) of Ch. 1 together with the  $\sigma$ -compactness of  $X$  to find a sequence  $K_1, K_2, \dots$  of compact sets in  $X$  with  $K_j \subset A$  for each  $j$  and  $\mathcal{H}^m(A \setminus (\cup_j K_j)) = 0$ . Then  $\mathcal{H}^m(f(A \setminus (\cup_j K_j))) = 0$  by 1.8, so  $f(A) = f(A \setminus (\cup_j K_j)) \cup (\cup_j f(K_j))$  is the union of a set of  $\mathcal{H}^m$ -measure zero and countably many compact (hence Borel) sets  $f(K_j)$ , so  $f(A)$  is  $\mathcal{H}^m$ -measurable as claimed. This completes the proof of (i).

To prove (ii) observe that, by the  $\sigma$ -compactness of  $X$ , for each  $i = 1, 2, \dots$  we can partition  $A$  into a disjoint union  $\cup_{j=1}^{\infty} A_{ij}$  where each  $A_{ij}$  is  $\mathcal{H}^m$ -measurable and  $\text{diam}(A_{ij}) < 1/i$ ; furthermore we can do this inductively, partitioning each  $A_{ik}$  to give the new sets  $A_{i+1,j}$ , so that each of the sets  $A_{i+1,j}$  is contained in one of the  $A_{ik}$ . Observe that then  $\sum_j \chi_{f(A_{ij})}$  is a non-negative function which is  $\mathcal{H}^m$ -measurable by (i) above and which increases pointwise (at every point  $y \in Y$ ) to  $\mathcal{N}(f, y)$ , and so  $\mathcal{N}(f, y)$  is  $\mathcal{H}^m$ -measurable. Also, by the monotone convergence theorem,

$$\int_Y \mathcal{N}(f, y) d\mathcal{H}^m(y) = \lim_{i \rightarrow \infty} \int_Y \sum_j \chi_{f(A_{ij})} d\mathcal{H}^m = \lim_{i \rightarrow \infty} \sum_j \mathcal{H}^m(f(A_{ij})),$$

and

$$\sum_j \mathcal{H}^m(f(A_{ij})) \leq (\text{Lip } f)^m \sum_j \mathcal{H}^m(A_{ij}) = (\text{Lip } f)^m \mathcal{H}^m(A)$$

by 1.8.  $\square$

Next, in the case when  $m \in \{1, 2, \dots\}$ , we want to extend the inequality of Theorem 1.9(ii) to the case when the  $k$ -dimensional Hausdorff measure of  $f^{-1}y$  (instead of  $\mathcal{H}^0(f^{-1}y)$ ) appears on the left. For this we assume for convenience that  $Y = \mathbb{R}^m$  (more general cases, e.g. when  $Y$  is a metric space such that each closed ball is compact, are discussed in [Fed69, 10.2.25], but the case  $Y = \mathbb{R}^m$  is adequate for the subsequent development here, and furthermore the proof is relatively elementary in this case).

**1.10 Theorem.** Suppose  $X$  is a  $\sigma$ -compact metric space,  $m \in \{1, 2, \dots\}$ ,  $k > 0$  ( $k$  need not be an integer),  $A \subset X$  is  $\mathcal{H}^{m+k}$ -measurable and  $\mathcal{H}^{m+k}(A) < \infty$ , and  $f : A \rightarrow \mathbb{R}^m$  is

*Lipschitz.* Then  $\mathcal{H}^k(f^{-1}y)$  is an  $\mathcal{L}^m$ -measurable function of  $y \in \mathbb{R}^m$  and

$$\int_{\mathbb{R}^m} \mathcal{H}^k(f^{-1}y) d\mathcal{L}^m(y) \leq \frac{\omega_m \omega_k}{\omega_{m+k}} (\text{Lip } f)^m \mathcal{H}^{m+k}(A).$$

In particular, for each  $R > 0$ ,  $\mathcal{H}^k(f^{-1}y) \leq R$  except possibly for an  $\mathcal{L}^m$ -measurable set  $E$  with  $\mathcal{L}^m(E) \leq \frac{\omega_m \omega_k}{\omega_{m+k}} (\text{Lip } f)^m \mathcal{H}^{m+k}(A)/R$ .

**1.11 Remark:** At one step in the proof below we are going to use the *upper Lebesgue integral*  $\int_{\mathbb{R}^m}^* f d\mathcal{L}^m$  of a not necessarily measurable function  $f : \mathbb{R}^m \rightarrow [0, \infty]$ . This is defined by

$$\int_{\mathbb{R}^m}^* f d\mathcal{L}^m = \inf_{\psi \geq f, \psi \text{ measurable}} \int_{\mathbb{R}^m} \psi d\mathcal{L}^m.$$

Observe that then there is always a measurable function  $\psi_f$  which attains the inf; that is,  $\psi_f \geq f$  and

$$\int_{\mathbb{R}^m}^* f d\mathcal{L}^m = \int_{\mathbb{R}^m} \psi_f d\mathcal{L}^m,$$

and if  $\int_{\mathbb{R}^m}^* f d\mathcal{L}^m < \infty$  the function  $\psi_f$  is unique up to change on a set of measure zero. Notice also that if  $\{f_i\}$  is an increasing sequence of maps  $\mathbb{R}^m \rightarrow [0, \infty]$  and if  $f = \lim_{i \rightarrow \infty} f_i$ , then  $\lim_{i \rightarrow \infty} \int_{\mathbb{R}^m}^* f_i d\mathcal{L}^m = \int_{\mathbb{R}^m}^* f d\mathcal{L}^m$ .

**Proof of 1.10:**  $f$  is Lipschitz, hence uniformly continuous on  $A$ , and also  $\mathbb{R}^m$  is complete. So if  $x_k \in A \rightarrow x \in \bar{A}$  then the sequence  $\{f(x_k)\}_{k=1,2,\dots}$  is Cauchy in  $\mathbb{R}^m$ , hence has a limit which we denote  $\bar{f}(x)$ , and evidently  $\bar{f} : \bar{A} \rightarrow \mathbb{R}^m$  so defined is a Lipschitz extension of  $f$  to  $\bar{A}$  with  $\text{Lip } \bar{f} = \text{Lip } f$ .

For each  $i = 1, 2, \dots$  pick closed subsets  $C_{i1}, C_{i2}, \dots$  of  $X$  with  $\text{diam } C_{ij} < 1/i$ ,  $A \subset \bigcup_j C_{ij}$  and

$$(1) \quad \sum_j \omega_{m+k} (\text{diam } C_{ij}/2)^{m+k} \leq \mathcal{H}_{1/i}^{m+k}(A) + 1/i.$$

Next, observe that

$$\begin{aligned} (2) \quad \mathcal{H}_{1/i}^k(f^{-1}y) &\leq \sum_{j: f^{-1}y \cap C_{ij} \neq \emptyset} \omega_k (\text{diam } C_{ij}/2)^k \\ &= \sum_{j: y \in f(A \cap C_{ij})} \omega_k (\text{diam } C_{ij}/2)^k \\ &\leq \sum_{j: y \in \bar{f}(\bar{A} \cap C_{ij})} \omega_k (\text{diam } C_{ij}/2)^k \\ &= \sum_j \omega_k (\text{diam } C_{ij}/2)^k \chi_{\bar{f}(\bar{A} \cap C_{ij})}(y). \end{aligned}$$

Notice that the right side here is a Borel measurable function of  $y$  (because  $\bar{f}$  is continuous and  $\bar{A} \cap C_{ij}$  can be written as a countable union of compact sets for each  $j$  by  $\sigma$ -compactness of  $X$ ), but the left side need not be measurable. Nevertheless (see the

discussion in Remark 1.11 above) (2) implies

$$\begin{aligned}
 (3) \quad & \int_{\mathbb{R}^m}^* \mathcal{H}_{1/i}^k(f^{-1}y) d\mathcal{L}^m(y) \\
 & \leq \sum_j \omega_k (\text{diam } C_{ij}/2)^k \mathcal{L}^m(\bar{f}(\bar{A} \cap C_{ij})) \leq \sum_j \omega_k \omega_m (\text{Lip } f)^m (\text{diam } C_{ij}/2)^{m+k} \\
 & \leq \left( \frac{\omega_m \omega_k}{\omega_{m+k}} \right) (\text{Lip } f)^m (\mathcal{H}_{1/i}^{m+k}(A) + 1/i),
 \end{aligned}$$

where we used  $\mathcal{L}^m(\bar{f}(\bar{A} \cap C_{ij})) \leq \omega_m \left( \frac{\text{diam } \bar{f}(\bar{A} \cap C_{ij})}{2} \right)^m$  (by the isodiametric inequality 2.7)  $\leq \omega_m (\text{Lip } f)^m \left( \frac{\text{diam } C_{ij}}{2} \right)^m$ . Letting  $i \rightarrow \infty$  (and noting the discussion in Remark 1.11), we conclude

$$(4) \quad \int_{\mathbb{R}^m}^* \mathcal{H}^k(f^{-1}y) d\mathcal{L}^m(y) \leq \left( \frac{\omega_m \omega_k}{\omega_{m+k}} \right) (\text{Lip } f)^m \mathcal{H}^{m+k}(A).$$

It remains to check that  $\mathcal{H}^k(f^{-1}y)$  is an  $\mathcal{H}^m$ -measurable function of  $y \in \mathbb{R}^m$  (which will enable us to replace the upper integral on the left of (4) with the standard integral). This is left as an exercise (Problem 2.8 in Ch.2 problems).  $\square$

We conclude this section with a discussion of Lipschitz domains in  $\mathbb{R}^n$ .

**1.12 Definition:** A bounded open set  $\Omega \subset \mathbb{R}^n$  is said to be a Lipschitz domain if there are constants  $0 < \sigma \leq \tau$  such that  $\forall y \in \partial\Omega$  there is a  $v \in \mathbb{S}^{n-1}$  and a Lipschitz function  $u : B_\sigma(0) \cap v^\perp \rightarrow (-\tau, \tau)$  such that

$$\begin{aligned}
 U_y \cap \Omega &= \{y + x + tv : x \in \check{B}_\sigma(0) \cap v^\perp, t < u(x)\} \\
 U_y \cap \partial\Omega &= \{y + x + tv : x \in \check{B}_\sigma(0) \cap v^\perp, t = u(x)\},
 \end{aligned}$$

where  $U_y$  is the open neighborhood of  $y$  given by

$$U_y = \{y + x + tv : x \in \check{B}_\sigma(0) \cap v^\perp, -\tau < t < \tau\}.$$

Thus, roughly speaking,  $\Omega$  is Lipschitz means that locally, near each of its points,  $\partial\Omega$  can be expressed as the graph of a Lipschitz function.

Of course the bounded open convex subsets of  $\mathbb{R}^n$  are automatically Lipschitz domains; more precisely, we have the following lemma:

**1.13 Lemma.** *Suppose that  $\Omega \subset \mathbb{R}^n$  is an open, bounded and convex. Then  $\Omega$  is Lipschitz. In fact if  $0 \in \Omega$ , and  $R > 0, \delta \in (0, 1)$  are such that  $B_{\delta R}(0) \subset \Omega \subset B_R(0)$ , then for each  $y \in \partial\Omega$  there is a Lipschitz function*

$$u : \check{B}_{\delta R/2}(0) \cap y^\perp \rightarrow (0, \infty) \text{ with } u(0) \in (\delta R, R], \text{ Lip } u \leq 2/\delta,$$

and

$$\begin{aligned}
 U_y^+ \cap \Omega &= \{x + ty : x \in \check{B}_{\delta R/2}(0) \cap y^\perp, 0 < t < u(x)\} \\
 U_y^+ \cap \partial\Omega &= \{x + ty : x \in \check{B}_{\delta R/2}(0) \cap y^\perp, t = u(x)\},
 \end{aligned}$$

where  $U_y^+$  is the open neighborhood of  $y$  defined by

$$U_y^+ = \{x + ty : x \in \check{B}_{\delta R/2}(0) \cap y^\perp, t > 0\}.$$

**Proof:** By scaling we can assume without loss of generality that  $R = 1$ , so  $B_\delta(0) \subset \Omega \subset B_1(0)$ . Let  $y \in \partial\Omega$ . By applying a suitable rotation we can also assume that  $y = \rho e_n$  with  $\rho \in (\delta, 1]$ . If  $p : \mathbb{R}^n = \mathbb{R}^{n-1} \times \mathbb{R} \rightarrow \mathbb{R}^{n-1}$  is the projection  $(x, t) \mapsto x$  and if  $U = \check{B}_{\delta/2}^{n-1}(0) \times (0, \infty)$  then evidently

$$(1) \quad p(U \cap \partial\Omega) = \check{B}_{\delta/2}^{n-1}(0).$$

Let  $(x_1, t_1), (x_2, t_2) \in U \cap \partial\Omega$  be arbitrary with  $t_2 \geq t_1$ , and let  $\pi$  be a supporting hyperplane for  $\bar{\Omega}$  at  $(x_1, t_1)$ , so that there is an open half space  $H$  with

$$\pi = \partial H, B_\delta(0) \subset \Omega \subset H, (x_1, t_1) \in \pi.$$

Then  $\pi \cap B_\delta(0) = \emptyset$ , so  $\pi$  is not a vertical hyperplane and we can write

$$\pi = \{(x, t) : t = t_1 + a \cdot (x - x_1)\} \text{ and } H = \{(x, t) : t < t_1 + a \cdot (x - x_1)\},$$

where  $a \in \mathbb{R}^{n-1}$ . We must also then have  $|a| \leq 2/\delta$ , since otherwise there is a point  $x \in \check{B}_\delta^{n-1}(0)$  with  $a \cdot (x - x_1) = -t_1$  which would imply  $(x, 0) \in \pi \cap \check{B}_\delta(0)$ , contradicting  $\pi \cap B_\delta(0) = \emptyset$ .

Finally  $(x_2, t_2) \in \bar{H}$ , so  $0 \leq t_2 - t_1 \leq a \cdot (x_2 - x_1)$  and hence

$$(2) \quad 0 \leq t_2 - t_1 \leq 2\delta^{-1}|x_2 - x_1|.$$

The existence of  $u : \check{B}_{\delta/2}^{n-1}(0) \rightarrow (0, \infty)$  with  $\text{Lip } u \leq 2/\delta$  and  $\check{B}_{\delta/2}^{n-1}(0) \times (0, \infty) \cap \partial\Omega = \text{graph } u$  is now a direct consequence of (1), (2).  $\square$

## 2 BV Functions

In this section we gather together the basic facts about locally BV (i.e. bounded variation) functions which will be needed later.

First recall that if  $U$  is open in  $\mathbb{R}^n$  and if  $u \in L_{\text{loc}}^1(U)$ , then  $u$  is said to be in  $BV_{\text{loc}}(U)$  if for each  $W \subset\subset U$  there is a constant  $c(W) < \infty$  such that

$$2.1 \quad \int_W u \operatorname{div} g \, d\mathcal{L}^n \leq c(W) \sup |g|$$

for all vector functions  $g = (g^1, \dots, g^n)$ ,  $g^j \in C_c^\infty(W)$ . Notice that this means that the functional  $\int_U u \operatorname{div} g$  extends uniquely to give a (real-valued) linear functional on

$\mathcal{K}(U, \mathbb{R}^n) \equiv \{\text{continuous } g = (g^1, \dots, g^n) : U \rightarrow \mathbb{R}^n \text{ with } \text{spt } |g| \text{ compact}\}$  which is bounded on

$$\mathcal{K}_W(U, \mathbb{R}^n) \equiv \{g \in \mathcal{K}(U, \mathbb{R}^n) : \text{spt } |g| \subset W\}$$

for every  $W \subset\subset U$ . Then, by the Riesz Representation Theorem 5.14 of Ch. 1, there is a Radon measure  $\mu$  on  $U$  and a Borel measurable function  $v = (v^1, \dots, v^n)$ ,  $|v| = 1$  a.e., such that

$$2.2 \quad \int_U u \operatorname{div} g \, d\mathcal{L}^n = \int_U g \cdot v \, d\mu.$$

Thus, in the language of distribution theory, the generalized derivatives  $D_j u$  of  $u$  are represented by the *signed measures*  $v_j \, d\mu$ ,  $j = 1, \dots, n$ . For this reason we often denote the total variation measure  $\mu$  of Ch. 1) by  $|Du|$ . In fact if  $u \in W_{\text{loc}}^{1,1}(U)$  we evidently do have  $d\mu = |Du| \, d\mathcal{L}^n$  and

$$2.3 \quad v_j = \begin{cases} \frac{D_j u}{|Du|} & \text{if } |Du| \neq 0 \\ 0 & \text{if } |Du| = 0. \end{cases}$$

Thus for  $u \in BV_{\text{loc}}(U)$ ,  $|Du|$  will henceforth denote the Radon measure on  $U$  which is uniquely characterized by

$$2.4 \quad |Du|(W) = \sup_{\substack{|g| \leq 1, \text{ spt } |g| \subset\subset W, g \text{ Lipschitz}}} \int_U u \operatorname{div} g \, d\mathcal{L}^n, \quad W \text{ open } \subset U.$$

The left side here is more usually denoted  $\int_W |Du|$ . Indeed if  $f$  is any non-negative Borel measurable function on  $U$ , then  $\int f \, d|Du|$  is more usually denoted simply by  $\int f |Du|$  ( $\equiv \int f |Du| \, d\mathcal{L}^n$  in case  $u \in W_{\text{loc}}^{1,1}(U)$ ). We shall henceforth adopt this notation.

There are a number of important results about  $BV$  functions which can be obtained by mollification. We let  $\varphi_\sigma(x) = \sigma^{-n} \varphi(x/\sigma)$ , where  $\varphi$  is a symmetric mollifier (so that  $\varphi \in C_c^\infty(\mathbb{R}^n)$ ,  $\varphi \geq 0$ ,  $\text{spt } \varphi \subset B_1(0)$ ,  $\int_{\mathbb{R}^n} \varphi = 1$ , and  $\varphi(x) = \varphi(-x)$ ), and for  $u \in L_{\text{loc}}^1(U)$  let  $u^{(\sigma)} = \varphi_\sigma * \tilde{u}$  be the mollified functions, where we set  $\tilde{u} = u$  on  $U_\sigma$ ,  $\tilde{u} = 0$  outside  $U_\sigma$ ,  $U_\sigma = \{x \in U : \text{dist}(x, \partial U) > \sigma\}$ . A key result concerning mollification is then as follows:

**2.5 Lemma.** *If  $u \in BV_{\text{loc}}(U)$ , then  $u^{(\sigma)} \rightarrow u$  in  $L_{\text{loc}}^1(U)$  and  $|Du^{(\sigma)}| \rightarrow |Du|$  in the sense of Radon measures in  $U$  (see 5.15 of Ch. 1) as  $\sigma \downarrow 0$ .*

The convergence of  $u^{(\sigma)}$  to  $u$  in  $L_{\text{loc}}^1(U)$  is standard. Thus it remains to prove

$$(1) \quad \lim_{\sigma \downarrow 0} \int f |Du^{(\sigma)}| = \int f |Du|$$

for each  $f \in C_c^0(U)$ ,  $f \geq 0$ . In fact by definition of  $|Du|$  it is rather easy to prove that

$$(2) \quad \int f |Du| \leq \liminf_{\sigma \downarrow 0} \int f |Du^{(\sigma)}|,$$

so we only have to check

$$(3) \quad \limsup_{\sigma \downarrow 0} \int f |Du^{(\sigma)}| \leq \int f |Du|$$

for each  $f \in C_c^0(U)$ ,  $f \geq 0$ .

This is achieved as follows: First note that

$$(4) \quad \int f |Du^{(\sigma)}| = \sup_{|g| \leq f, g \text{ smooth}} \int g \cdot \nabla u^{(\sigma)} d\mathcal{L}^n.$$

On the other hand for fixed  $g$  with  $g$  smooth and  $|g| \leq f$ , and for  $\sigma < \text{dist}\{\text{spt } f, \partial U\}$ , we have

$$\begin{aligned} \int g \cdot \nabla u^{(\sigma)} d\mathcal{L}^n &= - \int u^{(\sigma)} \text{div } g d\mathcal{L}^n \\ &= - \int \varphi_\sigma * u \text{div } g d\mathcal{L}^n \\ &= - \int u (\varphi_\sigma * \text{div } g) d\mathcal{L}^n \\ &= - \int u \text{div}(\varphi_\sigma * g) d\mathcal{L}^n. \end{aligned}$$

On the other hand by definition of  $|Du|$ , the right side here is

$$\leq \int_{W_\sigma} (f + \varepsilon(\sigma)) |Du|$$

where  $W_\sigma = \{x \in U : \text{dist}(x, \text{spt } f) < \sigma\}$ , because

$$\begin{aligned} |\varphi_\sigma * g| &\equiv |(\varphi_\sigma * g^1, \dots, \varphi_\sigma * g^n)| \\ &\leq \varphi_\sigma * |g| \leq \varphi_\sigma * f \end{aligned}$$

and because  $\varphi_\sigma * f \rightarrow f$  uniformly in  $W_{\sigma_0}$  as  $\sigma \downarrow 0$ , where  $\sigma_0 < \text{dist}(\text{spt } f, \partial U)$ . Thus (3) follows from (4).  $\square$

**2.6 Theorem (Compactness Theorem for BV Functions.)** *If  $\{u_k\}$  is a sequence of  $BV_{\text{loc}}(U)$  functions satisfying*

$$\sup_{k \geq 1} (\|u_k\|_{L^1(W)} + \int_W |Du_k|) < \infty$$

for each  $W \subset\subset U$ , then there is a subsequence  $\{u_{k'}\} \subset \{u_k\}$  and a  $BV_{\text{loc}}(U)$  function  $u$  such that  $u_{k'} \rightarrow u$  in  $L^1_{\text{loc}}(U)$  and

$$\int_W |Du| \leq \liminf \int_W |Du_{k'}| \quad \forall W \subset\subset U.$$

**Proof:** By virtue of the previous lemma, in order to prove  $u_{k'} \rightarrow u$  in  $L^1_{\text{loc}}(U)$  for some subsequence  $\{u_{k'}\}$ , it is enough to prove that the sets

$$\{u \in C^\infty(U) : \int_W (|u| + |Du|) d\mathcal{L}^n \leq c(W)\}, \quad W \subset\subset U,$$

(for given constants  $c(W) < \infty$ ) are precompact in  $L^1_{\text{loc}}(U)$ . For the simple proof of this (involving mollification and Arzela's theorem) see for example [GT01, Theorem 7.22].

Finally the fact that  $\int_W |Du| \leq \liminf \int_W |Du_{k'}|$  is a direct consequence of the definition of  $|Du|, |Du_{k'}|$ .  $\square$

Next we have the Poincaré inequality for BV functions.

**2.7 Lemma.** *Suppose  $U$  is bounded, open and convex, let  $\delta \in (0, 1)$  be such that there is  $R > 0$  and  $\xi \in U$  with  $B_{\delta R}(\xi) \subset U \subset B_R(\xi)$ , and let  $u \in BV(U)$ . Then for any  $\theta \in (0, 1)$  and any  $\beta \in \mathbb{R}$  with*

$$(\ddagger) \quad \min\{\mathcal{L}^n\{x \in U : u(x) \geq \beta\}, \mathcal{L}^n\{x \in U : u(x) \leq \beta\}\} \geq \theta \mathcal{L}^n(U).$$

*we have*

$$\int_U |u - \beta| d\mathcal{L}^n \leq CR \int_U |Du|,$$

*where  $C = C(\theta, \delta, n)$ .*

**Proof:** By rescaling  $x \mapsto R^{-1}(x - \xi)$  we can without loss of generality assume  $R = 1$  and  $\xi = 0$ .

Let  $\beta, \theta$  be as in 2.7  $(\ddagger)$  and choose convex  $W \subset U$  such that

$$(\dagger) \quad \int_W |u - \beta| d\mathcal{L}^n \geq \frac{1}{2} \int_U |u - \beta| d\mathcal{L}^n$$

and such that 2.7  $(\ddagger)$  holds with  $W$  in place of  $U$  and  $\theta/2$  in place of  $\theta$ . (For example we may take  $W = \{x \in U : \text{dist}(x, \partial U) > \eta\}$  with  $\eta$  small.)

Letting  $u_\sigma$  denote the mollified functions corresponding to  $u$ , note that for sufficiently small  $\sigma$  we must have 2.7  $(\ddagger)$  with  $u_\sigma$  in place of  $u$ ,  $\theta/4$  in place of  $\theta$ , and  $W$  in place of  $U$ . Hence by the usual Poincaré inequality for smooth functions (see e.g. [GT01]) we have, with suitable  $\beta^{(\sigma)} \rightarrow \beta$  in place of  $\beta$ ,

$$\int_W |u_\sigma - \beta^{(\sigma)}| d\mathcal{L}^n \leq c \int_W |Du_\sigma| d\mathcal{L}^n,$$

$c = c(n, \theta, \delta)$ , for all sufficiently small  $\sigma$ . The required inequality now follows by letting  $\sigma \downarrow 0$  and using  $(\dagger)$  above together with 2.5.  $\square$

**2.8 Lemma.** *Suppose  $U, \delta, \xi, R$  are as in 2.7,  $u \in BV(\mathbb{R}^n)$  with  $\text{spt } u \subset \bar{U}$ . Then*

$$\int_{\mathbb{R}^n} |Du| (= \int_{\bar{U}} |Du|) \leq C \left( \int_U |Du| + R^{-1} \int_U |u| d\mathcal{L}^n \right),$$

where  $C = C(\delta, n)$ .

**2.9 Remark:** Note that by combining this with the Poincaré inequality in 2.7, we conclude

$$R^{-1} \int_{\mathbb{R}^n} |u - \beta \chi_U| + \int_{\mathbb{R}^n} |D(u - \beta \chi_U)| \leq C \int_U |Du|,$$

$C = C(\theta, \delta)$ , whenever  $\beta$  is as in 2.7  $(\ddagger)$ .

**Proof of 2.8:** As in the proof of 2.7, we can assume without loss of generality that  $R = 1$  and  $\xi = 0$ .

Let  $d(x) = \text{dist}(x, \partial U)$ ,  $x \in \mathbb{R}^n$ . Observe  $x, z \in \bar{U}$  with  $d(x) \leq d(z) \Rightarrow d(x + t(z - x)) \geq d(x) \forall t \in [0, 1]$  (otherwise  $\min_{t \in [0, 1]} d(x + t(z - x)) < \min\{d(x), d(z)\}$ , which evidently contradicts the convexity of  $\bar{U}$ ). Thus  $d(x + t(z - x))|_{[0, 1]}$  attains its minimum value  $d(x)$  at  $t = 0$ , and hence

$$(1) \quad (z - x) \cdot Dd(x) = \frac{d}{dt} d(x + t(z - x)) \Big|_{t=0} \geq 0$$

for all pairs  $x, z \in \bar{U}$  such that  $d$  is differentiable at  $x$  and  $d(x) \leq d(z)$ . In particular since  $B_\delta(0) \subset U$  (recall we assume  $B_{\delta R}(\xi) \subset U \subset B_R(\xi)$  with  $R = 1$  and  $\xi = 0$ ), for any  $\sigma > 0$  such that  $B_{\delta+\sigma}(0) \subset U$  and any  $x \in U$  with  $d(x) < \sigma$  and  $d$  differentiable at  $x$  we can take  $z = -\delta Dd(x)$  in (1) (because then  $z \in B_\delta(0)$  and so  $d(z) > \sigma > d(x)$ ). Hence  $(-x - \delta Dd(x)) \cdot Dd(x) \geq 0$ , and so

$$(2) \quad -x \cdot Dd(x) \geq \delta, \text{ a.e. } x \in U \text{ with } d(x) < \sigma.$$

Then we let  $\gamma_\sigma : \mathbb{R} \rightarrow [0, 1]$  be an increasing  $C^1$  function with  $\gamma_\sigma(t) \equiv 0$  for  $t \leq \sigma/2$  and  $\gamma(t) \equiv 1$  for  $t \geq \sigma$ , and set

$$(3) \quad \varphi_\sigma = \gamma_\sigma \circ d$$

Then by (2) and (3) we have, for  $\sigma < \text{dist}(B_\delta(0), \partial U)$ ,

$$(4) \quad \delta |D\varphi_\sigma(x)| \leq -x \cdot D\varphi_\sigma(x), \quad x \in \bar{U}.$$

Now by definition of  $|Dw|$  for  $BV_{\text{loc}}(\mathbb{R}^n)$  functions  $w$ , we have

$$(5) \quad \int_{\mathbb{R}^n} |D(\varphi_\sigma u)| \leq \int_{\mathbb{R}^n} |D\varphi_\sigma| |u| d\mathcal{L}^n + \int_{\mathbb{R}^n} \varphi_\sigma |Du|$$

and by (4)

$$\begin{aligned}
 (6) \quad \delta \int_{\mathbb{R}^n} |D\varphi_\sigma| |u| d\mathcal{L}^n &\leq - \int x \cdot D\varphi_\sigma |u| d\mathcal{L}^n \\
 &= - \int |u| \operatorname{div}(x\varphi_\sigma) d\mathcal{L}^n + n \int |u| \varphi_\sigma d\mathcal{L}^n \\
 &\leq \int_U |D|u|| + n \int_{\mathbb{R}^n} |u| d\mathcal{L}^n \text{ (by definition of } |D|u||) \\
 &\leq \int_U |Du| + n \int_{\mathbb{R}^n} |u| d\mathcal{L}^n
 \end{aligned}$$

(because  $|D|u|| \leq |Du|$  by virtue of 2.5 and the fact that  $|D|u|| \leq \liminf_{\sigma \downarrow 0} |D|u_\sigma||$ ).

Finally, to complete the proof of 2.8, we note that (using the definition of  $|Dw|$  for the  $BV_{\text{loc}}(\mathbb{R}^n)$  functions  $w = u, \varphi_\sigma u$ , together with the fact that  $\varphi_\sigma u \rightarrow u$  in  $L^1(\mathbb{R}^n)$  as  $\sigma \downarrow 0$ )

$$\int_{\mathbb{R}^n} |Du| \leq \liminf_{\sigma \downarrow 0} \int_{\mathbb{R}^n} |D(\varphi_\sigma u)|.$$

Then 2.8 follows from (5), (6).  $\square$

### 3 The Area Formula

The area formula, which we establish in this section, generalizes the classical formula

$$3.1 \quad \mathcal{L}^n(\tau(A)) = |\det \tau| \mathcal{L}^n(A)$$

established in Corollary 1.20 of Ch.1, valid for any linear transformation  $\tau : \mathbb{R}^n \rightarrow \mathbb{R}^n$  and any subset  $A \subset \mathbb{R}^n$ .

In the statement below we assume  $f : U \rightarrow \mathbb{R}^m$  ( $U \subset \mathbb{R}^n$  open) is locally Lipschitz (i.e. Lipschitz on each ball  $B_\rho(y) \subset U$ ), with  $n \leq m$ , and we define

$$3.2 \quad J_f(x) = \sqrt{\det((d_x f)^*(d_x f))} = \sqrt{\det(D_i f(x) \cdot D_j f(x))}$$

at all points where this exists (which is for  $\mathcal{L}^n$ -a.e.  $x \in U$  by virtue of Rademacher's Theorem 1.4).

**3.3 Theorem (Area Formula.)** *Suppose  $U$  is open in  $\mathbb{R}^n$  and  $f : U \rightarrow \mathbb{R}^m$  is locally Lipschitz, with  $n \leq m$ , and  $J_f$  is as in 3.2. Then*

$$\int_A J_f d\mathcal{L}^n = \int_{\mathbb{R}^m} \mathcal{H}^0(A \cap f^{-1}y) d\mathcal{H}^n(y)$$

for each Lebesgue measurable  $A \subset U$ . In particular

$$\int_A J_F d\mathcal{L}^n = \mathcal{H}^n(f(A)) \text{ provided } f|_A \text{ is 1:1.}$$

**3.4 Remarks:** (1) Observe that  $\mathcal{H}^0(A \cap f^{-1}y)$  is by definition just the “multiplicity function of  $f|A$ ”—i.e. the number of points (possibly  $\infty$ ) in  $\{x \in A : f(x) = y\}$ . Part of the conclusion of the above theorem is that  $\mathcal{H}^0(A \cap f^{-1}y)$  is an  $\mathcal{H}^n$ -measurable function of  $y$ .

(2) If  $h$  is a non-negative  $\mathcal{L}^n$ -measurable function on  $U$  then we have the more general formula

$$\begin{aligned} \int_U h J_f d\mathcal{L}^n &= \int_{\mathbb{R}^m} \int_{f^{-1}y} h(x) d\mathcal{H}^0(x) d\mathcal{H}^n(y) \\ &= \int_{\mathbb{R}^m} (\sum_{x \in f^{-1}y} h(x)) d\mathcal{H}^n(y). \end{aligned}$$

This clearly follows from the above theorem by writing  $h$  as the pointwise limit (everywhere) of an increasing sequence of non-negative (real-valued) simple functions.

**Proof of Theorem 3.3:** Since both sides of the identity are additive with respect to pairwise disjoint unions, it suffices to give the proof for the case when  $f$  is Lipschitz (rather than merely locally Lipschitz) and  $\mathcal{L}^n(A) < \infty$ .

We first consider Case (i):

$$A \subset \{x \in U : f \text{ is differentiable at } x \text{ and } J_f(x) > 0\}.$$

Then for each  $x \in A$  and each  $\varepsilon \in (0, 1)$ ,  $d_x f$  exists,  $\|d_x f\| \leq \text{Lip } f$  and  $\min_{v \in S^{n-1}} |d_x f(v)| > 0$ , and there is  $\delta_x = \delta_x(\varepsilon) > 0$  such that  $|f(z) - f(x) - d_x f(z - x)| \leq \frac{1}{2}\varepsilon |d_x f(z - x)|$  for all  $z \in B_{\delta_x}(x)$ , hence in particular

$$(1) \quad (1 - \tfrac{1}{2}\varepsilon) |d_x f(z - x)| \leq |f(x) - f(z)| \leq (1 + \tfrac{1}{2}\varepsilon) |d_x f(z - x)| \quad \forall z \in B_{\delta_x}(x).$$

Observe also the general fact that if  $\lambda : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a rank  $n$  linear map, then  $\lambda^* \circ \lambda$  is a positive definite symmetric linear map  $\mathbb{R}^n \rightarrow \mathbb{R}^n$  so by the Spectral Theorem there is an orthogonal transformation  $q : \mathbb{R}^n \rightarrow \mathbb{R}^n$  with

$$(2) \quad q^* \circ \lambda^* \circ \lambda \circ q = \Lambda,$$

where  $\Lambda$  is the diagonal transformation  $\Lambda e_j = \mu_j e_j$ ,  $j = 1, \dots, n$ , with  $0 < \mu_1 \leq \mu_2 \leq \dots \leq \mu_n$  the eigenvalues of  $\lambda^* \circ \lambda$ . Thus

$$(3) \quad \lambda^* \circ \lambda = \tau^2,$$

where  $\tau : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is the symmetric linear transformation given by

$$(4) \quad \tau = q \circ \sqrt{\Lambda} \circ q^*,$$

and hence by (3)

$$(5) \quad |\tau(v)| = |\lambda(v)| \quad \forall v \in \mathbb{R}^n \text{ and } |\det \tau| = \sqrt{\det(\lambda^* \circ \lambda)}.$$

Let  $\{\lambda_j : j = 1, 2, \dots\}$  be a dense set of rank  $n$  linear maps  $\mathbb{R}^n \rightarrow \mathbb{R}^m$ . Then, since  $\min_{v \in S^{n-1}} |d_x f(v)| > 0$  for  $x \in A$ , (1) implies that for each  $x \in A$  there is some  $j$  with

$$(1 - \varepsilon)|\lambda_j(z - x)| \leq |f(x) - f(z)| \leq (1 + \varepsilon)|\lambda_j(z - x)| \quad \forall z \in B_{\delta_x}(x),$$

where  $\delta_x$  is as in (1), and

$$(1 - \varepsilon)\sqrt{\det(\lambda_j^* \circ \lambda_j)} \leq \sqrt{\det((d_x f)^* \circ (d_x f))} = J_f(x) \leq (1 + \varepsilon)\sqrt{\det(\lambda_j^* \circ \lambda_j)},$$

and, by (3) and (5), for each  $j$  there is a symmetric rank  $n$  linear map  $\tau_j$  with  $\tau_j^2 = \lambda_j^* \circ \lambda_j$ , and hence

$$|\tau_j(v)| = |\lambda_j(v)| \quad \forall v \in \mathbb{R}^n \text{ and } |\det \tau_j| = \sqrt{\det(\lambda_j^* \circ \lambda_j)}.$$

Thus, with such  $\tau_j$ , for each  $x \in A$  there is  $j$  with

$$(6) \quad (1 - \varepsilon)|\tau_j(z - x)| \leq |f(x) - f(z)| \leq (1 + \varepsilon)|\tau_j(z - x)| \quad \forall z \in B_{\delta_x}(x) \text{ and}$$

$$(7) \quad (1 - \varepsilon)|\det \tau_j| \leq J_f(x) \leq (1 + \varepsilon)|\det \tau_j|.$$

Thus we can decompose  $A$  into a disjoint union  $\cup_j A_j$  of  $\mathcal{H}^n$ -measurable sets  $A_j$  such that (6), (7) hold for each  $x \in A_j$ , and for each  $j$  we let  $A_{ji} = \{x \in A_j : \delta_x \geq 1/i\}$ ,  $i = 1, 2, \dots$ . Observe that then, by (6),

$$(1 - \varepsilon)|\tau_j(x_1 - x_2)| \leq |f(x_1) - f(x_2)| \leq (1 + \varepsilon)|\tau_j(x_1 - x_2)|$$

for each  $x_1, x_2 \in A_{ji}$  such that  $|x_1 - x_2| < 1/i$ , so we can select pairwise disjoint  $\mathcal{H}^n$ -measurable sets  $A_{ji\ell} \subset A_{ji}$  with  $\text{diam } A_{ji\ell} < 1/i$  for each  $\ell$  and  $\cup_\ell A_{ji\ell} = A_{ji}$ , giving

$$\text{Lip}(f \circ \tau_j^{-1}|_{\tau_j(A_{ji\ell})}) \leq (1 + \varepsilon), \quad \text{Lip}((f \circ \tau_j^{-1}|_{\tau_j(A_{ji\ell})})^{-1}) \leq 1/(1 - \varepsilon), \quad \ell = 1, 2, \dots$$

Since  $f(A_{ji\ell}) = (f \circ \tau_j^{-1})(\tau_j(A_{ji\ell}))$ , we can then use 1.8 to yield

$$(8) \quad (1 - \varepsilon)^n |\det \tau_j| \mathcal{L}^n(A_{ji\ell}) = (1 - \varepsilon)^n \mathcal{H}^n(\tau_j(A_{ji\ell})) \\ \leq \mathcal{H}^n(f(A_{ji\ell})) \leq (1 + \varepsilon)^n \mathcal{H}^n(\tau_j(A_{ji\ell})) = (1 + \varepsilon)^n |\det \tau_j| \mathcal{L}^n(A_{ji\ell}).$$

Hence, by (7),

$$(1 - \varepsilon)^{n+1} \int_{A_{ji\ell}} J_f(x) d\mathcal{L}^n(x) \leq \mathcal{H}^n(f(A_{ji\ell})) \leq (1 + \varepsilon)^{n+1} \int_{A_{ji\ell}} J_f(x) d\mathcal{L}^n(x),$$

and, since

$$\sum_\ell \chi_{f(A_{ji\ell})}(y) = \mathcal{H}^0(A_{ji} \cap f^{-1}y), \quad y \in \mathbb{R}^m,$$

by summing on  $\ell$  we obtain

$$(1 - \varepsilon)^{n+1} \int_{A_{ji}} J_f d\mathcal{L}^n \leq \int_{\mathbb{R}^m} \mathcal{H}^0(A_{ji} \cap f^{-1}y) d\mathcal{H}^n(y) \leq (1 + \varepsilon)^{n+1} \int_{A_{ji}} J_f d\mathcal{L}^n.$$

Then we get the Area Formula in Case (i) by letting  $i \rightarrow \infty$  (using  $A_{ji} \subset A_{ji+1}$  and  $A_j = \cup_i A_{ji}$ ), then summing over  $j$ , and finally letting  $\varepsilon \downarrow 0$ .

Case (ii):  $A \subset \{x \in U : f \text{ is differentiable at } x \text{ and } J_f(x) = 0\}$ .

In this case we define  $F : \mathbb{R}^n \rightarrow \mathbb{R}^{m+n}$  by

$$F(x) = (f(x), \varepsilon x).$$

$F$  is then a 1:1 rank  $n$  map, so by Case (i)

$$(9) \quad \mathcal{H}^n(F(A)) = \int_{\mathbb{R}^{m+n}} J_F(x) d\mathcal{H}^n(x).$$

Now  $p \circ F = f$  where  $p$  is the projection  $(x, y) \in \mathbb{R}^n \times \mathbb{R}^m \mapsto x \in \mathbb{R}^n$ , and  $|p(z_1) - p(z_2)| \leq |z_1 - z_2|$ , so, using 1.8 with  $f = p$ ,

$$(10) \quad \mathcal{H}^n(f(A)) = \mathcal{H}^n(p(F(A))) \leq \mathcal{H}^n(F(A)).$$

Also by direct computation we have

$$J_F(x) = \sqrt{\det(D_i f(x) \cdot D_j f(x) + \varepsilon^2 \delta_{ij})} = J_f(x) + E(x), \quad 0 \leq E(x) \leq C\varepsilon^2,$$

where  $C = C(n, m, \text{Lip } f)$ . But for  $x \in A$  we have  $J_f(x) = 0$ , so using (9), (10) and letting  $\varepsilon \downarrow 0$  we obtain the Area Formula (with both sides 0) in Case (ii).

Case (iii):  $A \subset \{x \in U : f \text{ is not differentiable at } x\}$ .

By Rademacher's Theorem 1.4,  $\mathcal{H}^n(A) = 0$  and, since  $f$  is Lipschitz, in this case we can apply 1.8 to conclude  $\mathcal{H}^n(f(A)) = 0$ , hence the Area Formula trivially holds, with both sides 0, in Case (iii).  $\square$

**3.5 Examples:** (1) *Space curves*: Using the above area formula we first check that  $\mathcal{H}^1$ -measure agrees with the usual arc-length measure for  $C^1$  curves in  $\mathbb{R}^n$ . In fact if  $\gamma : [a, b] \rightarrow \mathbb{R}^n$  is a 1:1  $C^1$  map then the Jacobian  $J_\gamma$  is just  $\sqrt{|\dot{\gamma}|^2} = |\dot{\gamma}|$ , so that the Area Formula 3.3 gives

$$\mathcal{H}^1(\gamma(A)) = \int_A |\dot{\gamma}| d\mathcal{L}^1$$

as required.

(2) *Submanifolds of  $\mathbb{R}^{n+k}$* : If  $M$  is any  $n$ -dimensional embedded  $C^1$  submanifold of  $\mathbb{R}^{n+k}$  (see next section for a systematic discussion of such submanifolds), we want to check that  $\mathcal{H}^n \llcorner M$  (where  $\mathcal{H}^n$  is  $n$ -dimensional Hausdorff measure in  $\mathbb{R}^{n+k}$ ) agrees with the usual  $n$ -dimensional volume measure on  $M$ , i.e. that if  $\text{vol}$  denotes the volume measure (in the

usual sense of Riemannian geometry) on the submanifold  $M$ , and if  $\mathcal{H}^n$  is Hausdorff measure on the ambient space  $\mathbb{R}^{n+k}$ , then for Borel sets  $A \subset M$  (or more generally for  $\mathcal{H}^n$ -measurable sets  $A \subset M$ ) we have

$$(\ddagger) \quad \text{vol}(A) = \mathcal{H}^n(A).$$

It is enough to check this in a region where a local coordinate representation (see the discussion in §4 below) applies, because we can decompose the Borel set  $A$  into a countable pairwise disjoint union of Borel sets  $A_j$ , each of which is contained in the image of a local coordinate representation. Thus we suppose  $U$  is open in  $\mathbb{R}^{n+k}$  and that there is a local representation  $\psi$  for  $M$  such that

$$\psi : V \rightarrow \mathbb{R}^{n+k} \text{ is } C^1, \quad \psi(V) = M \cap W \text{ and } A \subset M \cap W,$$

where  $W$  is open in  $\mathbb{R}^{n+k}$  and let  $A_0 = \psi^{-1}(A) \subset V$  be the preimage (of course  $A_0$  is then also Borel). By the Area Formula 3.3

$$\mathcal{H}^n(A) = \int_{A_0} J_\psi \, d\mathcal{L}^n,$$

where  $J_\psi = \sqrt{\det(D_i \psi \cdot D_j \psi)}$ . Now notice on the other hand that  $g_{ij} = D_i \psi \cdot D_j \psi$  is the metric for  $M$  (relative to the local coordinates in  $V$ ) in the usual sense of Riemannian geometry, so this says  $\mathcal{H}^n(A) = \int_{A_0} \sqrt{g} \, d\mathcal{L}^n$ , where  $g = \det(g_{ij})$ , and the right side here is indeed the usual definition of  $\text{vol}(A)$  in the sense of Riemannian geometry, so  $(\ddagger)$  is established.

(3) *n-dimensional graphs in  $\mathbb{R}^{n+1}$* : If  $\Omega$  is a domain in  $\mathbb{R}^n$  and if  $M = \text{graph } u$ , where  $u \in C^1(\Omega)$ , then  $M$  is globally represented by the “graph map”  $\psi : x \mapsto (x, u(x))$ ; in this case

$$J_\psi(x) \equiv \sqrt{\det(D_i \psi \cdot D_j \psi)} \equiv \sqrt{\det(\delta_{ij} + D_i u D_j u)} = \sqrt{1 + |Du|^2},$$

so the Area Formula 3.3 in this case gives  $\mathcal{H}^n(M) = \int_{\Omega} \sqrt{1 + |Du|^2} \, dx$ .

## 4 Submanifolds of $\mathbb{R}^{n+k}$

Let  $M$  denote an  $n$ -dimensional embedded  $C^q$  submanifold of  $\mathbb{R}^{n+k}$ ,  $0 \leq k, r \geq 1$ . By this we mean  $M$  is a subset of  $\mathbb{R}^{n+k}$  such that for each  $x \in M$  there are open sets  $V \subset \mathbb{R}^n$ ,  $W \subset \mathbb{R}^{n+k}$ , and a 1:1  $C^q$  map  $\psi : V \rightarrow W$  with

$$4.1 \quad x \in \psi(V) = W \cap M,$$

where  $D\psi(\xi) = (D_i \psi^j(\xi))_{i=1, \dots, n, j=1, \dots, n+k}$  has rank  $n$  at each point  $\xi \in V$ , and the inverse map  $\varphi : \psi(\xi) \in W \cap M \mapsto \xi \in V$  is continuous. Such a map  $\psi$  will be referred to as “a local coordinate representation of  $M$  near  $x$ .”

**4.2 Remarks:** (1) The condition in the above definition that the inverse map  $\varphi$  is continuous allows examples like  $M = \{(x, \sin 1/x : x > 0)\} \subset \mathbb{R}^2$  as  $C^\infty$  submanifolds, but eliminates examples such as  $M = (\{0\} \times (-1, 1)) \cup \{(x, \sin 1/x : x > 0)\}$ . In fact the above definition (which requires that the local representation  $\psi$  is a homeomorphism of  $V$  onto the image  $M \cap W$ ) ensures that  $\psi$  is an open map onto its image; that is

$$V_0 \subset V, V_0 \text{ open} \Rightarrow \exists \text{ an open } W_0 \subset W \text{ with } \psi(V_0) = M \cap W_0.$$

(2) If the notation is as in the 4.1 and  $y_0 \in M \cap W$ ,  $x_0 \in V$  with  $y_0 = \psi(x_0)$ , then since  $\psi$  has rank  $n$  we can select  $1 \leq \ell_1 < \ell_2 < \dots < \ell_n \leq n+k$  with  $\det(D_i \psi^{\ell_j}(x_0)) \neq 0$ , and so, by the inverse function theorem, there is  $\delta_{x_0} > 0$  such that, with  $\pi : (y^1, \dots, y^{n+k}) \mapsto (y^{\ell_1}, \dots, y^{\ell_n})$ , the map  $\pi \circ \psi|_{\check{B}_{\delta_{x_0}}(x_0)}$  is a 1:1  $C^q$  map onto an open set  $U_0 \subset \mathbb{R}^n$  such that the inverse  $(\pi \circ \psi)^{-1} : U_0 \rightarrow \check{B}_{\delta_{x_0}}(x_0)$  is also  $C^q$ . Observe that then  $\varphi_{y_0} = (\pi \circ \psi)^{-1} \circ \pi : \pi^{-1}U_0 \rightarrow \check{B}_{\delta_0}(x_0)$  is also a  $C^q$  map and it agrees with  $\varphi$  (the inverse of  $\psi$ ) on  $\psi(\check{B}_{\delta_0}(x_0))$ , because  $\varphi_{y_0}(\psi(x)) = (\pi \circ \psi)^{-1} \circ (\pi \circ \psi)(x) = x$  for  $x \in \check{B}_{\delta_0}(x_0)$ . So  $\varphi_{y_0}$  is a  $C^q$  extension of  $\varphi|_{\psi(\check{B}_{\delta_{x_0}}(x_0))}$  to the open set  $\pi^{-1}(U_0)$ .

(3) Local graphical representations for  $M$ : Using the notation of Remark (2) above, in the special case when  $\ell_j = j$ ,  $j = 1, \dots, n$ , so  $\pi$  is just the projection of  $y = (y^1, \dots, y^{n+k})$  onto the first  $n$  coordinates  $(y^1, \dots, y^n)$ , we have  $\psi \circ (\pi \circ \psi)^{-1}(x) = (x, u(x))$ , with  $u = (u^1, \dots, u^k)$ ,  $u^j = \psi^{n+j} \circ (\pi \circ \psi)^{-1} : U_0 \rightarrow \mathbb{R}$ ,  $j = 1, \dots, k$ , and

$$\psi(\check{B}_{\delta_{x_0}}(x_0)) = \text{graph } u = \{(x, u(x)) : x \in U_0\}.$$

Also, by Remark (1) above,  $\psi(\check{B}_{\delta_{x_0}}(x_0)) = M \cap W_0$  for some open  $W_0$ , so then the “graph map”  $G : x \mapsto (x, u(x))$ ,  $x \in U_0$ , defines an alternate local representation for  $M$  near  $y_0$ . Without the assumption  $\ell_j = j$ ,  $j = 1, \dots, n$ , this of course remains true modulo composition with a permutation map (permuting the coordinates  $x^1, \dots, x^{n+k}$  in  $\mathbb{R}^{n+k}$  so that the coordinates  $x^{\ell_1}, \dots, x^{\ell_n}$  are moved to the first  $n$  slots), so for each  $y_0 \in M$  there is an open  $W_0$  with  $y_0 \in W_0$  and

$$(\ddagger) \quad M \cap W_0 = Q(\text{graph } u)$$

for some orthogonal transformation  $Q$  (where  $Q$  is in fact just a permutation of coordinates in  $\mathbb{R}^{n+k}$ ) and for some  $C^q$  vector function  $u = (u^1, \dots, u^k)$  defined on an open set  $U_0 \subset \mathbb{R}^n$ . Thus  $M$  is a  $C^q$  embedded submanifold of  $\mathbb{R}^{n+k}$  if and only if  $M$  is locally representable, near each of its points, as the graph of a  $C^q$  function  $u$ ; i.e. each  $y_0 \in M$  lies on some open  $W_0$  such that  $(\ddagger)$  holds, with  $u = (u^1, \dots, u^k)$  a  $C^q$  vector function on some open  $U_0 \subset \mathbb{R}^n$ .

If  $\psi : V \rightarrow W$  is as in 4.1 above, if  $A \subset M \cap W$  and  $h : A \rightarrow [0, \infty)$  are  $\mathcal{H}^n$ -measurable,

then we have the formula

$$4.3 \quad \int_A h d\mathcal{H}^n = \int_{\psi^{-1}(A)} h \circ \psi J_\psi d\mathcal{L}^n,$$

where  $J_\psi = \sqrt{\det(D_i \psi \cdot D_j \psi)}$ , which is checked using approximation of  $h$  by simple functions and using the special case of the area formula in Example 3.5(2) above.

Then if, for  $j = 1, 2$ ,  $M_j$  is a  $C^q$  submanifold of dimension  $n_j$  in  $\mathbb{R}^{n_j+k_j}$  with local representation  $\psi_j : V_j \rightarrow \mathbb{R}^{n_j+k_j}$  such that (as in 4.1)  $\psi_j(V_j) = M_j \cap W_j$ , then  $M_1 \times M_2$  is a  $C^q$  submanifold of dimension  $n = n_1 + n_2$  in  $\mathbb{R}^{n+k}$  ( $k = k_1 + k_2$ ) with local representation

$$\Psi : (x, y) \in V_1 \times V_2 \mapsto (\psi_1(x), \psi_2(y)) \in (M_1 \times M_2) \cap (W_1 \times W_2) \subset \mathbb{R}^{n+k},$$

and evidently

$$4.4 \quad J_\Psi(x, y) = J_{\psi_1}(x) J_{\psi_2}(y), \quad (x, y) \in V_1 \times V_2,$$

so by 4.3 and Fubini's theorem we see that

$$4.5 \quad H^n(A_1 \times A_2) = \mathcal{H}^{n_1}(A_1) \mathcal{H}^{n_2}(A_2)$$

for any  $\mathcal{H}^{n_j}$ -measurable subsets  $A_j \subset M_j$ ,  $j = 1, 2$ . In particular

$$4.6 \quad \mathcal{H}^n(M_1 \times M_2) = \mathcal{H}^{n_1}(M_1) \mathcal{H}^{n_2}(M_2).$$

**4.7 Definition:** The *tangent space*  $T_x M$  of  $M$  at  $x \in M$  is the subspace of  $\mathbb{R}^{n+k}$  consisting of those  $\tau \in \mathbb{R}^{n+k}$  such that  $\tau = \dot{\gamma}(0)$  for some  $C^1$  curve  $\gamma : (-\varepsilon, \varepsilon) \rightarrow \mathbb{R}^{n+k}$  (for some  $\varepsilon > 0$ ) with  $\gamma((-\varepsilon, \varepsilon)) \subset M$ ,  $\gamma(0) = x$ .

Note that

$$4.8 \quad T_x M \text{ is a linear subspace of } \mathbb{R}^{n+k} \text{ with basis } D_1 \psi(\xi), \dots, D_n \psi(\xi),$$

where  $\psi$  is any local representation as in 4.1 above with  $\psi(\xi) = x$ . Indeed if  $\gamma : (-\varepsilon, \varepsilon) \rightarrow \mathbb{R}^{n+k}$  is  $C^1$  with  $\gamma((-\varepsilon, \varepsilon)) \subset M$  and  $\gamma(0) = x$  then for all sufficiently small  $t$  we have  $\gamma(t) = \psi(\varphi(\gamma(t)))$ , where  $\varphi$  is the inverse of  $\psi$  on  $M \cap W$ . With  $\varphi_x$  the  $C^q$  extension of  $\varphi$  to a neighborhood of  $x$  defined in Remark 4.2(2) above, we thus have  $\gamma(t) = \psi(\varphi_x(\gamma(t)))$  for sufficiently small  $t$  and hence by the chain rule  $\gamma'(0) = \sum_{j=1}^n c^j D_j \psi(\xi)$ , where  $(c^1, \dots, c^n) = \frac{d}{dt} \varphi_x(\gamma(t))|_{t=0}$ . So  $T_x M \subset \text{span}\{D_1 \psi(\xi), \dots, D_n \psi(\xi)\}$  and of course the reverse inclusion is trivial because  $\sum_{j=1}^n c^j D_j \psi(\xi) = \frac{d}{dt} \psi(\xi + tc)|_{t=0}$  for any  $c = (c^1, \dots, c^n) \in \mathbb{R}^n$ .

**4.9 Definition:** A function  $f : M \rightarrow \mathbb{R}^{n+k}$  ( $k \geq 0$ ) is said to be  $C^q$  on  $M$  if  $f$  is the restriction to  $M$  of a  $C^q$  function  $\bar{f} : U \rightarrow \mathbb{R}^{n+k}$ , where  $U$  is an open set in  $\mathbb{R}^{n+k}$  containing  $M$ .

We next want to discuss some differentiability properties for locally Lipschitz maps  $f :$

$M \rightarrow \mathbb{R}^P$  with  $P \geq 1$ . Thus for each  $x \in M$  we assume there are  $\rho, L > 0$  with

$$4.10 \quad |f(y) - f(z)| \leq L|y - z| \quad y, z \in M \cap B_\rho(x).$$

First we discuss directional derivatives of such an  $f$ : For given  $\tau \in T_x M$  the directional derivative  $D_\tau f \in \mathbb{R}^P$  is defined by

$$4.11 \quad D_\tau f = \left. \frac{d}{dt} f(\gamma(t)) \right|_{t=0}$$

for any  $C^1$  curve  $\gamma : (-1, 1) \rightarrow M$  with  $\gamma(0) = x, \dot{\gamma}(0) = \tau$ , whenever this derivative exists. Of course it is easy to see that existence and the actual value is independent of the particular curve  $\gamma$  we use to represent  $\tau$  because if  $\tilde{\gamma}$  is another such curve then, by 4.10,

$$4.12 \quad \lim_{t \downarrow 0} t^{-1} |f(\gamma(t)) - f(\tilde{\gamma}(t))| \leq L \lim_{t \downarrow 0} t^{-1} |\gamma(t) - \tilde{\gamma}(t)| = 0$$

because  $\gamma(0) = \tilde{\gamma}(0) (= x)$  and  $\gamma'(0) = \tilde{\gamma}'(0) (= \tau)$ .

We claim that in fact there is a set  $E$  of  $\mathcal{H}^n$ -measure zero such that  $\forall x \in M \setminus E$

$$4.13 \quad D_\tau f(x) \text{ exists and the map } \tau \mapsto D_\tau f(x) \text{ is a linear map } T_x M \rightarrow \mathbb{R}^P,$$

so we can define the induced linear map  $d_x^M f : T_x M \rightarrow \mathbb{R}^P$  by

$$4.14 \quad d_x^M f(v) = D_v f(x), \quad v \in T_x M.$$

Indeed 4.13 is a consequence of the Rademacher theorem in  $\mathbb{R}^n$  proved in 1.4, as follows: Let  $x \in M$  and let  $\psi : V \rightarrow \mathbb{R}^{n+k}$  be  $C^1$  with  $x \in \psi(V) = M \cap W$  as in 4.1. Then according to 1.4 there is  $E_0 \subset V$  with  $\mathcal{H}^n(E_0) = 0$  such that  $f \circ \psi$  is differentiable at every point of  $V \setminus E_0$  and in particular for every  $\eta \in \mathbb{R}^n$  and  $x \in V \setminus E_0$  we have  $D_\eta(f \circ \psi)(x) = \left. \frac{d}{dt} f(\psi(x + t\eta)) \right|_{t=0}$  exists and is linear in  $\eta$ . But  $\gamma(t) = \psi(x + t\eta)$  is a curve as in 4.11 with  $\tau = \sum_{j=1}^n \eta_j D_j \psi(x)$ , so in fact this says that the directional derivatives

$$4.15 \quad D_{\sum_{j=1}^n \eta_j D_j \psi(\xi)} f(x) \text{ exist and } = D_\eta(f \circ \psi)(\xi)$$

and that furthermore this is linear in  $\eta$  for all  $\xi \in V \setminus E_0$  and  $x = \psi(\xi) \in W \cap M \setminus \psi(E_0)$ . Hence, since  $D_1 \psi(\xi), \dots, D_n \psi(\xi)$  is a basis for  $T_{\psi(\xi)} M$  by 4.8, this says that indeed 4.13 does hold at points of  $W \cap M \setminus \psi(E_0)$ , and of course  $\psi(E_0)$  is a set of  $\mathcal{H}^n$ -measure zero by 1.8 because  $\psi$  is locally Lipschitz in  $V$ .

Notice also that if in fact  $f$  is the restriction of a locally Lipschitz function  $\tilde{f}$  defined in an open set  $U \supset M$  then (by the same argument as in 4.12 with  $\tilde{\gamma}(t) = x + t\tau$ ) we have (for each given  $x \in M$  and  $\tau \in T_x M$ )

$$4.16 \quad D_\tau f(x) \text{ exists} \iff \left. \frac{d}{dt} \tilde{f}(x + t\tau) \right|_{t=0} \text{ exists},$$

and in that case the two quantities are equal; furthermore (by 4.13) for  $\mathcal{H}^n$ -a.e.  $x \in M$ ,  $D_\tau f(x)$  does exist for each  $\tau \in T_x M$  and it is a linear function of  $\tau \in T_x M$ .

Taking the particular choice  $\eta = e_i$ ,  $x = \psi(\xi) \in W \cap M \setminus \psi(E_0)$  in 4.15, and letting  $\tau_1, \dots, \tau_n$  be an orthonormal basis for  $T_x M$ , so that

$$D_i \psi(x) = \sum_{\ell=1}^n D_i \psi(x) \cdot \tau_\ell \tau_\ell,$$

we then have

$$D_i(f \circ \psi)(x) = \sum_{\ell=1}^n D_i \psi(\xi) \cdot \tau_\ell D_{\tau_\ell} f(x), \quad i = 1, \dots, n.$$

Thus

$$D_i(f \circ \psi)(x) \cdot D_j(f \circ \psi)(x) = \sum_{\ell, m=1}^n (D_i \psi(x) \cdot \tau_\ell) (D_j \psi(x) \cdot \tau_m) D_{\tau_\ell} f(x) \cdot D_{\tau_m} f(x).$$

Since  $\det AB = \det A \det B$  for square matrices  $A, B$ , we thus have

$$\begin{aligned} 4.17 \quad J_{f \circ \psi}(x) &= |\det(D_\ell \psi(x) \cdot \tau_m)| \sqrt{\det(D_{\tau_\ell} f(x) \cdot D_{\tau_m} f(x))} \\ &= J_\psi(x) J_f^M(x) \end{aligned}$$

with  $x = \psi(\xi)$ , where  $J_\psi = \sqrt{\det(D_i \psi \cdot D_j \psi)}$  (in accordance with 3.2) and

$$4.18 \quad J_f^M(x) = \sqrt{\det(D_{\tau_\ell} f(x) \cdot D_{\tau_m} f(x))},$$

with  $\tau_1, \dots, \tau_n$  any orthonormal basis for  $T_x M$ ;  $J_f^M$  is the “Jacobian of  $f : M \rightarrow \mathbb{R}^m$ ” (which is consistent with the terminology introduced in §3 in the special case when  $k = 0$  and  $M$  is an open subset of  $\mathbb{R}^n$ ).

Using 4.17 we now want to discuss the natural extension of the area formula 3.3 to the case when  $f : M \rightarrow \mathbb{R}^m$  is locally Lipschitz and  $m \geq n$ . We claim that in this case we have the general area formula

$$4.19 \quad \int_{f(A)} \mathcal{H}^0(A \cap f^{-1}y) d\mathcal{H}^n(y) = \int_A J_f^M d\mathcal{H}^n.$$

Since both sides here are additive with respect to disjoint unions, it is evidently enough to check this under the assumption that  $A = \psi(A_0)$  with  $\psi : V \rightarrow \mathbb{R}^{n+k}$  a local representation for  $M$  as in 4.1 and  $A_0 \subset V$  an  $\mathcal{H}^n$ -measurable set. Since  $f \circ \psi$  is locally Lipschitz on  $V$  we can use the Area Formula 3.3 with  $f \circ \psi$  in place of  $f$  and  $A_0$  in place

of  $A$  to give

$$\begin{aligned}
 \int_{f(A)} \mathcal{H}^0(A \cap f^{-1}y) d\mathcal{H}^n(y) &= \int_{f \circ \psi(A_0)} \mathcal{H}^0(A_0 \cap \psi^{-1}f^{-1}y) d\mathcal{H}^n(y) \\
 &= \int_{A_0} J_{f \circ \psi}(x) d\mathcal{H}^n(x) \\
 &= \int_{A_0} J_f^M(\psi(x)) J_\psi(x) d\mathcal{H}^n(x) \quad (\text{by 4.17}) \\
 &= \int_A J_f^M(y) d\mathcal{H}^n(y),
 \end{aligned}$$

where at the last step we used Remark 3.4(2) with  $\psi$  in place of  $f$  and with  $h = J_f^M \circ \psi$ . Thus 4.19 is proved.

More generally (by approximating  $h$  by an increasing sequence of non-negative simple functions and using 4.19),

$$4.20 \quad \int_{f(M)} (\sum_{x \in f^{-1}(y)} h(x)) d\mathcal{H}^n(y) = \int_M h(x) J_f^M(x) d\mathcal{H}^n(x)$$

for any non-negative  $\mathcal{H}^n$  measurable function  $h$  on  $M$ .

We can also (for any  $m \geq 1$ ) define the induced linear map  $df_x^M : T_x M \rightarrow \mathbb{R}^m$  just as we did in  $\mathbb{R}^n$  by

$$4.21 \quad df_x^M(\tau) = D_\tau f(x), \quad \tau \in T_x M.$$

In case  $f$  is real-valued (i.e.  $m = 1$ ) then we define the gradient  $\nabla^M f$  of  $f$  by

$$4.22 \quad \nabla^M f(x) = \sum_{j=1}^n (D_{\tau_j} f(x)) \tau_j, \quad x \in M,$$

where  $\tau_1, \dots, \tau_n$  is any orthonormal basis for  $T_x M$ . If we let  $\nabla_j^M f = e_j \cdot \nabla^M f$  ( $e_j = j$ -th standard basis vector in  $\mathbb{R}^{n+k}$ ,  $j = 1, \dots, n+k$ ) then

$$4.23 \quad \nabla^M f(x) = \sum_{j=1}^{n+k} \nabla_j^M f(x) e_j.$$

If  $f$  is the restriction to  $M$  of a  $C^1$  function  $\bar{f}$  on  $U$ , where  $U$  is an open subset of  $\mathbb{R}^{n+k}$  containing  $M$ , then

$$\nabla^M f(x) = p_{T_x M}(\nabla_{\mathbb{R}^{n+k}} \bar{f}(x)), \quad x \in M,$$

where  $\nabla_{\mathbb{R}^{n+k}} \bar{f}$  is the usual  $\mathbb{R}^{n+k}$  gradient  $(D_1 \bar{f}, \dots, D_{n+k} \bar{f})$  on  $U$ . Indeed, with  $\tau_1, \dots, \tau_n$  any orthonormal basis for  $T_x M$ ,  $p_{T_x M}(\nabla_{\mathbb{R}^{n+k}} \bar{f}(x)) = \sum_{i=1}^n \tau_i \cdot \nabla_{\mathbb{R}^{n+k}} \bar{f}(x) \tau_i = \sum_{i=1}^n D_{\tau_i} \bar{f}(x) \tau_i = \sum_{i=1}^n D_{\tau_i} f(x) \tau_i = \nabla^M f(x)$ .

Now given a vector function (“vector field”)  $X = (X^1, \dots, X^{n+k}) : M \rightarrow \mathbb{R}^{n+k}$  with each component  $X^j$  a Lipschitz function on  $M$ , we define

$$4.24 \quad \operatorname{div}_M X = \sum_{j=1}^{n+k} \nabla_j^M X^j$$

on  $M$ . (Notice that we do *not* require  $X_x \in T_x M$ .) Then, at  $x \in M$ , we have

$$4.25 \quad \begin{aligned} \operatorname{div}_M X &= \sum_{j=1}^{n+k} e_j \cdot (\nabla^M X^j) \\ &= \sum_{j=1}^{n+k} e_j \cdot \left( \sum_{i=1}^n (D_{\tau_i} X^j) \tau_i \right), \end{aligned}$$

so that (since  $X = \sum_{j=1}^{n+k} X^j e_j$ )

$$4.26 \quad \operatorname{div}_M X = \sum_{i=1}^n \tau_i \cdot D_{\tau_i} X,$$

where  $\tau_1, \dots, \tau_n$  is any orthonormal basis for  $T_x M$ .

Recall that the classical *Divergence Theorem* of Riemannian geometry states that if  $M$  is an  $n$ -dimensional  $C^2$  submanifold of  $\mathbb{R}^{n+k}$  and if  $X$  is a  $C^1$  *tangent* vector field on  $M$  with compact support in  $M$  (i.e.  $X(y) \in T_y M$  for each  $y \in M$  and  $\{y \in M : X(y) \neq 0\}$  is contained in a compact subset  $K$  of  $M$ ), then

$$4.27 \quad \int_M \operatorname{div}_M X \, d\mathcal{H}^n = 0.$$

This can be proved using local coordinates and a partition of unity, but to better motivate our later discussion of first variation of varifolds we give a more intrinsic proof as follows:

**Proof of 4.27:** Let  $K$  be a compact subset of  $M$  containing  $\{y \in M : X(y) \neq 0\}$  and let  $\varphi(t, x)$ ,  $(t, x) \in (-\varepsilon, \varepsilon) \times M$ , be the geometric flow on  $M$  generated by the tangent vector field  $X$ . Thus

$$\begin{cases} \partial \varphi(t, x) / \partial t = X(\varphi(t, x)) \\ \varphi(0, x) = x \end{cases}$$

By ODE theory (see e.g. [HL])  $\varphi$  and its velocity  $\partial \varphi(t, x) / \partial t$  exist and are  $C^1$  on  $(-\varepsilon, \varepsilon) \times M$  for small enough  $\varepsilon$ . Furthermore if  $\varphi_t(x) = \partial \varphi(t, x) / \partial t$ , then, for  $|t| < \delta$ ,  $\delta > 0$  sufficiently small,  $\varphi_t$  is a  $C^1$  diffeomorphism of  $M$  onto  $M$  and  $\varphi_t$  the identity on  $M \setminus K$ . So in particular  $\varphi_t(M \cap K) = M \cap K$  for all sufficiently small  $|t|$ , and hence trivially

$$(\ddagger) \quad \frac{d}{dt} \mathcal{H}^n(\varphi_t(M \cap K))|_{t=0} = 0.$$

But on the other hand the area formula gives

$$\mathcal{H}^n(\varphi_t(M \cap K)) = \int_{M \cap K} J_{\varphi_t} \, d\mathcal{H}^n,$$

where  $J_{\varphi_t}(y) = \sqrt{\det(D_{\tau_i}\varphi_t(y) \cdot D_{\tau_j}\varphi_t(y))}$ , and, since  $\varphi(t, y) = y + t(X(y) + E(t, y))$  with  $\sup_{y \in K} (|E(t, y)| + \sup_{\tau \in T_y M, |\tau|=1} |D_\tau E(t, y)|) \rightarrow 0$  as  $t \rightarrow 0$ , we have  $D_{\tau_i}\varphi(y) = \tau_i + t(D_{\tau_i}X(y) + E_i(t, y))$ , where  $\sup_{y \in K} |E_i(t, y)| \rightarrow 0$  as  $t \downarrow 0$ . Hence

$$J_{\varphi_t} = \sqrt{\det(\delta_{ij} + t(\tau_i \cdot D_{\tau_j}X(y) + \tau_j \cdot D_{\tau_i}X(y) + E_{ij}(t, y))},$$

where  $\sup_{y \in K} |E_{ij}(t, y)| \rightarrow 0$  as  $t \rightarrow 0$ . Using the formula  $\det(I + tA) = 1 + t \operatorname{trace} A + O(t^2)$  as  $t \rightarrow 0$ , we thus conclude

$$\frac{d}{dt} J_{\varphi_t}(y)|_{t=0} = \sum_{i=1}^n \tau_i \cdot D_{\tau_i}X(y) = \operatorname{div}_M X(y).$$

Hence by  $(\ddagger)$  we have 4.27 as claimed.  $\square$

**4.28 Remarks:** (1)  $M$  need *not* be orientable in the above discussion.

(2) If we drop the condition that  $\{y \in M : X(y) \neq 0\}$  is contained in a compact subset of  $M$ , and instead assume  $\overline{M}$  (the closure of  $M$  in  $\mathbb{R}^{n+k}$ ) is a compact manifold with  $C^1$  boundary  $\partial M = \overline{M} \setminus M$  and if we let  $X$  be any  $C^1$  vector field on  $\overline{M}$ , still with  $X(y) \in T_y M$  for each  $y \in M$ , then in place of 4.27 we get

$$(\ddagger) \quad \int_M \operatorname{div}_M X \, d\mathcal{H}^n = - \int_{\partial M} X \cdot \eta \, d\mathcal{H}^{n-1},$$

where  $\eta$  is the inward pointing unit co-normal of  $\partial M$ ; that is,  $|\eta| = 1$ ,  $\eta$  is normal to  $\partial M$ , tangent to  $M$ , and points into  $M$  at each point of  $\partial M$ .

(3) In general the closure  $\overline{M}$  of  $M$  will not be a nice manifold with boundary; indeed it can certainly happen that  $\mathcal{H}^n(\overline{M} \setminus M) > 0$ . (For example consider  $M = \{(x, y) \in \mathbb{R}^2 : x > 0, y = \sin(1/x)\}$ , in which case  $M$  is a  $C^\infty$  1-dimensional embedded submanifold of  $\mathbb{R}^2$  in the sense of the above definitions, but  $\overline{M} \setminus M$  is the interval  $\{0\} \times [-1, 1]$  on the  $y$ -coordinate axis.) Nevertheless, as we have shown above, 4.27 does hold provided  $\{y \in M : X(y) \neq 0\}$  is contained in a compact subset of  $M$  and  $X_y \in T_y M \, \forall y \in M$ .

In case  $M$  is at least  $C^2$  we define the second fundamental form of  $M$  at  $y$  to be the bilinear form

$$4.29 \quad B_y : T_y M \times T_y M \rightarrow (T_y M)^\perp$$

such that

$$4.30 \quad B_y(\tau, \eta) = -\sum_{\alpha=1}^k (\eta \cdot D_\tau v^\alpha) v^\alpha|_y, \quad \tau, \eta \in T_y M,$$

where  $v^1, \dots, v^k$  are (locally defined, near  $y$ ) vector fields with  $v^\alpha(z) \cdot v^\beta(z) = \delta_{\alpha\beta}$  and  $v^\alpha(z) \in (T_z M)^\perp$  for every  $z$  in some neighborhood of  $y$ . Of course such  $v^\alpha$  exist in a

neighborhood of any given  $y_0 \in M$ , because we can use a local representation  $\psi : V \rightarrow \mathbb{R}^n$  for  $M$  with  $y_0 = \psi(x_0)$  for some  $x_0 \in V$ , and then choose vectors  $\eta_1, \dots, \eta_k \in \mathbb{R}^{n+k}$  such that  $D_1\psi(x_0), \dots, D_n\psi(x_0), \eta_1, \dots, \eta_k$  are linearly independent. Then, for  $\delta > 0$  small enough,  $D_1\psi(x), \dots, D_n\psi(x), \eta_1, \dots, \eta_k$  are still linearly independent vectors in  $\mathbb{R}^{n+k}$  for all  $x \in \check{B}_\delta(x_0)$ , so the Gram-Schmidt orthogonalization process gives orthonormal  $C^1$  vector fields  $\tau_1(y), \dots, \tau_{n+k}(y)$  on  $M \cap \psi(\check{B}_\delta(x_0))$  (where  $y = \psi(x)$ ) such that  $\tau_1(y), \dots, \tau_n(y)$  is an orthonormal basis for  $T_y M$  and  $\nu^j(y) = \tau_{n+j}(y)$ ,  $j = 1, \dots, k$ , is an orthonormal basis for  $(T_y M)^\perp$  for each  $y \in \psi(\check{B}_\delta(x_0))$ .

The geometric significance of  $B$  is as follows: If  $\tau \in T_y M$  with  $|\tau| = 1$  and  $\gamma : (-\varepsilon, \varepsilon) \rightarrow \mathbb{R}^{n+k}$  (for some  $\varepsilon > 0$ ) is a  $C^2$  curve with  $\gamma(0) = y$ ,  $\gamma(-\varepsilon, \varepsilon) \subset M$ , and  $\dot{\gamma}(0) = \tau$ , then

$$4.31 \quad B_y(\tau, \tau) = (\ddot{\gamma}(0))^\perp,$$

which is just the normal component (relative to  $M$ ) of the curvature of  $\gamma$  at 0,  $\gamma$  being considered as an ordinary space-curve in  $\mathbb{R}^{n+k}$ . (Thus  $B_y(\tau, \tau)$  measures the “normal curvature” of  $M$  in the direction  $\tau$ .) To check this, simply note that  $\nu^\alpha(\gamma(t)) \cdot \dot{\gamma}(t) \equiv 0$ ,  $|t| < 1$ , because  $\dot{\gamma}(t) \in T_{\gamma(t)} M$  and  $\nu^\alpha(\gamma(t)) \in (T_{\gamma(t)} M)^\perp$ . Differentiating this relation with respect to  $t$ , we get (after setting  $t = 0$ )

$$\nu^\alpha(y) \cdot \ddot{\gamma}(0) = -(D_\tau \nu^\alpha) \cdot \tau$$

and hence (multiplying by  $\nu^\alpha(y)$  and summing over  $\alpha$ ) we have

$$\begin{aligned} (\ddot{\gamma}(0))^\perp &= -\sum_{\alpha=1}^k (\tau \cdot D_\tau \nu^\alpha) \nu^\alpha(y) \\ &= B_y(\tau, \tau) \end{aligned}$$

as required. (Note that the parameter  $t$  here need *not* be arc-length for  $\gamma$ ; it suffices that  $\dot{\gamma}(0) = \tau$ ,  $|\tau| = 1$ .) More generally, by a similar argument, if  $\tau, \eta \in T_y M$  and if  $\varphi : U \rightarrow \mathbb{R}^{n+k}$  is a  $C^2$  mapping of a neighborhood  $U$  of 0 in  $\mathbb{R}^2$  with  $\varphi(U) \subset M$ ,  $\varphi(0, 0) = y$ ,  $\frac{\partial \varphi}{\partial s}(0, 0) = \tau$ ,  $\frac{\partial \varphi}{\partial t}(0, 0) = \eta$ , then

$$4.32 \quad B_y(\tau, \eta) = -\left( \frac{\partial^2 \varphi}{\partial s \partial t}(0, 0) \right)^\perp.$$

Of course such maps  $\varphi$  do exist for any given  $\tau, \eta \in T_y M$ —for example we can let  $\psi : V \rightarrow W$  be a  $C^2$  local representation for  $M$ ,  $y = \psi(x)$ , and select  $z = (z^1, \dots, z^n)$   $w = (w^1, \dots, w^n) \in \mathbb{R}^n$  such that  $\tau = \sum_{j=1}^n z^j D_j \psi(x)$  and  $\eta = \sum_{j=1}^n w^j D_j \psi(x)$ , and then  $\varphi(s, t) = \psi(x + s z + t w)$  is a suitable choice for  $\varphi$ . Since  $\partial^2 \varphi / \partial s \partial t = \partial^2 \varphi / \partial t \partial s$ , 4.32 implies in particular that  $B_y(\tau, \eta) = B_y(\eta, \tau)$ ; that is  $B_y$  is a *symmetric* bilinear form with values in  $(T_y M)^\perp$ .

We define the mean curvature vector  $\underline{H}$  of  $M$  at  $y$  to be trace  $B_y$ ; thus

$$4.33 \quad \underline{H}(y) = \sum_{i=1}^n B_y(\tau_i, \tau_i) \in (T_y M)^\perp,$$

where  $\tau_1, \dots, \tau_n$  is an orthonormal basis for  $T_y M$ . Notice that then  $(v^1, \dots, v^k \in (T_y M)^\perp)$  are as in 4.30)

$$\underline{H}(y) = -\sum_{\alpha=1}^k \sum_{i=1}^n (\tau_i \cdot D_{\tau_i} v^\alpha) v^\alpha(y),$$

so that

$$4.34 \quad \underline{H}(y) = -\sum_{\alpha=1}^k (\operatorname{div}_M v^\alpha) v^\alpha$$

near  $y$ .

## 5 First Variation of a Submanifold

Let  $M$  be an  $n$ -dimensional  $C^2$  submanifold embedded in  $\mathbb{R}^{n+k}$  as in the previous section. We want to compute the initial rate of change (“first variation”) of  $\mathcal{H}^n(M)$  when  $M$  is undergoes a compactly supported perturbation via a 1-parameter family of maps  $\varphi_t : M \rightarrow \mathbb{R}^{n+k}$  with  $\varphi_0$  equal to the identity on  $M$ . So let  $K \subset M$  be compact and let  $\varphi : (-\varepsilon, \varepsilon) \times M \rightarrow \mathbb{R}^{n+k}$  (where  $\varepsilon > 0$ ) be a map such that

$$5.1 \quad \begin{cases} \varphi(0, x) = x, & \forall x \in M \\ \varphi(t, x) = x, & \forall (t, x) \in (-\varepsilon, \varepsilon) \times (M \setminus K), \end{cases}$$

and such that the velocity  $\partial\varphi(t, x)/\partial t$  is  $C^1$  on  $(\varepsilon, \varepsilon) \times M$ . Then the initial velocity vector  $X(y) = \partial\varphi(t, y)/\partial t|_{t=0}$  is a  $C^1$  vector field on  $M$  and we can write

$$5.2 \quad \varphi(t, y) = y + t(X(y) + E(t, y)),$$

where  $\sup_{y \in K} (|E(t, y)| + \sup_{\tau \in T_x M, |\tau| \leq 1} |D_\tau E(t, y)|) \rightarrow 0$  as  $t \rightarrow 0$ .

For example given any  $C^1$  vector function  $X : M \rightarrow \mathbb{R}^{n+k}$  with  $\{y \in M : X(y) \neq 0\}$  contained in a compact set  $K$  of  $M$ , we can construct such a  $\varphi$  simply by taking  $\varphi(t, y) = y + tX(y)$ ,  $|t| < 1$ .

We want to compute the first variation

$$\frac{d}{dt} \mathcal{H}^n(\varphi_t(M \cap K))|_{t=0},$$

where  $\varphi_t(y) = \varphi(t, y)$  with  $\varphi, K$  as in 5.1. To do this we first note that, by the area formula,

$$5.3 \quad \mathcal{H}^n(\varphi_t(M \cap K)) = \int_{M \cap K} J_{\varphi_t} d\mathcal{H}^n,$$

and, by 5.2,  $D_{\tau_i} \varphi(y) = \tau_i + t(D_{\tau_i} X(y) + E_i(t, y))$ , where  $\sup_{y \in K} |E_i(t, y)| \rightarrow 0$  as  $t \downarrow 0$ , and hence

$$J_{\varphi_t} = \sqrt{\det(\delta_{ij} + t(\tau_i \cdot D_{\tau_j} X(y) + \tau_j \cdot D_{\tau_i} X(y) + E_{ij}(t, y)))},$$

where  $\sup_{y \in K} |E_{ij}(t, y)| \rightarrow 0$  as  $t \rightarrow 0$ . Using the formula  $\det(I + tA) = 1 + t \operatorname{trace} A + O(t^2)$  as  $t \rightarrow 0$ , we thus conclude

$$\frac{d}{dt} J_{\varphi_t}(y)|_{t=0} = \sum_{i=1}^n \tau_i \cdot D_{\tau_i} X(y) = \operatorname{div}_M X(y).$$

Hence, by 5.3,

$$5.4 \quad \frac{d}{dt} \mathcal{H}^n(\varphi_t(M \cap K))|_{t=0} = \int_M \operatorname{div}_M X,$$

where  $X$  is the initial velocity vector of  $\varphi_t$ :  $X(y) = \frac{\partial}{\partial t} \varphi(t, y)|_{t=0}$ .

If we now decompose  $X$  into its tangent and normal parts:

$$X = X^\top + X^\perp$$

where (at least locally, in the notation introduced in 4.30 above)

$$X^\perp = \sum_{\alpha=1}^k (v^\alpha \cdot X) v^\alpha.$$

and  $X^\top = p_{T_x M}(X)$ . Then we have (near  $y$ )

$$\operatorname{div}_M X^\perp = \sum_{\alpha=1}^k (v^\alpha \cdot X) \operatorname{div} v^\alpha,$$

so that by 4.34

$$5.5 \quad \operatorname{div}_M X^\perp = -X \cdot \underline{H}$$

at each point of  $M$ . On the other hand  $\int_M \operatorname{div}_M X^\top = 0$  by 4.27. Hence, since  $\operatorname{div}_M X = \operatorname{div}_M X^\top + \operatorname{div}_M X^\perp$ , we obtain

$$5.6 \quad \int_M \operatorname{div}_M X \, d\mathcal{H}^n = - \int_M X \cdot \underline{H} \, d\mathcal{H}^n$$

for any  $C^1$  vector function  $X : M \rightarrow \mathbb{R}^{n+k}$  with  $\{y \in M : X(y) \neq 0\}$  contained in a compact subset  $K$  of  $M$ ; this identity is sometimes referred to as “the first variation formula” for the submanifold  $M$ .

**5.7 Remarks:** (1) Observe then that  $M$  is “stationary,” i.e. has first variation zero, i.e.  $\int_M \operatorname{div}_M X \, d\mathcal{H}^n = 0$  whenever  $X$  has compact support in  $M$  (which by 5.4 is equivalent to  $\frac{d}{dt} \mathcal{H}^n(\varphi_t(M \cap K))|_{t=0} = 0$  whenever  $\varphi$  is as in 5.1) if and only if the mean curvature

of  $M$  is identically zero. Such  $C^2$  submanifolds are usually referred to as “minimal submanifolds.”

(2) We should explain the appropriateness (or otherwise) of the terminology “minimal submanifold” introduced in the above remark: Observe that if  $M$  is *area minimizing* in the sense that  $\mathcal{H}^n(M \cap K) \leq \mathcal{H}^n(\varphi_t(M \cap K))$  whenever  $\varphi$  and  $K$  are as in 5.1, then  $\mathcal{H}^n(\varphi_t(M \cap K))$  has a minimum at  $t = 0$  and hence we do have stationarity  $\frac{d}{dt} \mathcal{H}^n(\varphi_t(M \cap K))|_{t=0} = 0$ . The converse is false though (as the example of the catenoid shows), so one could perhaps criticize the terminology “minimal” on this basis; nevertheless the converse is true (in the present context when  $M$  is  $C^2$ ) at least locally, i.e. provided we stipulate that the compact set  $K = B_\rho(y_0)$  in 5.1 above is a ball of sufficiently small radius  $\rho$  (depending on  $M$  and  $y_0 \in M$ ). See problem 7 of Ch. 2 problems.

(3) In case the situation is as in Remark 4.28(2), so that we drop the condition that  $\{y \in M : X(y) \neq 0\}$  is contained in a compact subset of  $M$  and instead assume  $\overline{M}$  (the closure of  $M$  in  $\mathbb{R}^{n+k}$ ) is a compact manifold with  $C^1$  boundary  $\partial M = \overline{M} \setminus M$  and  $X$  is any  $C^1$  vector field on  $\overline{M}$ , still with  $X(y) \in T_y M$  for each  $y \in M$ , then in place of 5.7 we get

$$\int_M \operatorname{div}_M X \, d\mathcal{H}^n = - \int_M X \cdot \underline{H} \, d\mathcal{H}^n - \int_{\partial M} X \cdot \eta \, d\mathcal{H}^{n-1}.$$

So far we have only discussed submanifolds of  $\mathbb{R}^{n+k}$ , and the concept of first variation using ambient space  $\mathbb{R}^{n+k}$ . For some applications it is important to allow the ambient space to be a complete  $(n+k)$ -dimensional Riemannian submanifold  $N$  rather than  $\mathbb{R}^{n+k}$ . By the Nash embedding theorem there is no loss of generality in assuming that  $N$  is (isometrically) embedded in  $\mathbb{R}^{n+L}$  for some  $L \geq k$ . So suppose  $N$  is a  $C^2$   $(n+k)$ -dimensional embedded submanifold of  $\mathbb{R}^{n+L}$ ,  $0 \leq k \leq L$ , let  $K$  be any compact subset of  $N$ , and let  $\varphi : (-\varepsilon, \varepsilon) \times N \rightarrow N$  (for some  $\varepsilon > 0$ ) be a  $C^1$  map with velocity vector  $\partial\varphi(t, y)/\partial t$  also  $C^1$  on  $(-\varepsilon, \varepsilon) \times M$ , and

$$5.8 \quad \varphi(0, y) = y \text{ for all } y \in N, \quad \varphi(t, y) = y \text{ for all } (t, y) \in (-\varepsilon, \varepsilon) \times N \setminus K.$$

Then we have the following definition for a  $C^2$  submanifold  $M$  of  $N$ :

**5.9 Definition:**  $M \subset N$  is a *stationary in  $N$*  (or “a minimal submanifold of  $N$ ”) if  $\frac{d}{dt} \mathcal{H}^n(\varphi_t(M \cap K))|_{t=0} = 0$  whenever  $K, \varphi$  are as in 5.8 and  $\varphi_t(y) = \varphi(t, y)$ .

**5.10 Remark:** In view of the fact that for each given compact  $K \subset M$  and each  $C^1$  vector field  $X$  on  $N$  with compact support  $K$  such that  $X_x \in T_x N$  at each  $x \in N$  (i.e.  $X$  is a *tangent* vector field on  $N$  with support in  $K$ ), there is (cf. the discussion in the proof of 4.27) a 1-parameter family  $\varphi_t$  as in 5.8 above with  $\frac{\partial}{\partial t} \varphi(y, t)|_{t=0} = X|_y$  at each point

$y \in N$ , we see that  $M$  is stationary in  $N$  as in 5.9 above if and only if

$$(\ddagger) \quad \int_M \operatorname{div}_M X = 0$$

whenever  $X$  is a  $C^1$  on  $N$  with compact support  $K$  and  $X|_y \in T_y N \forall y \in N$ .

If we let  $v^1, \dots, v^L$  be an orthonormal family (defined locally near a point  $y \in M$ ) of vector fields normal to  $M$ , such that  $v^1, \dots, v^k$  are tangent to  $N$  and  $v^{k+1}, \dots, v^L$  are normal to  $N$ , then for any vector field  $X$  on  $M$  we can write  $X = X^\top + X^\perp$ , where  $X_z^\top \in T_z N$  and  $X^\perp = \sum_{j=k+1}^L (v^j \cdot X) v^j$  (= the part of  $X$  normal to  $N$ ). Then if  $\tau_1, \dots, \tau_n$  is any orthonormal basis for  $T_y M$ , we have, at the point  $y$ ,

$$\begin{aligned} 5.11 \quad \operatorname{div}_M X &= \operatorname{div}_M X^\top + \sum_{j=k+1}^L (v^j \cdot X) \operatorname{div}_M v^j \\ &= \operatorname{div}_M X^\top + \sum_{j=k+1}^L (v^j \cdot X) \sum_{i=1}^n \tau_i \cdot D_{\tau_i} v^j \\ &= \operatorname{div}_M X^\top - \sum_{i=1}^n X \cdot B_y^N(\tau_i, \tau_i), \end{aligned}$$

where  $B_y^N$  is the second fundamental form of  $N$  at  $y$  and where we used the definition of second fundamental form as in 4.30 (with  $N$  in place of  $M$ ) and hence by virtue of Remark 5.10 (with  $X^\top$  in place of  $X$ ) we conclude:

**5.12 Lemma.** *If  $N$  is an  $(n+k)$ -dimensional embedded  $C^2$  submanifold of  $\mathbb{R}^{n+L}$  and if  $M \subset N$  is an  $n$ -dimensional embedded  $C^2$  submanifold of  $N$ , then  $M$  is stationary in  $N$  (i.e.  $\frac{d}{dt} \mathbb{M}(\varphi_t(M \cap K))|_{t=0} = 0$  whenever the  $\varphi_t$  and  $M \cap K$  are as in 5.8) if and only if*

$$\int_M \operatorname{div}_M X = - \int_M \underline{H}_M^N \cdot X$$

for each  $C^1$  vector field  $X$  with  $\{y \in M : X|_y \neq 0\}$  contained in a compact subset of  $M$ . Here

$$\underline{H}_M^N(y) = \sum_{i=1}^n B_y^N(\tau_i, \tau_i), \quad y \in M,$$

where  $B_y^N$  denotes the second fundamental form of  $N$  at  $y$  and  $\tau_1, \dots, \tau_n$  is any orthonormal basis of  $T_y M (\subset T_y N \subset \mathbb{R}^{n+L})$ .

## 6 Second Variation

We continue to suppose that  $M$  is an  $n$ -dimensional  $C^2$  embedded submanifold of  $\mathbb{R}^{n+k}$ . Also, let  $K \subset M$  be compact and  $\varphi : (-\varepsilon, \varepsilon) \times M \rightarrow \mathbb{R}^{n+k}$  as in 5.1 except that now we require also that  $\partial^2 \varphi(t, y) / \partial t^2$  exists and is  $C^1$  on  $(-\varepsilon, \varepsilon) \times M$ , so both the initial velocity  $X(y) = \partial \varphi(t, y) / \partial t|_{t=0}$  and the initial acceleration  $Z(x) = \partial^2 \varphi(t, y) / \partial t^2|_{t=0}$  will be  $C^1$  vector fields on  $M$ . Clearly then

$$6.1 \quad \varphi_t(x) = x + tX_x + \frac{1}{2}t^2(Z_x + E(t, x)) \text{ where } E(t, x) \rightarrow 0 \text{ as } t \rightarrow 0$$

in the sense that  $\sup_{x \in K} (|E(t, x)| + \sup_{\tau \in T_x M, |\tau|=1} |D_\tau E(t, x)|) \rightarrow 0$  as  $t \rightarrow 0$ .

Thus  $\{\varphi_t(M \cap K)\}_{|t| < \varepsilon}$  is a 1-parameter family of submanifolds, each with finite  $\mathcal{H}^n$ -measure, which agrees with  $M \cap K$  at  $t = 0$ . We computed the first variation  $\frac{d}{dt} \mathcal{H}^n(\varphi_t(M \cap K))|_{t=0}$  in the previous section: by 5.4 we have

$$6.2 \quad \frac{d}{dt} \mathcal{H}^n(\varphi_t(M \cap K))|_{t=0} = \int_M \operatorname{div}_M X.$$

Here we want to compute  $\frac{d^2}{dt^2} \mathcal{H}^n \varphi_t(M \cap K)|_{t=0}$  (i.e. the “second variation” of  $M$ ), which, as for the first variation, involves using the area formula

$$\mathcal{H}^n(\varphi_t(M \cap K)) = \int_M J_{\varphi_t} d\mathcal{H}^n.$$

This time we need to compute the terms up of second order in the Taylor series expansion (in the variable  $t$ ) of  $J_{\varphi_t}$ . With  $\tau_1, \dots, \tau_n$  any orthonormal basis for  $T_x M$ , we have by 6.1

$$D_{\tau_i} \varphi_t(x) = \tau_i + t D_{\tau_i} X + \frac{t^2}{2} (D_{\tau_i} Z + E_i(t, x))$$

for  $i = 1, \dots, n$ , where  $\sup_{x \in K \cap M} |E_i(t, x)| \rightarrow 0$  as  $t \rightarrow 0$ , so  $J_{\varphi_t}^2(x) = \det(D_{\tau_i} \varphi_t(x) \cdot D_{\tau_j} \varphi_t(x))$  has the form

$$\begin{aligned} (J_{\varphi_t}(x))^2 &= \delta_{ij} + t(\tau_i \cdot D_{\tau_j} X + \tau_j \cdot D_{\tau_i} X) \\ &\quad + t^2 \left( \frac{1}{2} (\tau_i \cdot D_{\tau_j} Z + \tau_j \cdot D_{\tau_i} Z) + (D_{\tau_i} X) \cdot (D_{\tau_j} X) + E_{ij}(t, x) \right), \end{aligned}$$

where  $\sup_{x \in K} |E_{ij}(t, x)| \rightarrow 0$  as  $t \rightarrow 0$ . By the general formula

$$\det(I + A) = 1 + \operatorname{trace} A + \frac{1}{2} (\operatorname{trace} A)^2 - \frac{1}{2} \operatorname{trace}(A^2) + O(|A|^3),$$

we then have

$$\begin{aligned} (J_{\varphi_t}(x))^2 &= 1 + 2t \operatorname{div}_M X + t^2 (\operatorname{div}_M Z + \sum_{i=1}^n |D_{\tau_i} X|^2 \\ &\quad + 2(\operatorname{div}_M X)^2 - \frac{1}{2} \sum_{i,j=1}^n (\tau_i \cdot D_{\tau_j} X + \tau_j \cdot D_{\tau_i} X)^2 + F(t, x)) \\ &= 1 + 2t \operatorname{div}_M X + t^2 (\operatorname{div}_M Z + \sum_{i=1}^n |(D_{\tau_i} X)^\perp|^2 \\ &\quad + 2(\operatorname{div}_M X)^2 - \sum_{i,j=1}^n (\tau_i \cdot D_{\tau_j} X)(\tau_j \cdot D_{\tau_i} X)) + t^2 F(t, x), \end{aligned}$$

where  $(D_{\tau_i} X)^\perp (= \text{the normal part of } D_{\tau_i} X) = D_{\tau_i} X - \sum_{j=1}^n (\tau_j \cdot D_{\tau_i} X) \tau_j$ , and where  $\sup_{x \in K} |F(t, x)| \rightarrow 0$  as  $t \rightarrow 0$ . Using  $\sqrt{1+x} = 1 + \frac{1}{2}x - \frac{1}{8}x^2 + O(x^3)$ , we thus get

$$\begin{aligned} J_{\varphi_t} &= 1 + t \operatorname{div}_M X + \frac{t^2}{2} (\operatorname{div}_M Z + (\operatorname{div}_M X)^2 + \sum_{i=1}^n |(D_{\tau_i} X)^\perp|^2 \\ &\quad - \sum_{i,j=1}^n (\tau_i \cdot D_{\tau_j} X)(\tau_j \cdot D_{\tau_i} X) + \widetilde{F}(t, x)), \end{aligned}$$

where again  $\sup_{x \in K} |\widetilde{F}(t, x)| \rightarrow 0$ . Thus

$$\begin{aligned} 6.3 \quad \frac{d^2}{dt^2} \mathcal{H}^n(\varphi_t(M \cap K))|_{t=0} &= \int_M (\operatorname{div}_M Z + (\operatorname{div}_M X)^2 + \sum_{i=1}^n |(D_{\tau_i} X)^\perp|^2 \\ &\quad - \sum_{i,j=1}^n (\tau_i \cdot D_{\tau_j} X)(\tau_j \cdot D_{\tau_i} X)) d\mathcal{H}^n. \end{aligned}$$

Finally, we shall need later the following important fact about the second variation formula 6.3.

**6.4 Lemma.** *If  $M$  is a  $C^2$  minimal submanifold (as in Definition 5.9) and if  $\{y \in M : X|_y \neq 0\}$  is contained in a compact subset of  $M$  with  $X|_y \in (T_y M)^\perp \forall y \in M$  (thus  $X|_M$  is a compactly supported tangent vector field on  $M$ ), then 6.3 says*

$$\left. \frac{d^2}{dt^2} \mathcal{H}^n(\varphi_t(M \cap K)) \right|_{t=0} = \int_M (\sum_{i=1}^n |(D_{\tau_i} X)^\perp|^2 - \sum_{i,j=1}^n (X \cdot B(\tau_i, \tau_j))^2) d\mathcal{H}^n.$$

**6.5 Remark:** In case  $k = 1$  and  $M$  is orientable, with continuous unit normal  $\nu$ , then  $X = \zeta \nu$  for some scalar function  $\zeta$  with compact support on  $M$ , and the above identity has the simple form

$$(*) \quad \left. \frac{d^2}{dt^2} \mathcal{H}^n(\varphi_t(M \cap K)) \right|_{t=0} = \int_M (|\nabla^M \zeta|^2 - \zeta^2 |B|^2) d\mathcal{H}^n,$$

where  $|B|^2 = \sum_{i,j=1}^n |B(\tau_i, \tau_j)|^2 \equiv \sum_{i,j=1}^n |\nu \cdot B(\tau_i, \tau_j)|^2$ . This is clear, because  $(D_{\tau_i}(\nu \zeta))^\perp = \nu D_{\tau_i} \zeta$  by virtue of the fact that  $D_{\tau_i} \nu|_y \in T_y M \forall y \in M$ .

**Proof of 6.4:** First we note that  $\int_M \operatorname{div}_M Z d\mathcal{H}^n = 0$  by virtue of the fact that  $M$  is stationary, and  $\operatorname{div}_M X = -X \cdot \underline{H} = 0$  by virtue of 5.5. The proof is then completed by noting that  $\tau_i \cdot D_{\tau_j} X = -X \cdot B(\tau_i, \tau_j)$  by virtue of the fact that  $X$  is normal to  $M$  together with 4.30.  $\square$

## 7 Co-Area Formula and $C^1$ Sard Theorem

Let  $M$  be a  $C^1$  submanifold of a Euclidean space  $\mathbb{R}^{n+\ell}$ , where  $\ell \in \{0, 1, \dots\}$  (note the case  $\ell = 0$  is included, which is the case when  $M$  is an open subset  $U$  of  $\mathbb{R}^n$ ), and let  $f = (f^1, \dots, f^m) : M \rightarrow \mathbb{R}^m$  be locally Lipschitz, with  $m \leq n$ , so that  $n = m + k$ ,  $k \in \{0, 1, \dots\}$ .

From 1.10 in case  $X = \mathbb{R}^{n+\ell}$ ,  $\mathcal{H}^k(M \cap f^{-1}z)$  is an  $\mathcal{H}^m$ -measurable function of  $z$  and

$$7.1 \quad \int_{\mathbb{R}^m} \mathcal{H}^k(A \cap f^{-1}z) d\mathcal{L}^m(z) \leq C \mathcal{H}^n(A), \quad C = C(n, \ell),$$

for each  $\mathcal{H}^n$ -measurable set  $A \subset M$ . The coarea formula, which we now present, enables us to replace this inequality with an exact identity. In the statement, we use the notation

$$J_f^M(x) = \sqrt{\det((df_x) \circ (df_x)^*)} = \sqrt{\det(\nabla^M f^i(x) \cdot \nabla^M f^j(x))},$$

when these quantities exist (which is  $\mathcal{H}^n$ -a.e. in  $M$  by 4.13); in terms of an orthonormal basis  $\tau_1, \dots, \tau_n$  for  $T_x M$  we thus have, by 4.22,

$$7.2 \quad J_f^M(x) = \sqrt{\det((\sum_{q=1}^n D_{\tau_q} f^i(x) D_{\tau_q} f^j(x))_{i,j=1,\dots,m})}.$$

**7.3 Theorem (Coarea Formula.)** *As above, let  $M$  be an  $n$ -dimensional  $C^1$  submanifold of  $\mathbb{R}^{n+\ell}$  and let  $f : M \rightarrow \mathbb{R}^m$  be locally Lipschitz with  $m = n - k$ , where  $k \in \{0, 1, 2, \dots\}$  (so  $m \leq n$  in contrast to the Area Formula 4.19, where we assumed  $m \geq n$ ). Then*

$$\int_{\mathbb{R}^m} \mathcal{H}^k(A \cap f^{-1}y) d\mathcal{L}^m(y) = \int_A J_f^M d\mathcal{H}^n$$

for any  $\mathcal{H}^n$ -measurable set  $A \subset M$ , where  $J_f^M$  is as in 7.2.

**7.4 Remarks: (1)** We then have the general formula

$$\int_M h J_f^M d\mathcal{H}^n = \int_{\mathbb{R}^m} \int_{f^{-1}(y)} h(x) d\mathcal{H}^k(x) d\mathcal{L}^m(y).$$

for any non-negative  $\mathcal{H}^n$ -measurable function  $h : M \rightarrow \mathbb{R}^m$ , which follows directly from the above theorem by approximating  $h$  pointwise (everywhere) by an increasing sequence of non-negative simple functions.

(2) Observe that if  $\ell = 0$  (so that  $M$  is an open subset  $U$  of  $\mathbb{R}^n$ ) and if  $f = p$ , where  $p : x = (x^1, \dots, x^n) \mapsto (x^1, \dots, x^m)$  is the projection onto the first  $m$  coordinates, then  $J_f^M \equiv 1$  and the above is just Fubini's Theorem. Thus the coarea formula can be viewed as a generalization of Fubini's Theorem. It is then not surprising that the proof given below depends in part of Fubini's Theorem.

**Proof of Theorem 7.3:** If  $n = m$  the result of the theorem is covered by the Area Formula 4.19, so we assume  $k = n - m \geq 1$ . Using the additivity of the relevant integrals on each side of the identity with respect to decompositions of  $A$  into pairwise disjoint unions, we can also assume without loss of generality that  $f$  is Lipschitz (rather than merely locally Lipschitz) and  $\mathcal{H}^n(A) < \infty$ .

The proof will be based on the Area Formula 4.19 and Fubini's Theorem.

Let  $M_+$  be the set of points  $x \in M$  such that  $T_x M$  and the directional derivatives  $D_\tau f(x)$  exist for all  $\tau \in T_x M$  and are linear in  $\tau \in T_x M$ , and recall that (by 4.13)

$$(1) \quad \mathcal{H}^n(M \setminus M_+) = 0.$$

We claim that for each  $\varepsilon > 0$  and each  $x \in M_+$  there is  $\delta_x > 0$  such that

$$(2) \quad z \in M \cap B_{\delta_x}(x) \setminus \{x\} \text{ with } f(z) = f(x) \Rightarrow |p_{K_x(f)}(z-x) - (z-x)| \leq \varepsilon|z-x|,$$

which follows from the fact that if  $x_j \in M \setminus \{x\}$  with  $x_j \rightarrow x$ , if  $f(x_j) = f(x) \forall j$  and if  $|x_j - x|^{-1}(x_j - x) \rightarrow v$ , then  $v \in T_x M$  and  $D_v f(x) = 0$ . (See problem 2.6 of Ch.2 problems.)

Since  $A = \{x \in A \cap M_+ : J_f(x) > 0\} \cup \{x \in A \cap M_+ : J_f(x) = 0\} \cup (A \cap (M \setminus M_+))$ , it suffices to consider just Case 1:  $A \subset \{x \in A \cap M_+ : J_f(x) > 0\}$ , Case 2:  $A \subset \{x \in A \cap M_+ : J_f(x) = 0\}$ , Case 3:  $A \subset M \setminus M_+$ .

In Case 1  $d_x^M f$  has rank  $m$  and  $K_x(f)$  has dimension  $k$  at each point of  $A$ . Let  $L$  be any  $k$ -dimensional subspace of  $\mathbb{R}^{n+\ell}$  with orthonormal basis  $\eta_1, \dots, \eta_k$ , let  $\varepsilon \in (0, \frac{1}{2})$ , define

$$(3) \quad A_{L,\varepsilon} = \{x \in A : \|p_{K_x(f)} - p_L\| < \varepsilon\}.$$

Then for each  $x \in A_{L,\varepsilon}$  there is an orthonormal basis  $\tau_1, \dots, \tau_n$  for  $T_x M$  with

$$(4) \quad K_x(f) = \text{span}\{\tau_{m+1}, \dots, \tau_n\} \text{ and } |\eta_i - \tau_{m+i}| < C\varepsilon, \quad i = 1, \dots, k,$$

with  $C = C(m, k)$ , and we can define  $g : M \rightarrow \mathbb{R}^n$  by

$$(5) \quad g(x) = (f(x), x \cdot \eta_1, \dots, x \cdot \eta_k), \quad x \in M.$$

Now  $J_g^M(x) = (\det(D_{\tau_i} g(x) \cdot D_{\tau_j} g(x)))^{1/2}$  and, by (4), for  $x \in A_{L,\varepsilon}$ ,

$$(D_{\tau_i} g(x) \cdot D_{\tau_j} g(x)) = \begin{pmatrix} (D_{\tau_i} f(x) \cdot D_{\tau_j} f(x))_{i,j=1,\dots,m} & O_{m \times k} \\ O_{k \times m} & I_{k \times k} \end{pmatrix} + E, \quad |E| \leq C\varepsilon,$$

where  $O_{k \times m}$  is the  $k \times m$  zero matrix,  $I_{k \times k}$  is the  $k \times k$  identity matrix and  $C = C(m, k, \text{Lip } f)$ . Also, by 7.2 and (4),  $J_f^M f(x) = (\det((D_{\tau_i} f(x) \cdot D_{\tau_j} f(x))_{i,j=1,\dots,m}))^{1/2}$ , so, for each  $x \in A_{L,\varepsilon}$ ,

$$(6) \quad J_g^M(x) = J_f^M(x) + e, \quad |e| \leq C\varepsilon, \quad C = C(m, k, \text{Lip } f).$$

Now, with  $\delta_x$  as in (2), we write

$$A_{L,\varepsilon,j} = \{x \in A_{L,\varepsilon} : \delta_x \geq 1/j\}, \quad j = 1, 2, \dots$$

Since  $|(\eta_1 \cdot y, \dots, \eta_k \cdot y)| = |p_L(y)| = |y + (p_{K_x(f)} y - y) + (p_L - p_{K_x(f)})y|$  for  $y \in \mathbb{R}^{n+\ell}$ , we then have, by (2), (3) and (5),

$$(7) \quad x_1, x_2 \in A_{L,\varepsilon,j} \text{ with } f(x_1) = f(x_2) \text{ and } |x_1 - x_2| < 1/j \Rightarrow \\ (1 - 2\varepsilon)|x_1 - x_2| \leq |g(x_1) - g(x_2)| \leq (1 + 2\varepsilon)|x_1 - x_2|.$$

Next, for any  $\mathcal{H}^n$ -measurable subset  $B \subset A_{L,\varepsilon}$ , choose pairwise disjoint  $\mathcal{H}^n$ -measurable subsets  $B_1, B_2, \dots$  of  $A_{L,\varepsilon,j}$  with diameter  $B_i < 1/j$  for each  $i$  and  $\cup_i B_i = B \cap A_{L,\varepsilon,j}$ . Then, by the Area Formula 4.19 and Fubini's Theorem,

$$\begin{aligned}
 (8) \quad \int_{B_i} J_g^M d\mathcal{H}^n &= \mathcal{H}^n(g(B_i)) \\
 &= \int_{\mathbb{R}^m} \mathcal{H}^k(g(B_i) \cap (\{y\} \times \mathbb{R}^k)) dy \\
 &= \int_{\mathbb{R}^m} \mathcal{H}^k(g(B_i \cap f^{-1}y)) dy,
 \end{aligned}$$

and, since  $\text{Lip } g|_{B_i \cap f^{-1}y} \leq (1 + 2\varepsilon)$  and  $\text{Lip}(g|_{B_i \cap f^{-1}y})^{-1} \leq (1 - 2\varepsilon)^{-1}$  by (7), we use 1.8 to conclude

$$(1 - 2\varepsilon)^k \mathcal{H}^k(B_i \cap f^{-1}y) \leq \mathcal{H}^k(g(B_i \cap f^{-1}y)) \leq (1 + 2\varepsilon)^k \mathcal{H}^k(B_i \cap f^{-1}y).$$

Hence, in view of (6), (8) gives

$$\left| \int_{B_i} J_f^M d\mathcal{H}^n - \int_{\mathbb{R}^m} \mathcal{H}^k(B_i \cap f^{-1}y) dy \right| \leq C\varepsilon \mathcal{H}^n(B_i), \quad C = C(m, k, \text{Lip } f).$$

Hence, by first summing on  $i$  and then letting  $j \rightarrow \infty$ , we obtain, for any  $\mathcal{H}^n$ -measurable  $B \subset A_{L,\varepsilon}$ ,

$$(9) \quad \left| \int_B J_f^M d\mathcal{H}^n - \int_{\mathbb{R}^m} \mathcal{H}^k(B \cap f^{-1}y) dy \right| \leq C\varepsilon \mathcal{H}^n(B)$$

Now, still with  $\varepsilon \in (0, \frac{1}{2})$  arbitrary, let  $L_1, \dots, L_N$  ( $N = N(n, \ell, \varepsilon)$ ) be  $k$ -dimensional subspaces of  $\mathbb{R}^{n+\ell}$  such that for every  $k$ -dimensional subspace  $L$  of  $\mathbb{R}^{n+\ell}$  there is  $j \in \{1, \dots, N\}$  with  $|p_L - p_{L_j}| < \varepsilon$ . Then we can decompose  $A$  into a disjoint union  $\cup_{j=1}^N A_j$  of  $\mathcal{H}^n$ -measurable subsets such that  $A_j \subset A_{L_j,\varepsilon}$  for each  $j$ , and hence, using (9) with  $L_j$  in place of  $L$  and  $A_j$  in place of  $B$ , we have

$$\left| \int_{A_j} J_f^M d\mathcal{H}^n - \int_{\mathbb{R}^m} \mathcal{H}^k(A_j \cap f^{-1}y) dy \right| \leq C\varepsilon \mathcal{H}^n(A_j),$$

and the required identity follows by first summing over  $j$  and then letting  $\varepsilon \downarrow 0$ . The completes the proof of Case 1.

In Case 2 we are assuming  $A \subset \{x \in M_+ : J_f^M(x) = 0\}$ , and we can apply Case 1 with  $\widetilde{M} = M \times (0, 1)^m$ ,  $\widetilde{A} = A \times (0, 1)^m$  and  $m + k$  in place of  $M$ ,  $A$  and  $k$  respectively, and  $F : \widetilde{M} \rightarrow \mathbb{R}^m$  in place of  $f$ , where

$$F(x, z) = f(x) + \varepsilon z, \quad x \in M, \quad z \in (0, 1)^m \quad (\varepsilon \in (0, 1)).$$

Notice that  $d_{x,z}^{\widetilde{M}}F$  evidently has rank  $m$  at each point  $(x, z) \in \widetilde{M}$ , so Case 1 is indeed applicable and gives

$$(10) \quad \int_{\mathbb{R}^m} \mathcal{H}^{m+k}(\widetilde{A} \cap F^{-1}y) d\mathcal{L}^m(y) = \int_{\widetilde{A}} J_F^{\widetilde{M}} d\mathcal{H}^{m+n}.$$

Using the formula  $J_F^{\widetilde{M}}(x, z) = \sqrt{\det(\sum_{q=1}^{n+m} D_{\eta_q} F^i(x, z) \cdot D_{\eta_q} F^j(x, z))}$  (Cf. 7.2), where  $\eta_q = (\tau_q, 0)$  for  $q = 1, \dots, n$  and  $\eta_{n+q} = (0, e_q)$  for  $q = 1, \dots, m$ , with  $\tau_1, \dots, \tau_n$  an orthonormal basis for  $T_x M$  and  $e_1, \dots, e_m$  the standard basis for  $\mathbb{R}^m$ , we see that

$$\begin{aligned} J_F^{\widetilde{M}}(x, z) &= \sqrt{\det((\sum_{q=1}^n D_{\tau_q} f^i(x) D_{\tau_q} f^j(x) + \varepsilon^2 \delta_{ij})_{i,j=1,\dots,m})} \\ &= J_f^M(x) + E(x, z) \text{ by 7.2,} \end{aligned}$$

where  $|E(x, z)| \leq C\varepsilon$ ,  $C = C(n, \ell, \text{Lip } f)$ . Since  $J_f^M|_A = 0$ , (10) then implies

$$(11) \quad \int_{\mathbb{R}^m} \mathcal{H}^{m+k}(\widetilde{A} \cap F^{-1}y) d\mathcal{L}^m(y) \leq C\varepsilon \mathcal{H}^n(A)$$

With  $p$  is the projection  $p : (x, z) \in \mathbb{R}^{n+\ell} \times \mathbb{R}^m \mapsto z \in \mathbb{R}^m$  we have

$$(\widetilde{A} \cap F^{-1}y) \cap p^{-1}(z) = \{(x, z) : x \in A, f(x) = y - \varepsilon z\} = (A \cap f^{-1}(y - \varepsilon z)) \times \{z\}$$

for  $z \in (0, 1)^m$ , so, by 7.1 with  $\widetilde{A} \cap F^{-1}y$  in place of  $A$  and  $p$  in place of  $f$ ,

$$\int_{(0,1)^m} \mathcal{H}^k(A \cap f^{-1}(y - \varepsilon z)) d\mathcal{L}^m(z) \leq C \mathcal{H}^{m+k}(\widetilde{A} \cap F^{-1}y), \quad y \in \mathbb{R}^m.$$

Integrating this inequality with respect to  $y \in \mathbb{R}^m$ , using Fubini's Theorem to change the order of integration on the left, and noting that  $\int_{\mathbb{R}^m} \mathcal{H}^k(A \cap f^{-1}(y - \varepsilon z)) d\mathcal{L}^m(y) = \int_{\mathbb{R}^m} \mathcal{H}^k(A \cap f^{-1}(y)) d\mathcal{L}^m(y)$  (by change of variable  $y \mapsto y - \varepsilon z$ ), we conclude

$$\int_{\mathbb{R}^m} \mathcal{H}^k(A \cap f^{-1}(y)) d\mathcal{L}^m(y) \leq \int_{\mathbb{R}^m} \mathcal{H}^{m+k}(\widetilde{A} \cap F^{-1}y) d\mathcal{L}^m(y) \leq C\varepsilon \mathcal{H}^n(A) \quad (\text{by (11)}).$$

So the Coarea Formula (with both sides = 0) is proved in Case 2 by letting  $\varepsilon \downarrow 0$ .

In Case 3,  $A \subset M \setminus M_+$ , so  $\mathcal{H}^n(A) = 0$  by (1), and the Coarea Formula holds (again with each side = 0) by virtue of 7.1.

This completes the proof of the Coarea Formula.  $\square$

In case  $f : M \rightarrow \mathbb{R}^m$  is  $C^1$  (rather than merely locally Lipschitz) with  $m < n$ , there is an important additional consequence of 7.3: namely if  $C = \{x \in M : J_f^M(x) = 0\}$ , then (by using 7.3 with  $A = C$ )  $\mathcal{H}^k(C \cap f^{-1}(y)) = 0$  for  $\mathcal{L}^m$ -a.e.  $y \in \mathbb{R}^m$ . Also, since

$J_f^M(x) \neq 0$  precisely when  $d_x^M f$  has rank  $m$ , the implicit function theorem implies that either  $f^{-1}(y) \setminus C$  is empty or else is a  $k$ -dimensional  $C^1$  embedded submanifold in the sense of §4 above.

In summary we thus have the following important result.

**7.5 Theorem ( $C^1$  Sard-type Theorem.)** *Suppose  $f : M \rightarrow \mathbb{R}^m$ ,  $m \leq n$ , is  $C^1$ , with  $M$  is an  $n$ -dimensional  $C^1$  embedded submanifold of  $\mathbb{R}^N$ . Then for  $\mathcal{L}^m$ -a.e.  $y \in f(M)$ ,  $f^{-1}(y)$  decomposes into a  $k$ -dimensional  $C^1$  embedded submanifold and a closed set of  $\mathcal{H}^k$ -measure zero, where  $k = n - m$ . Specifically,*

$$f^{-1}(y) = (f^{-1}(y) \setminus C) \cup (f^{-1}(y) \cap C),$$

$C = \{x \in M : J_f^M(x) = 0\} (\equiv \{x \in M : \text{rank}(df_x) < m\})$ ,  $\mathcal{H}^k(f^{-1}(y) \cap C) = 0$ ,  $\mathcal{L}^m$ -a.e.  $y \in \mathbb{R}^m$ , and  $f^{-1}(y) \setminus C$  is either empty or an  $k$ -dimensional  $C^1$  embedded submanifold.

**7.6 Remark:** If  $f$  and  $M$  are of class  $C^{k+1}$ , then *Sard's Theorem* asserts the stronger result that in fact  $f^{-1}(y) \cap C = \emptyset$  for  $\mathcal{L}^m$ -a.e.  $y \in \mathbb{R}^m$ , so that  $f^{-1}(y)$  is either empty or a  $k$ -dimensional  $C^{k+1}$  embedded submanifold for  $\mathcal{L}^m$ -a.e.  $y \in \mathbb{R}^m$ .

We conclude this section with some important remarks about selection of “good” slices by a given locally Lipschitz function  $f : M \rightarrow \mathbb{R}^m$ :

**7.7 Remarks:** (1) First notice that the formula 7.4(1) enables us to bound the  $\mathcal{H}^k$  measure of the “slices”  $f^{-1}y$  for a good set of  $y$ . Specifically if  $|f| \leq R$  and  $g$  is as in 7.4 ( $g \equiv 1$  is an important case), then there must be set  $S \subset B_R(0) (\subset \mathbb{R}^m)$ ,  $S = S(g, f, M)$ , with  $\mathcal{L}^m(S) \geq \frac{1}{2}\mathcal{L}^m(B_R(0))$  and with

$$\int_{f^{-1}(y)} g d\mathcal{H}^k \leq \frac{2}{\mathcal{L}^m(B_R(0))} \int_M g J_f^* d\mathcal{H}^n$$

for each  $y \in S$ . For otherwise there would be a set  $T \subset B_R(0)$  with  $\mathcal{L}^m(T) > \frac{1}{2}\mathcal{L}^m(B_R(0))$  and

$$\int_{f^{-1}(y)} g d\mathcal{H}^k \geq \frac{2}{\mathcal{L}^m(B_R(0))} \int_M g J_f^* d\mathcal{H}^n, \quad y \in T,$$

so that, integrating over  $T$  we obtain a contradiction to 7.4 if  $\int_M g J_f^* d\mathcal{H}^n > 0$ . On the other hand if  $\int_M g J_f^* d\mathcal{H}^n = 0$  then the required result is a trivial consequence of 7.4.

(2) The above has an important extension to the case when we have  $f : \mathbb{R}^N \rightarrow \mathbb{R}^n$  and sequences  $\{M_j\}$ ,  $\{g_j\}$  satisfying the conditions of  $M$ ,  $g$  above. In this case there is a set  $S \subset B_R(0)$  with  $\mathcal{L}^m(S) \geq \frac{1}{2}\mathcal{L}^m(B_R(0))$  such that for each  $y \in S$  there is a *subsequence*

$\{j'\}$  (depending on  $y$ ) with

$$\int_{M_{j'} \cap f^{-1}(y)} g_{j'} d\mathcal{H}^k \leq \frac{2}{\mathcal{L}^m(B_R(0))} \int_{M_{j'}} g_{j'} J_f^* d\mathcal{H}^n.$$

Indeed otherwise there is a set  $T$  with  $\mathcal{L}^m(T) > \frac{1}{2}\mathcal{L}^m(B_R(0))$  so that for each  $y \in T$  there is  $\ell(y)$  such that

$$(\ddagger) \quad \int_{M_j \cap f^{-1}(y)} g_j d\mathcal{H}^k > \frac{2}{\mathcal{L}^m(B_R(0))} \int_{M_j} g_j J_f^* d\mathcal{H}^n$$

for each  $j > \ell(y)$ . But  $T = \cup_{j=1}^{\infty} T_j$ ,  $T_j = \{y \in T : \ell(y) \leq j\}$ , and hence there must exist  $j$  so that  $\mathcal{L}^m(T_j) > \frac{1}{2}\mathcal{L}^m(B_R(0))$ . Then, integrating  $(\ddagger)$  over  $y \in T_j$ , we obtain a contradiction to 7.4 as before.

## CHAPTER 2 PROBLEMS

2.1  $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$  is said to be approximately differentiable at  $x \in \mathbb{R}^m$  with respect to Lebesgue measure if there is a linear map  $v_x : \mathbb{R}^m \rightarrow \mathbb{R}^n$  such that

$$\lim_{\rho \downarrow 0} \rho^{-m} \mathcal{L}^m(\{y \in B_\rho(x) \setminus \{x\} : |y-x|^{-1}|f(y) - f(x) - v_x(y-x)| \geq \varepsilon\}) = 0 \quad \forall \varepsilon > 0.$$

If  $f$  is locally Lipschitz near  $x$  (i.e.  $\exists K, R > 0$  such that  $|f(z) - f(y)| \leq K|z - y|$  for all  $y, z \in B_R(x)$ ), prove that approximate differentiability at  $x$  is equivalent to differentiability at  $x$ .

(Recall differentiability at  $x$  means that there is a linear map  $v_x : \mathbb{R}^m \rightarrow \mathbb{R}^n$  such that  $\lim_{y \rightarrow x} |x - y|^{-1}|f(y) - f(x) - v_x(y - x)| = 0$ ; i.e. for each  $\varepsilon > 0$  there is  $\delta > 0$  such that  $|x - y|^{-1}|f(y) - f(x) - v_x(y - x)| < \varepsilon$  for all  $y \in B_\delta(x) \setminus \{x\}$ .)

2.2 (Tietze extension theorem) Assume  $X$  is an arbitrary metric space,  $A \subset X$  is closed,  $\lambda > 0$ , and  $f : A \rightarrow \mathbb{R}$  is bounded continuous with  $\sup_A |f| \leq \lambda$ .

(i) Let  $A_+ = \{x : f(x) \geq \lambda/3\}$ ,  $A_- = \{x : f(x) \leq -\lambda/3\}$  and check that if  $A_\pm$  are non-empty then  $h_1(x) = \frac{\lambda}{3}(d(x, A_-) - d(x, A_+))/(d(x, A_+) + d(x, A_-))$  defines a continuous function  $h_1 : X \rightarrow [-\lambda/3, \lambda/3]$  on  $X$  such that  $h_1 \equiv \lambda/3$  on  $A_+$  and  $h_1 \equiv -\lambda/3$  on  $A_-$ ; note also that such a function  $h_1$  is obtained by taking  $h_1 \equiv \lambda/3$  if  $A_- = \emptyset$  and  $h_1 \equiv -\lambda/3$  if  $A_+ = \emptyset$ .

(ii) Prove by (i) and induction on  $k$  that for each  $k = 1, 2, \dots$  there exist continuous  $h_1, h_2, \dots, h_k$  on  $X$  such that  $\sup_A |f - \sum_{j=1}^k h_j| \leq (2/3)^k \lambda$  and  $|h_j| \leq 2^{j-1} \lambda/3^j$  on  $X$ ,  $j = 1, 2, \dots, k$ .

(iii) By letting  $k \rightarrow \infty$  in (ii), prove there is a continuous  $H : X \rightarrow [-\lambda, \lambda]$  such that  $H|_A = f$ .

(iv) Prove there is a continuous  $H : X \rightarrow \mathbb{R}$  with  $H|_A = f$  even if no boundedness hypothesis is assumed for  $f$ .

Hint for (iv): Start by applying (iii) with  $\arctan f$  in place of  $f$ . Caution: In this case (iii) would be applied with  $\lambda = \pi/2$  and the extension  $H$  may possibly have a non-empty set  $C = \{x : |H(x)| = \pi/2\}$ , so you cannot get the required extension for  $f$  simply by using  $\tan H$ . (Note however that  $C$  is a closed set disjoint from  $A$ .)

2.3 Suppose  $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$  is continuous on  $\mathbb{R}^m$  and satisfies  $\limsup_{y \rightarrow x} |y - x|^{-1}|f(y) - f(x)| < \infty$  at almost all points  $x \in \mathbb{R}^m$  (that is,  $\lim_{\rho \downarrow 0} \sup_{0 < |y-x| < \rho} |y - x|^{-1}|f(y) - f(x)| < \infty$  a.e.)

(i) If  $C_j = \{x : |f(y) - f(x)| \leq j|y - x| \text{ whenever } |y - x| < 1/j\}$ , prove that  $C_j$  is closed and that  $\mathcal{L}^n(\mathbb{R}^n \setminus (\cup_j C_j)) = 0$ .

(ii) Let  $\cup_i C_{j,i}$  be a decomposition of  $C_j$  into closed (not necessarily disjoint) subsets of diameter  $< 1/j$ . Prove that  $f|_{C_{j,i}}$  is Lipschitz.

(iii) Prove that  $f$  is approximately differentiable (see 2.1 above) at  $\mathcal{L}^n$ -a.e. point  $x \in \mathbb{R}^n$

2.4 (Chain rule for composite of a Lipschitz and an AC function.) If  $g : \mathbb{R} \rightarrow \mathbb{R}$  is Lipschitz and  $f : [a, b] \rightarrow \mathbb{R}$  is absolutely continuous (AC) (hence both  $f, g \circ f$  are AC on  $(a, b)$  and so both  $(g \circ f)'(x), f'(x)$  exist a.e.  $x \in (a, b)$ ), prove:

(i)  $x \in (a, b)$  and  $f'(x) = 0 \Rightarrow (g \circ f)'(x)$  exists and is equal to zero.

(ii) If  $\gamma(y) = g'(y)$  when this exists, and  $\gamma(y) = 0$  at points  $y$  where  $g'(y)$  does not exist, prove that  $(g \circ f)'(x) = \gamma(f(x))f'(x)$  for a.e.  $x \in (a, b)$ .

Hint for (ii): By (i),  $(g \circ f)'(x)$  exists and is equal to zero at all points of  $F_0 = \{x \in (a, b) : f'(x) = 0\}$ . Then show, directly by using difference quotients, that  $(g \circ f)'(x)$  and  $g'(f(x))f'(x)$  both exist and are equal at every point of  $(a, b) \setminus (E \cup F_0)$ , where  $E = \{x \in (a, b) : f'(x) \text{ does not exist}\} \cup \{x \in (a, b) : (g \circ f)'(x) \text{ does not exist}\}$ .

2.5 Let  $X, Y$  be metric spaces with  $X$   $\sigma$ -compact and let  $f : A \rightarrow Y$  be Lipschitz with  $A \subset X$  and  $A = \bigcup_{j=1}^{\infty} A_j$  with  $\mathcal{H}^m(A_j) < \infty$  for each  $j$ .

(i) Prove that  $\mathcal{H}^m(\{y : \mathcal{H}^0(A_j \cap f^{-1}y) = \infty\}) = 0$  for each  $j$ .

(ii) Give an example to show that  $\mathcal{H}^m(\{y : \mathcal{H}^0(A \cap f^{-1}y) = \infty\}) = \infty$  is possible with the stated hypotheses.

2.6 Let  $M$  be an  $n$ -dimensional embedded  $C^1$  submanifold of  $\mathbb{R}^{n+k}$ , and  $x \in M$ .

(i) If  $x_j \in M \setminus \{x\}$  with  $x_j \rightarrow x$  and  $|x_j - x|^{-1}(x_j - x) \rightarrow v \in \mathbb{S}^{n+k-1}$ , prove  $v \in T_x M$ .

(ii) If  $f : \mathbb{R}^{n+k} \rightarrow \mathbb{R}^m$  is Lipschitz, if  $x_j, v$  are as in (i) above with  $f(x_j) = f(x)$  for each  $j$ , then  $D_v f(x) = 0$ . Note: Part of what is to be proved is that  $D_v f(x)$  exists.

2.7 Suppose  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is Lipschitz and  $\int_{\mathbb{R}^n} |Df| \leq 1$ . Prove that for each  $K > 0$  the slices  $\{x \in \mathbb{R}^n : f(x) = t\}$  have  $(n-1)$ -dimensional Hausdorff measure  $\leq K$  with the exception of a set of  $t$  of Lebesgue measure  $\leq 1/K$ . Hint: Coarea formula.

2.8 Suppose  $X$  is a  $\sigma$ -compact metric space,  $m \in \{1, 2, \dots\}$ ,  $k > 0$ ,  $A \subset X$  is  $\mathcal{H}^{m+k}$ -measurable with  $\mathcal{H}^{m+k}(A) < \infty$ , and  $f : A \rightarrow \mathbb{R}^m$  is continuous. Prove

(i) If  $A$  is compact and  $U$  is an open subset of  $X$ , then  $\{y \in X : f^{-1}y \subset U\}$  is an open subset of  $\mathbb{R}^m$ .

(ii) If  $A$  is compact then  $\{y \in \mathbb{R}^m : \mathcal{H}_\delta^k(f^{-1}(y)) < t\}$  is open for each  $t \in \mathbb{R}$  and each  $\delta > 0$ .

(iii)  $\mathcal{H}^k(f^{-1}y)$  is an  $\mathcal{H}^m$ -measurable function of  $y \in \mathbb{R}^m$ , provided  $f : A \rightarrow \mathbb{R}^m$  is Lipschitz.

Hint for (iii): Using the inequality (4) in the proof of Theorem 1.10, start by showing that if  $\mathcal{H}^{m+k}(A) = 0$  then  $\mathcal{H}^k(f^{-1}y) = 0$  for  $\mathcal{L}^m$ -a.e.  $y \in \mathbb{R}^m$ .

2.9 Let  $\gamma_j : [a_j, b_j] \rightarrow \mathbb{R}^n$  be absolutely continuous and such that  $\gamma_j(a_j) = 0$  and length  $\gamma_j = 1$  for each  $j = 1, 2, \dots$

(i) Prove there is a Lipschitz map  $\gamma : [0, 1] \rightarrow \mathbb{R}^n$  with  $\text{Lip } \gamma \leq 1$  such that a subsequence of  $\gamma_j([a_j, b_j]) \rightarrow \gamma([0, 1])$  in the Hausdorff distance sense.

Note on terminology: Given sets  $A, B$  in a metric space  $X$ , the Hausdorff distance between  $A, B$  is defined as the inf of the set of  $\lambda > 0$  such that  $A \subset \{x \in X : d(x, B) < \lambda\}$  and  $B \subset \{x \in X : d(x, A) < \lambda\}$ .

(ii) Construct an example of a sequence  $\gamma_j$  which shows that  $\gamma$  in (i) may have length strictly less than 1.

2.10 For each  $N = 2, 3, \dots$  let  $M_N$  be the 2-dimensional  $C^\infty$  submanifold of  $\mathbb{R}^3$  defined by

$$M_N = \cup_{j,k \in \{0, \pm 1, \pm 2, \dots\}} \{(x, y) \in \mathbb{R}^2 \times \mathbb{R} : |x - (j/N, k/N)| = 1/N^2\}.$$

(Thus  $M_N$  is a countable pairwise disjoint union of cylinders with axes parallel to the third coordinate axis.)

Prove that  $\mathcal{H}^2 \llcorner M_N \rightarrow 2\pi \mathcal{L}^3$  (i.e.  $2\pi$  times Lebesgue measure on  $\mathbb{R}^3$ ) as  $N \rightarrow \infty$ ; i.e. prove

$$\int_{M_N} f d\mathcal{H}^2 \rightarrow 2\pi \int_{\mathbb{R}^3} f d\mathcal{L}^3 \text{ for each } f \in C_c^0(\mathbb{R}^3).$$

Hint: First show that  $N^{-2} \sum_{j,k=0, \pm 1, \pm 2, \dots} \int_{\mathbb{R}} f(j/N, k/N, y) dy \rightarrow \int_{\mathbb{R}^3} f d\mathcal{L}^3$  for each  $f \in C_c^0(\mathbb{R}^3)$ .

2.11 With  $M_N$  as in Q 2.10 above, prove that  $\int_{M_N} \omega \rightarrow 0$  as  $N \rightarrow \infty$  for each continuous 2-form  $\omega$  on  $\mathbb{R}^3$  with compact support in  $\mathbb{R}^3$ .

Note: Here we use the usual definition of  $\int_M \omega$  for a 2-dimensional oriented  $C^1$  submanifold of  $\mathbb{R}^3$  and a continuous 2-form  $\omega = \omega_1 dx^2 \wedge dx^3 + \omega_2 dx^1 \wedge dx^3 + \omega_3 dx^1 \wedge dx^2$ ; namely, we assume that we have selected a continuous unit normal  $\nu = (\nu_1, \nu_2, \nu_3)$  for  $M$  and then  $\int_M \omega = \int_M \omega^* \cdot \nu d\mathcal{H}^2$ , where  $\omega^* = (\omega_1, -\omega_2, \omega_3)$  the vector field dual to  $\omega$ .

# Chapter 3

## Countably $n$ -Rectifiable Sets

§ 1	Basic Notions, Tangent Properties . . . . .	85
§ 2	Gradients, Jacobians, Area, Co-Area . . . . .	92
§ 3	Purely Unrectifiable Sets, Structure Theorem . . . . .	94
§ 4	Sets of Locally Finite Perimeter . . . . .	96

### 1 Basic Notions, Tangent Properties

Firstly, a set  $M \subset \mathbb{R}^{n+k}$  is said to be countably  $n$ -rectifiable if

$$1.1 \quad M \subset M_0 \cup \left( \bigcup_{j=1}^{\infty} F_j(\mathbb{R}^n) \right),$$

where  $\mathcal{H}^n(M_0) = 0$  and  $F_j : \mathbb{R}^n \rightarrow \mathbb{R}^{n+k}$  are Lipschitz functions for  $j = 1, 2, \dots$ <sup>1</sup>  
 Notice also that by the extension theorem 1.2 of Ch.2 this is equivalent to saying

$$M = M_0 \cup \left( \bigcup_{j=1}^{\infty} F_j(A_j) \right)$$

where  $\mathcal{H}^n(M_0) = 0$ ,  $F_j : A_j \rightarrow \mathbb{R}^{n+k}$  Lipschitz,  $A_j \subset \mathbb{R}^n$ . More importantly, we have the following lemma.

**1.2 Lemma.**  *$M$  is countably  $n$ -rectifiable if and only if  $M \subset \bigcup_{j=0}^{\infty} N_j$ , where  $\mathcal{H}^n(N_0) = 0$  and where each  $N_j$ ,  $j \geq 1$ , is an  $n$ -dimensional embedded  $C^1$  submanifold of  $\mathbb{R}^{n+k}$ .*

**Proof:** The “if” part is essentially trivial because if  $N$  is an  $n$ -dimensional  $C^1$  embedded submanifold, then using local representations for  $N$  as in Remark 4.2(3) of Ch.2 we see

---

<sup>1</sup>Notice that this differs slightly from the terminology of [Fed69] in that we allow a set  $M_0$  with  $\mathcal{H}^n(M_0) = 0$ .

that for each  $x \in N$  there is  $\rho_x > 0$  such that  $B_{\rho_x}(x) \cap N = \psi(V)$  for suitable  $C^1$  map  $\psi : V \rightarrow \mathbb{R}^{n+k}$ ,  $V \subset \mathbb{R}^n$  open. Since such  $C^1$  maps are automatically Lipschitz in each closed ball  $\subset V$  it is then clear that  $M$  satisfies the definition 1.1.

The “only if” part is a consequence of the  $C^1$  Approximation Theorem 1.5 of Ch.2, which guarantees that if  $F_j$  are Lipschitz functions as in the Definition 1.1 above, then for each  $j \in \{1, 2, \dots\}$  we can choose  $C^1$  functions  $G_{1j}, G_{2j}, \dots : \mathbb{R}^n \rightarrow \mathbb{R}^k$  such that  $\mathcal{H}^n(\{x : F_j(x) \neq G_{ij}(x)\}) < 1/i$ . So, with

$$Z_j = \mathbb{R}^n \setminus (\cup_{i=1}^{\infty} \{x : F_j(x) = G_{ij}(x)\}),$$

we have  $\mathcal{H}^n(Z_j) = 0$ , in which case

$$(1) \quad F_j(\mathbb{R}^n) \subset F_j(Z_j) \cup (\cup_{i=1}^{\infty} G_{ij}(\mathbb{R}^n)), \quad j = 1, 2, \dots$$

Then  $\mathcal{H}^n(F_j(Z_j)) = 0$  because  $F_j$  is Lipschitz and  $\mathcal{H}^n(Z_j) = 0$ , so

$$(2) \quad \mathcal{H}^n(N_0) = 0, \text{ where } N_0 = (\cup_{j=1}^{\infty} F_j(Z_j)),$$

and we have proved

$$M \subset M_0 \cup N_0 \cup (\cup_{i,j=1}^{\infty} G_{ij}(\mathbb{R}^n)).$$

Let  $C_{ij}$  be the critical set of  $G_{ij}$ ; i.e.  $C_{ij} = \{x \in \mathbb{R}^n : J_{G_{ij}}(x) = 0\}$ . By the area formula  $\mathcal{H}^n(G_{ij}(C_{ij})) = 0$ , whereas if  $x \in \mathbb{R}^n \setminus C_{ij}$ , then by an inverse function theorem argument similar to that in Remark 4.2(2), there is a  $\rho > 0$  such that  $G_{ij}(\check{B}_{\rho}(x))$  is an  $n$ -dimensional  $C^1$  embedded submanifold of  $\mathbb{R}^n$  (with  $G_{ij}|_{\check{B}_{\rho}(x)}$  providing a local representation in a neighborhood of the point  $y = G_{ij}(x)$ ). So  $\cup_{ij} G_{ij}(\mathbb{R}^n)$  can be written as the union of a set of measure zero and countably many  $n$ -dimensional  $C^1$  embedded submanifolds of  $\mathbb{R}^{n+k}$ .  $\square$

**1.3 Remark:** If  $M$  is countably  $n$ -rectifiable, the above lemma guarantees that we can find  $N_0$  with  $\mathcal{H}^n$  measure zero and  $n$ -dimensional  $C^1$  embedded submanifolds  $N_1, N_2, \dots$  with  $M \subset \cup_{j=0}^{\infty} N_j$ , and so we can write  $M$  as a disjoint union  $M = \cup_{j=0}^{\infty} M_j$  with  $M_j \subset N_j$  for each  $j = 0, 1, 2, \dots$ . To achieve this, just define the  $M_j$  inductively by  $M_0 = M \cap N_0$  and  $M_j = M \cap N_j \setminus \cup_{i=0}^{j-1} M_i$ ,  $j \geq 1$ . Of course the sets  $M_j$  so constructed are all  $\mathcal{H}^n$ -measurable if  $M$  is.

We now want to give an important characterization of countably  $n$ -rectifiable sets in terms of *approximate tangent spaces*, which we first define:

**1.4 Definition:** If  $M$  is an  $\mathcal{H}^n$ -measurable subset of  $\mathbb{R}^{n+k}$  with  $\mathcal{H}^n(M \cap K) < \infty \forall$  compact  $K$ , then we say that an  $n$ -dimensional subspace of  $P$  of  $\mathbb{R}^{n+k}$  is the approximate tangent space for  $M$  at  $x$  ( $x$  a given point in  $\mathbb{R}^{n+k}$ ) if

$$\lim_{\lambda \downarrow 0} \int_{\eta_{x,\lambda}(M)} f(y) d\mathcal{H}^n(y) = \int_P f(y) d\mathcal{H}^n(y) \quad \forall f \in C_c^0(\mathbb{R}^{n+k}).$$

(Recall  $\eta_{x,\lambda} : \mathbb{R}^{n+k} \rightarrow \mathbb{R}^{n+k}$  is defined by  $\eta_{x,\lambda}(y) = \lambda^{-1}(y - x)$ ,  $x, y \in \mathbb{R}^{n+k}$ ,  $\lambda > 0$ .)

**1.5 Remarks:** (1) Of course  $P$  is unique if it exists; we shall denote it by  $T_x M$ .

(2) We show below (in the proof of the “ $\Rightarrow$ ” part of 1.6) that, with  $M_j, N_j$  as in Remark 1.3 above,

$$T_x M = T_x N_j, \quad \mathcal{H}^n\text{-a.e. } x \in M_j, \quad j = 1, 2, \dots$$

This is a very useful fact.

(3) By choosing  $f : \mathbb{R}^{n+k} \rightarrow [0, 1] \in C_c^0(\mathbb{R}^{n+k})$  with  $f \equiv 1$  on  $B_1(0)$  and  $f \equiv 0$  on  $\mathbb{R}^{n+k} \setminus B_{1+\varepsilon}(0)$  in Definition 1.4, we see (after letting  $\varepsilon \downarrow 0$ ) that  $T_x M$  exists  $\Rightarrow$

$$\lim_{\rho \downarrow 0} (\omega_n \rho^n)^{-1} \mathcal{H}^n(M \cap B_\rho(x)) = 1.$$

and similarly, if  $T_x M$  exists, if  $0 < \alpha < 1$ , and if we let  $f : \mathbb{R}^{n+k} \rightarrow [0, 1] \in C_c^0(\mathbb{R}^{n+k})$  in Definition 1.4 approximate the indicator function of  $\{y \in \mathbb{R}^{n+k} : \text{dist}(y, (T_x M)^\perp) \leq \alpha|y|\} \cap B_1(0)$  then

$$\lim_{\rho \downarrow 0} (\omega_n \rho^n)^{-1} \mathcal{H}^n(M \cap \{y \in \mathbb{R}^{n+k} : \text{dist}(y - x, (T_x M)^\perp) \leq \alpha|y - x|\} \cap B_\rho(x)) = 0.$$

The following theorem gives the important characterization of countably  $n$ -rectifiable sets in terms of existence of approximate tangent spaces.

**1.6 Theorem.** *Suppose  $M$  is  $\mathcal{H}^n$ -measurable with  $\mathcal{H}^n(M \cap K) < \infty$  for each compact  $K \subset \mathbb{R}^{n+k}$ . Then  $M$  countably  $n$ -rectifiable  $\iff$  the approximate tangent space  $T_x M$  exists for  $\mathcal{H}^n$ -a.e.  $x \in M$ .*

**Proof of 1.6 “ $\Rightarrow$ ”:** As described in Remark 1.3 above, we may write  $M$  as the disjoint union  $\cup_{j=0}^\infty M_j$ , where  $\mathcal{H}^n(M_0) = 0$ ,  $M_j \subset N_j$ ,  $j \geq 1$ ,  $N_j$  embedded  $C^1$  submanifolds of dimensions  $n$ , and  $M_j$   $\mathcal{H}^n$ -measurable. Let  $R > 0$  and  $f \in C_c^0(\mathbb{R}^{n+k})$  with  $f \equiv 0$  in  $\mathbb{R}^{n+k} \setminus B_R(0)$ . For each  $j = 1, 2, \dots$  we can write

$$M = (M \setminus N_j) \cup (M \cap N_j) = (M \setminus N_j) \cup (N_j \setminus (N_j \setminus M)),$$

and hence

$$\int_{\eta_{x,\lambda}(M)} f d\mathcal{H}^n = \int_{\eta_{x,\lambda}(N_j)} f d\mathcal{H}^n - \int_{\eta_{x,\lambda}(N_j \setminus M)} f d\mathcal{H}^n + \int_{\eta_{x,\lambda}(M \setminus N_j)} f d\mathcal{H}^n$$

If  $x \in M_j$ , then  $x \in N_j$  and  $N_j$  is a  $C^1$  embedded submanifold, so

$$\lim_{\lambda \downarrow 0} \int_{\eta_{x,\lambda}(N_j)} f d\mathcal{H}^n = \int_{T_x N_j} f d\mathcal{H}^n.$$

Also, by the Upper Density Theorem 3.8 of Ch.1,

$$\begin{aligned} \left| \int_{\eta_{x,\lambda}(M \setminus N_j)} f d\mathcal{H}^n \right| &\leq \sup |f| \mathcal{H}^n(B_R(0) \cap \eta_{x,\lambda}(M \setminus N_j)) \\ &= \sup |f| \lambda^{-n} \mathcal{H}^n(B_{\lambda R}(x) \cap M \setminus N_j) \rightarrow 0 \text{ for } \mathcal{H}^n\text{-a.e. } x \in M_j. \end{aligned}$$

Similarly, again by the Upper Density Theorem,

$$\left| \int_{\eta_{x,\lambda}(N_j \setminus M)} f d\mathcal{H}^n \right| \rightarrow 0 \text{ for } \mathcal{H}^n\text{-a.e. } x \in M_j.$$

Thus we have shown that  $T_x M$  exists and  $= T_x N_j$  for  $\mathcal{H}^n$ -a.e.  $x \in M_j$ . In particular Remark 1.5(2) is checked.  $\square$

**Proof of 1.6 “ $\Leftarrow$ ”:** We can of course assume  $\mathcal{H}^n(M) > 0$ . Define  $\mu = \mathcal{H}^n \llcorner (M \cap B_R(0))$ , with any  $R > 0$  such that  $M \cap B_R(0)$  has positive measure. Then  $\mu$  is Borel regular with  $0 < \mu(\mathbb{R}^{n+k}) < \infty$ .

Given any  $k$ -dimensional subspace  $\pi \subset \mathbb{R}^{n+k}$  and any  $\alpha \in (0, 1)$  we let  $X_\alpha(\pi, x)$  denote the double cone

$$(1) \quad X_\alpha(\pi, x) = \{y \in \mathbb{R}^{n+k} : \text{dist}(y - x, \pi) \leq \alpha|y - x|\},$$

which can alternatively be written

$$(2) \quad X_\alpha(\pi, x) = \{y \in \mathbb{R}^{n+k} : |q_\pi(y - x)| \leq \alpha|y - x|\},$$

where  $q_\pi$  denotes orthogonal projection of  $\mathbb{R}^{n+k}$  onto  $\pi^\perp$ , with

$$\pi^\perp = \{z \in \mathbb{R}^{n+k} : z \cdot w = 0 \forall w \in \pi\}.$$

For  $k$ -dimensional subspaces  $\pi, \pi'$  we define the distance between  $\pi, \pi'$ , denoted  $d(\pi, \pi')$ , by

$$(3) \quad d(\pi, \pi') = \sup_{|x|=1} |p_\pi(x) - p_{\pi'}(x)|,$$

where  $p_\pi$  denotes orthogonal projection of  $\mathbb{R}^{n+k}$  to  $\pi$ , so that in fact  $d(\pi, \pi')$  is just the norm  $\|p_\pi - p_{\pi'}\|$  of the linear map  $p_\pi - p_{\pi'}$ .

By Remark 1.5(3) we have

$$(4) \quad \lim_{\rho \downarrow 0} \frac{\mu(B_\rho(x))}{\omega_n \rho^n} = 1$$

and

$$(5) \quad \lim_{\rho \downarrow 0} \frac{\mu(X_{\frac{1}{2}}(\pi_x, x) \cap B_\rho(x))}{\omega_n \rho^n} = 0,$$

for  $\mu$ -a.e.  $x \in M \cap B_R(0)$ , where  $\pi_x = (P_x)^\perp$ .

For  $k = 1, 2, \dots$  and  $\mathcal{H}^n$ -a.e.  $x \in M \cap B_R(0)$ , define

$$f_k(x) = \inf_{0 < \rho < \frac{1}{k}} \frac{\mu(B_\rho(x))}{\omega_n \rho^n}$$

and

$$q_k(x) = \sup_{0 < \rho < \frac{1}{k}} \frac{\mu(X_{\frac{1}{2}}(\pi_x, x) \cap B_\rho(x))}{\omega_n \rho^n}.$$

Then

$$(6) \quad \lim f_k(x) = 1 \text{ and } \lim q_k(x) = 0 \text{ } \mu\text{-a.e. } x \in M \cap B_R(0),$$

and hence by Egoroff's Theorem (1.12 of Ch. 1) we can choose a Borel set  $E \subset M \cap B_R(0)$  with

$$(7) \quad \mu(\mathbb{R}^{n+k} \setminus E) \leq \frac{1}{2} \mu(\mathbb{R}^{n+k})$$

and with (6) holding *uniformly* for  $x \in E$ . Thus for each  $\varepsilon > 0$  there is a  $\delta > 0$  such that

$$(8) \quad \frac{\mu(B_\rho(x))}{\omega_n \rho^n} \geq 1 - \varepsilon, \quad \frac{\mu(X_{\frac{1}{2}}(\pi_x, x) \cap B_\rho(x))}{\omega_n \rho^n} \leq \varepsilon$$

$x \in E, 0 < \rho \leq \delta$ .

Now choose  $k$ -dimensional subspaces  $\pi_1, \dots, \pi_N$  of  $\mathbb{R}^{n+k}$  ( $N = N(n, k)$ ) such that for *each*  $k$ -dimensional subspace  $\pi$  of  $\mathbb{R}^{n+k}$ , there is a  $j \in \{1, \dots, N\}$  such that  $d(\pi, \pi_j) < \frac{1}{16}$ , and let  $E_1, \dots, E_N$  be the subsets of  $E$  defined by

$$E_j = \{x \in E : d(\pi_j, \pi_x) < \frac{1}{16}\}.$$

Then  $E = \cup_{j=1}^N E_j$  and we claim that if we take  $\delta > 0$  such that (8) holds with  $\varepsilon = 1/16^{n+1}$ , then

$$(9) \quad X_{\frac{1}{4}}(\pi_j, x) \cap E_j \cap B_{\delta/2}(x) = \{x\}, \quad \forall x \in E_j, \quad j = 1, \dots, N.$$

Indeed otherwise we could find a point  $x \in E_j$  and a  $y \in X_{\frac{1}{4}}(\pi_j, x) \cap E_j \cap \partial B_\rho(x)$  for some  $0 < \rho \leq \delta/2$ . But since  $x \in E$  and  $2\rho \leq \delta$ , we have (by (8) with  $\varepsilon = 1/16^{n+1}$ )

$$(10) \quad \mu(X_{\frac{1}{2}}(\pi_x, x) \cap B_{2\rho}(x)) < 16^{-n-1} \omega_n (2\rho)^n < \frac{1}{2} (\rho/8)^n.$$

On the other hand  $B_{\rho/8}(y) \subset X_{\frac{1}{2}}(\pi_j, x) \cap B_{2\rho}(x)$ , because  $|z - y| < \rho/8 \Rightarrow d(z - x, \pi_x) \leq \rho/8 + d(y - x, \pi_x) = \rho/8 + |(y - x) - p_{\pi_x}(y - x)| \leq \rho/8 + |(y - x) -$

$p_{\pi_j}(y-x)| + |(p_{\pi_j} - p_{\pi_x})(y-x)| \leq 7\rho/16 \leq |z-x|/2$ . Hence we have also (again by (8))

$$\mu(X_{\frac{1}{2}}(\pi_x, x) \cap B_{2\rho}(x)) \geq \mu(B_{\rho/8}(y)) > \frac{1}{2}\omega_n(\rho/8)^n,$$

which contradicts (10). We have therefore proved (9).

Take any fixed  $x_0 \in E_j$ . Since  $B_{\delta/4}(x_0) \subset B_{\delta/2}(x)$  for each  $x \in B_{\delta/4}(x_0)$ , (9) implies

$$(11) \quad X_{\frac{1}{4}}(\pi_j, x) \cap (E_j \cap B_{\delta/4}(x_0)) = \{x\}, \quad \forall x \in E_j \cap B_{\delta/4}(x_0), \quad j = 1, \dots, N.$$

If  $Q$  is an orthogonal transformation of  $\mathbb{R}^{n+k}$  with  $\pi_j = Q(\{0\} \times \mathbb{R}^k)$ , (11) evidently implies that  $E_j \cap B_{\delta/4}(x_0)$  is contained in the graph of a Lipschitz function defined over a domain in  $\mathbb{R}^n$ , and hence by the extension theorem 1.2 of Ch.2, we have

$$E_j \cap B_{\delta/4}(x_0) \subset Q(\text{graph } f),$$

where  $f = (f^1, \dots, f^k) : \mathbb{R}^n \rightarrow \mathbb{R}^k$  is Lipschitz.

Since  $j \in \{1, \dots, N\}$  and  $x_0 \in E_j$  are arbitrary and since we can cover  $E$  by finitely many balls  $B_{\delta/4}(x_i)$ , where  $x_i \in E$ , we conclude that there are finitely many Lipschitz functions  $f_1, \dots, f_J : \mathbb{R}^n \rightarrow \mathbb{R}^k$  and orthogonal transformations  $Q_1, \dots, Q_J$  of  $\mathbb{R}^{n+k}$  such that

$$E \subset \cup_{j=1}^J Q_j(\text{graph } f_j).$$

Thus by (7) we have

$$\mu(\mathbb{R}^{n+k} \setminus \cup_{j=1}^J Q_j(\text{graph } f_j)) \leq \frac{1}{2}\mu(\mathbb{R}^{n+k}).$$

Since we can now repeat the argument, starting with  $M \cap B_R(0) \setminus (\cup_{j=1}^J Q_j(\text{graph } f_j))$  in place of  $M \cap B_R(0)$ , we thus deduce that there are countably many Lipschitz graphs  $\text{graph } f_j$ ,  $j = 1, 2, \dots$ ,  $f_j : \mathbb{R}^n \rightarrow \mathbb{R}^k$ , and corresponding orthogonal transformations  $Q_1, Q_2, \dots$  with  $\mu(\mathbb{R}^{n+k} \setminus \cup_{j=1}^\infty Q_j(\text{graph } f_j)) = 0$ . Taking  $G_j$  to be the graph map  $x \mapsto (x, f_j(x))$  and  $F_j = Q_j \circ G_j$  we then have  $F_j : \mathbb{R}^n \rightarrow \mathbb{R}^{n+k}$  Lipschitz and  $\mathcal{H}^n(M \cap B_R(0) \setminus (\cup_{j=1}^\infty F_j(\mathbb{R}^n))) = 0$ , so, since  $M = \cup_{j=1}^\infty M \cap B_j(0)$ , we conclude  $M$  is countably  $n$ -rectifiable.  $\square$

It is often convenient to be able to relax the condition  $\mathcal{H}^n(M \cap K) < \infty \quad \forall$  compact  $K$  in 1.4 and 1.6 and consider instead sets  $M$  which can be written as the countable union  $\cup_{j=1}^\infty M_j$  of  $\mathcal{H}^n$ -measurable sets  $M_j$  with  $\mathcal{H}^n(M_j \cap K) < \infty$  for each  $j$  and each compact  $K \subset \mathbb{R}^{n+k}$ . This is evidently equivalent to the requirement that  $M$  is  $\mathcal{H}^n$ -measurable and there exists a positive  $\mathcal{H}^n$ -measurable function  $\theta$  on  $M$  such that  $\int_{M \cap K} \theta \, d\mathcal{H}^n < \infty$  for each compact  $K$ , so we proceed to discuss this situation, starting with the definition of approximate tangent space in such a setting:

**1.7 Definition:** Let  $M$  be an  $\mathcal{H}^n$ -measurable subset of  $\mathbb{R}^{n+k}$  and let  $\theta$  be a positive  $\mathcal{H}^n$ -measurable function on  $M$  with  $\int_{M \cap K} \theta d\mathcal{H}^n < \infty$  for each compact  $K \subset \mathbb{R}^{n+k}$ . We say an  $n$ -dimensional subpace  $P_x$  is an approximate tangent space of  $M$  with respect to  $\theta$  at the point  $x \in M$  if

$$(\ddagger) \quad \lim_{\lambda \downarrow 0} \int_{\eta_{x,\lambda}(M)} f(y) \theta(x + \lambda y) d\mathcal{H}^n(y) = \theta(x) \int_{P_x} f(y) d\mathcal{H}^n(y)$$

for each  $f \in C_c^0(\mathbb{R}^{n+k})$ . Evidently  $P_x$  is unique if it exists at all so we denote it  $T_x M$ , and also  $T_x M$  agrees  $\mathcal{H}^n$ -a.e. with our previous notion of approximate tangent space in case  $\theta$  is equal to 1  $\mathcal{H}^n$ -a.e. in  $M$ .

**1.8 Remark:** By taking a  $C^0$  function  $f : \mathbb{R}^{n+k} \rightarrow [0, 1]$  with  $f \equiv 1$  in  $B_1(0)$  and  $f \equiv 0$  in  $\mathbb{R}^{n+k} \setminus B_{1+\varepsilon}(0)$ , and then letting  $\varepsilon \downarrow 0$ , we see that the definition  $(\ddagger)$  implies in particular that

$$\lim_{\rho \downarrow 0} (\omega_n \rho^n)^{-1} \int_{M \cap B_\rho(y)} \theta d\mathcal{H}^n = \theta(y)$$

whenever  $M$  has an approximate tangent space with respect to  $\theta$  at  $y$ .

We now have the following generalization of Theorem 1.6:

**1.9 Theorem.** Suppose  $M \subset \mathbb{R}^{n+k}$  is  $\mathcal{H}^n$ -measurable and  $\theta$  is a positive  $\mathcal{H}^n$ -measurable function on  $M$  with  $\int_{M \cap K} \theta d\mathcal{H}^n < \infty$  for each compact  $K \subset \mathbb{R}^{n+k}$ . Then  $M$  is countably  $n$ -rectifiable  $\iff M$  has an approximate tangent space  $T_x M$  with respect to  $\theta$  for  $\mathcal{H}^n$ -a.e.  $x \in M$ .

**Proof:** Let  $\mu = \mathcal{H}^n \llcorner \theta$ . By Lusin's Theorem 1.24 and Remark 1.25 of Ch. 1 there is an increasing sequence  $\{M_j\}_{j=1,2,\dots}$  of closed sets with  $M_j \subset M$ ,  $\mu(M \setminus (\cup_j M_j)) = 0$  and  $\theta|_{M_j}$  continuous for each  $j$ , hence of course then  $\inf_{M_j \cap K} \theta > 0$  for each compact  $K \subset \mathbb{R}^{n+k}$  and in particular  $\mathcal{H}^n(M_j \cap K) < \infty$  for each compact  $K \subset \mathbb{R}^{n+k}$ ,  $j = 1, 2, \dots$

Using the continuity of  $\theta|_{M_j}$ ,

- (1)  $\forall x \in M_j : T_x M_j$  exists (as in Definition 1.4)  $\iff T_x M_j$  is the approximate tangent space of  $M_j$  with respect to  $\theta$  (as in Definition 1.7).

Also the Upper Density Theorem 3.8 of Ch. 1 implies that  $\lim_{\rho \downarrow 0} \rho^{-n} \mu((M \setminus M_j) \cap B_\rho(x)) = 0$  for  $\mathcal{H}^n$ -a.e.  $x \in M_j$  and hence, for  $\mathcal{H}^n$ -a.e.  $x \in M_j$ ,

- (2)  $T_x M_j$  is the approximate tangent space of  $M_j$  with respect to  $\theta \iff T_x M_j$  is the approximate tangent space of  $M$  with respect to  $\theta$ .

But, according to Theorem 1.6,  $M_j$  countable rectifiable  $\iff T_x M_j$  exists for  $\mathcal{H}^n$ -a.e.  $x \in M_j$ , and of course  $M$  countably rectifiable  $\iff M_j$  is countably rectifiable for each

$j$ . So by (1) and (2) we have  $M$  countably  $n$ -rectifiable  $\iff M$  has an approximate tangent space with respect to  $\theta$  at  $x$  (as in Definition 1.7) for  $\mathcal{H}^n$ -a.e.  $x \in M$ .  $\square$

## 2 Gradients, Jacobians, Area, Co-Area

Throughout this section  $M$  is supposed to be  $\mathcal{H}^n$ -measurable with locally finite  $\mathcal{H}^n$  measure, and countably  $n$ -rectifiable, so that we can express  $M$  as the disjoint union  $\cup_{j=0}^{\infty} M_j$  (as in Remark 1.3 of the present chapter), where  $\mathcal{H}^n(M_0) = 0$ ,  $M_j$  is  $\mathcal{H}^n$ -measurable of finite  $\mathcal{H}^n$ -measure, and  $M_j \subset N_j$ ,  $j \geq 1$ , where  $N_j$  are  $n$ -dimensional  $C^1$  embedded submanifolds of  $\mathbb{R}^{n+k}$ .

Let  $f$  be a locally Lipschitz function on  $U$ , where  $U$  is an open set in  $\mathbb{R}^{n+k}$  containing  $M$ . Then according to the discussion in §4 of Ch.2 we can define the gradient of  $f$ ,  $\nabla^M f$ ,  $\mathcal{H}^n$ -a.e.  $y \in M$  by

### 2.1 Definition:

$$\nabla^M f(y) = \nabla^{N_j} f(y), \quad y \in M_j,$$

where the notation is as above.

Notice that, up to change on sets of  $\mathcal{H}^n$ -measure zero, this is independent of the decomposition  $M = \cup M_j$  (and independent of the choice of the  $C^1$  submanifolds  $N_j$ ). Because for  $\mathcal{H}^n$ -a.e.  $y \in M$  we have  $D_{\tau} f(y) = \frac{d}{dt} f(y + t\tau)|_{t=0}$  for all  $\tau \in T_y M$ , and is a linear function of  $\tau \in T_y M$ , by 4.16 of Ch.2 and the fact that  $T_y M$  agrees with  $T_y N_j$  for  $\mathcal{H}^n$ -a.e.  $y \in M_j (= M \cap N_j)$ . In particular  $\nabla^M f(x) = \sum_{i=1}^n D_{\tau_i} f(x) \tau_i$ , where  $\tau_1, \dots, \tau_n$  is an orthonormal basis for  $T_y M$ , is well defined as an  $L^1$  function with respect to Hausdorff measure  $\mathcal{H}^n$  on  $M$ .

Having defined  $\nabla^M f$ , we can now define the linear  $d^M f_x : T_x M \rightarrow \mathbb{R}$  induced by  $f$  by setting

$$d^M f_x(\tau) = D_{\tau} f(y) (= \langle \tau, \nabla^M f(x) \rangle), \quad \tau \in T_x M$$

at all points where  $T_x M$  and  $\nabla^M f(x)$  exist.

More generally, if  $f = (f^1, \dots, f^Q)$  takes values in  $\mathbb{R}^Q$  ( $f^j$  still locally Lipschitz on  $U$ ,  $j = 1, \dots, Q$ ), we define  $d^M f_x : T_x M \rightarrow \mathbb{R}^Q$  by

$$2.2 \quad d^M f_x(\tau) = D_{\tau} f(x).$$

With such an  $f$ , in case  $Q = n + k$  ( $k \geq 0$ ), we define the Jacobian  $J_f^M(x)$  for  $\mathcal{H}^n$ -a.e.  $x \in M$  as in 4.18 of Ch.2; thus

$$2.3 \quad J_f^M(x) = \sqrt{\det((d^M f_x)^* \circ (d^M f_x))} = \sqrt{\det(D_{\tau_i} f(x) \cdot D_{\tau_j} f(x))},$$

where  $\tau_1, \dots, \tau_n$  is any orthonormal basis for  $T_x M$  and  $(d^M f_x)^* : \mathbb{R}^{n+k_1} \rightarrow T_x M$  denotes the adjoint of  $d^M f_x$ .

We then have the general area formula (still assuming  $Q \geq n$ )

$$2.4 \quad \int_A J_f^M d\mathcal{H}^n = \int_{\mathbb{R}^{n+k}} \mathcal{H}^0(A \cap f^{-1}(y)) d\mathcal{H}^n(y)$$

for any  $\mathcal{H}^n$ -measurable set  $A \subset M$ . Indeed by 4.19 of Ch.2 we do have 2.4 with  $N_j$  in place of  $M$  and  $A \cap M_j \subset N_j$  in place of  $A$  and  $j \geq 1$ , because, for  $j \geq 1$ ,  $J_f^{N_j} = J_f^M$   $\mathcal{H}^n$ -a.e. on  $M_j$ . We then conclude 2.4 by summing over  $j \geq 1$  and using the (easily checked) fact that if  $\psi : U \rightarrow \mathbb{R}^m$  is locally Lipschitz and  $B$  has  $\mathcal{H}^n$ -measure zero, then  $\mathcal{H}^n(\psi(B)) = 0$ .

We note also that if  $h$  is any non-negative  $\mathcal{H}^n$ -measurable function on  $M$ , then, by approximation of  $h$  by simple functions, 2.4 implies the more general formula

$$2.5 \quad \int_M h J_f^M d\mathcal{H}^n = \int_{\mathbb{R}^{n+k}} \left( \int_{f^{-1}(y)} h d\mathcal{H}^0 \right) d\mathcal{H}^n(y).$$

In case  $f|_M$  is 1:1 this becomes

$$2.6 \quad \int_M h J_f^M d\mathcal{H}^n = \int_{f(M)} h \circ f^{-1} d\mathcal{H}^n.$$

There is also a version of the co-area formula in case  $M$  is merely  $\mathcal{H}^n$ -measurable, countably  $n$ -rectifiable and  $f : U \rightarrow \mathbb{R}^m$  ( $U$  open  $U \supset M$ ) is locally Lipschitz with  $m < n$ , so that  $n = m + k$  with  $k \in \{0, 1, \dots\}$ .

In fact we can define (Cf. the smooth case described in §7 of Ch.2)

$$2.7 \quad J_f^M(x) = \sqrt{\det(d^M f_x \circ (d^M f_x)^*)} = \sqrt{\det(\nabla^M f^i(x) \cdot \nabla^M f^j(x))},$$

with  $d^M f_x$  as in 2.2 and  $(d^M f_x)^* = \text{adjoint of } d^M f_x$ . Then, for any  $\mathcal{H}^n$ -measurable set  $A \subset M$ ,

$$2.8 \quad \int_A J_f^M d\mathcal{H}^n = \int_{\mathbb{R}^m} \mathcal{H}^k(A \cap f^{-1}(y)) d\mathcal{L}^m(y).$$

This follows from the  $C^1$  case (see §7 of Ch.2) by using the decomposition  $M = \cup_{j=0}^{\infty} M_j$  of Remark 1.3 and the  $C^1$  Approximation Theorem 1.5 of Ch.2 in a similar manner to the procedure used for the discussion of the area formula above.

As for the smooth case, approximating a given non-negative  $\mathcal{H}^n$ -measurable function  $g$  by simple functions, we deduce directly from 2.8 the more general formula

$$2.9 \quad \int_A g J_f^M d\mathcal{H}^n = \int_{\mathbb{R}^m} \left( \int_{f^{-1}(y) \cap M} g d\mathcal{H}^k \right) d\mathcal{L}^m(y).$$

**2.10 Remarks:** (1) Note that Remark 7.7 of Ch.2 carries over without change to this setting.

(2) The “slices”  $M \cap f^{-1}(y)$  are countably  $k$ -rectifiable subsets of  $\mathbb{R}^{n+k}$  for  $\mathcal{L}^m$ -a.e.  $y \in \mathbb{R}^m$ . This follows directly from the decomposition  $M = \bigcup_{j=0}^{\infty} M_j$  of Remark 1.3 together with the  $C^1$  Sard-type Theorem 7.5 of Ch.2 and the  $C^1$  Approximation Theorem 1.5 of Ch.2.

### 3 Purely Unrectifiable Sets, Structure Theorem

**3.1 Definition:** A subset  $S \subset \mathbb{R}^{n+k}$  is said to be purely  $n$ -unrectifiable if  $P$  contains no countably  $n$ -rectifiable subsets of positive  $\mathcal{H}^n$ -measure.

**3.2 Lemma.** *If  $A$  is an arbitrary  $\mathcal{H}^n$   $\sigma$ -finite subset of  $\mathbb{R}^{n+k}$  (i.e.  $A = \bigcup_{j=1}^{\infty} A_j$  with  $\mathcal{H}^n(A_j) < \infty$  for each  $j$ ), it is always possible to decompose  $A$  into a disjoint union*

$$A = R \cup P,$$

*where  $R$  is countably  $n$ -rectifiable and  $P$  is purely  $n$ -unrectifiable. Also  $R$  can be chosen to be a Borel set if  $A$  is  $\mathcal{H}^n$  measurable.*

**Proof:** First observe that in case  $A$  is  $\mathcal{H}^n$ -measurable we can also take each  $A_j$  to be  $\mathcal{H}^n$ -measurable (e.g. first take a Borel set  $B_j \supset A_j$  with  $\mathcal{H}^n(B_j) = \mathcal{H}^n(A_j)$  and then replace  $A_j$  by  $A \cap B_j$ ), then by Theorem 1.22(2) of Ch.1 we can take a Borel set  $C_j \subset A_j$  with  $\mu(A_j \setminus C_j) = 0$ . In this case we let

$$\alpha_j = \sup\{\mathcal{H}^n(S) : S \subset C_j, S \text{ a countably } n\text{-rectifiable Borel set}\}.$$

By definition of  $\alpha_j$  we can select countably  $n$ -rectifiable Borel sets  $R_{ij} \subset C_j$  with  $\mathcal{H}^n(R_{ij}) > \alpha_j - 1/i$  and let  $R_j = \bigcup_i R_{ij}$ . Evidently  $R_j$  is a countably  $n$ -rectifiable Borel set and  $C_j \setminus R_j$  is purely unrectifiable, because if  $C_j \setminus R_j$  contains a countably  $n$ -rectifiable set of positive measure then  $C_j \setminus R_j$  contains a countably  $n$ -rectifiable Borel set  $B_j$  of positive measure and hence  $R \cup B_j \subset C_j$  with  $\mathcal{H}^n$ -measure  $> \alpha_j$ , contradicting the definition of  $\alpha_j$ . So 3.2 is proved with  $R = \bigcup_j R_j$  Borel and  $P = A \setminus R$   $\mathcal{H}^n$ -measurable.

To handle the case when  $A$  is not necessarily  $\mathcal{H}^n$ -measurable, we first pick a Borel set  $B = \bigcup_j B_j$ , where each  $B_j$  is a Borel set containing  $A_j$  with the same  $\mathcal{H}^n$ -measure as  $A_j$ . Then by the case of the theorem when  $A$  is  $\mathcal{H}^n$ -measurable which we established above, we have  $B = R \cup P$  (disjoint union) with  $R$  countably  $n$ -rectifiable Borel and  $P$  purely  $n$ -unrectifiable, and then  $A = (A \cap R) \cup (A \cap P)$  is a suitable decomposition of  $A$ .  $\square$

The following lemma gives a simple and convenient sufficient condition for checking if a set is purely  $n$ -unrectifiable. In this lemma we adopt the notation that  $p_L$  denotes the orthogonal projection of  $\mathbb{R}^{n+k}$  onto  $L$  for any  $n$ -dimensional subspace  $L \subset \mathbb{R}^{n+k}$ .

**3.3 Lemma.** *For  $1 \leq j_1 < j_2 < \cdots < j_n \leq n+k$  let  $p_{j_1, \dots, j_n}$  denote the orthogonal projection of  $\mathbb{R}^{n+k}$  onto  $\text{span}\{e_{j_1}, \dots, e_{j_n}\}$ , and suppose  $S \subset \mathbb{R}^{n+k}$  has the property that  $\mathcal{H}^n(p_{j_1, \dots, j_n}(S)) = 0$  for each  $1 \leq j_1 < \cdots < j_n \leq n+k$ . Then  $S$  is purely  $n$ -unrectifiable.*

**Proof of 3.3:** Suppose on the contrary that  $S$  is not purely  $n$ -unrectifiable. Then Lemma 1.2 implies there is an  $n$ -dimensional  $C^1$  embedded submanifold  $N$  with  $\mathcal{H}^n(S \cap N) > 0$ , so there must be some  $x \in S \cap N$  with  $\mathcal{H}^n(S \cap N \cap B_\rho(x)) > 0$  for all  $\rho > 0$ . With such an  $x$  we see that, by Remark 4.2(2) of Ch. 2 (with  $M = N$ ) that there is  $1 \leq j_1 < j_2 < \cdots < j_n \leq n+k$  and  $\rho > 0$  such that  $p_{j_1, \dots, j_n}|_{N \cap \check{B}_\rho(x)}$  is a  $C^1$  diffeomorphism onto an open  $W \subset \text{span}\{e_{j_1}, \dots, e_{j_n}\}$  and so  $\mathcal{H}^n(p_{j_1, \dots, j_n}(S \cap N \cap \check{B}_\rho(x))) > 0$ .  $\square$

**3.4 Example.** A simple example (in the case  $n = k = 1$ ) of the use of Lemma 3.3 is the following: Let  $C_0 = [0, 1] \times [0, 1]$ ,  $C_1$  = the union of the 4 sub-squares of  $C_0$  with edge length  $\frac{1}{4}$  each sharing one corner with  $C_0$ . Observe that the orthogonal projection  $p$  onto the line  $y = \frac{1}{2}x$  projects  $C_1$  onto a full line segment  $\sigma$  of length  $\frac{3}{\sqrt{5}}$ . Thus if we inductively define a sequence  $C_n$  of sets, each of which is the union of  $4^n$  squares with edge length  $4^{-n}$  and if we stipulate that  $C_{n+1}$  is obtained from  $C_n$  by replacing each square  $s$  of  $C_n$  with 4 squares of edge-length  $4^{-n-1}$ , each sharing a corner with  $s$ , then  $C_{n+1} \subset C_n$  and each  $C_n$  projects via the orthogonal projection  $p$  onto the full line segment  $\sigma$ , and hence so does the compact set  $C = \bigcap_{n=0}^{\infty} C_n$ . Furthermore  $\mathcal{H}^1(C) \geq \mathcal{H}^1(p(C)) = \frac{3}{\sqrt{5}} > 0$ , and also  $\mathcal{H}^1(C) \leq \sqrt{2} < \infty$  because each of the  $4^n$  squares comprising  $C_n$  has diameter  $4^{-n}\sqrt{2}$ . Finally, each  $C_n$  projects via orthogonal projection  $p_x$  of  $\mathbb{R}^2$  onto the  $x$ -axis to a union of  $2^n$  closed intervals each of length  $4^{-n}$ , and hence  $\mathcal{L}^1(p_x(C)) = \lim \mathcal{L}^1(p_x(C_n)) = 0$ . Similarly  $\mathcal{L}^1(p_y(C)) = 0$ , where  $p_y$  denotes orthogonal projection onto the  $y$ -axis. Evidently then Lemma 3.3 is applicable with  $n = k = 1$ , so  $C$  is purely 1-unrectifiable.

A very non-trivial theorem (the Structure Theorem) due to Besicovitch [Bes28, Bes38, Bes39] in case  $n = k = 1$  and Federer [Fed69] in general, says that the purely unrectifiable sets  $Q$  of  $\mathbb{R}^{n+k}$  which (like the subset  $P$  in 3.2) can be written as the countable union of sets of finite  $\mathcal{H}^n$ -measure, are characterized by the fact that they have  $\mathcal{H}^n$ -null projection via almost all orthogonal projections onto  $n$ -dimensional subspaces of  $\mathbb{R}^{n+k}$ . More precisely:

**3.5 Theorem.** *Suppose  $Q$  is a purely  $n$ -unrectifiable subset of  $\mathbb{R}^{n+k}$  with  $Q = \bigcup_{j=1}^{\infty} Q_j$ ,  $\mathcal{H}^n(Q_j) < \infty \forall j$ . Then  $\mathcal{H}^n(p(Q)) = 0$  for  $\sigma$ -almost all  $p \in O(n+k, n)$ . Here  $\sigma$  is Haar*

measure for  $O(n+k, n)$ , the orthogonal projections of  $\mathbb{R}^{n+k}$  onto  $n$ -dimensional subspaces of  $\mathbb{R}^{n+k}$ .

For the proof of this theorem see [Fed69] or [Ros84].

**3.6 Remark:** Of course only the purely  $n$ -unrectifiable subsets could possibly have the null projection property described in 3.5, by virtue of Lemma 3.3 above.

Notice that, by combining 3.2 and 3.6, we get the following *Rectifiability Theorem*, which is of fundamental importance in understanding the structure of subsets of  $\mathbb{R}^{n+k}$ :

**3.7 Theorem (Rectifiability Theorem for sets.)** *If  $A$  is an arbitrary subset of  $\mathbb{R}^{n+k}$  which can be written as a countable union  $\cup_{j=1}^{\infty} A_j$  with  $\mathcal{H}^n(A_j) < \infty \forall j$ , and if every subset  $B \subset A$  with positive  $\mathcal{H}^n$ -measure has the property that  $\mathcal{H}^n(p(B)) > 0$  for a set of  $p \in O(n+k, n)$  with  $\sigma$ -measure  $> 0$ , then  $A$  is countably  $n$ -rectifiable.*

## 4 Sets of Locally Finite Perimeter

An important class of countably  $n$ -rectifiable sets comes from the sets of locally finite perimeter in  $\mathbb{R}^{n+1}$ . (Or Cacciopoli sets—see De Giorgi [DG61], Giusti [Giu84].) First we need some definitions.

If  $U \subset \mathbb{R}^{n+1}$  is open and if  $E$  is an  $\mathcal{L}^{n+1}$ -measurable subset of  $\mathbb{R}^{n+1}$ , we say that  $E$  has locally finite perimeter in  $U$  if the indicator function  $\chi_E$  of  $E$  is in  $BV_{\text{loc}}(U)$ . (See §2 of Ch.2.)

Thus  $E$  has locally finite perimeter in  $U$  if there is a Radon measure  $\mu_E (= |D\chi_E|$  in the notation of §2 of Ch.2) on  $U$  and a Borel measurable function  $\nu = (\nu^1, \dots, \nu^{n+1})$  with  $|\nu| = 1$   $\mu_E$ -a.e. in  $U$ , such that

$$4.1 \quad \int_{E \cap U} \operatorname{div} g \, d\mathcal{L}^{n+1} = - \int_U g \cdot \nu \, d\mu_E$$

for each  $g = (g^1, \dots, g^{n+1})$  with  $g^j \in C_c^1(U)$ ,  $j = 1, \dots, n+1$ . Notice that if  $E$  is open and  $\partial E \cap U$  is an  $n$ -dimensional embedded  $C^1$  submanifold of  $\mathbb{R}^{n+1}$ , then the divergence theorem tells us that 4.1 holds with  $\mu_E = \mathcal{H}^n \llcorner (\partial E \cap U)$  and with  $\nu$  = the inward pointing unit normal to  $\partial E$ . Thus in general we interpret  $\mu_E$  as a “generalized boundary measure” and  $\nu$  as a “generalized inward unit normal”. It turns out (see 4.4 below) that in fact this interpretation is quite generally correct in a rather precise (and concrete) sense.

We now want to define the *reduced boundary*  $\partial^*E$  of a set  $E$  of finite perimeter by

$$4.2 \quad \partial^*E = \left\{ x \in U : \lim_{\rho \downarrow 0} \frac{\int_{B_\rho(x)} \nu \, d\mu_E}{\mu_E(B_\rho(x))} \text{ exists and has length 1} \right\}.$$

We henceforth use the notation

$$4.3 \quad v_E(x) = \lim_{\rho \downarrow 0} \frac{\int_{B_\rho(x)} v \, d\mu_E}{\mu_E(B_\rho(x))}, \quad x \in \partial^* E.$$

By virtue of the Lebesgue Theorem 4.10 of Ch.1 we have  $v_E = v \, \mu_E$ -a.e. in  $U$ , hence  $\mu_E(U \setminus \partial^* E) = 0$  and  $\mu_E = \mu_E \llcorner \partial^* E$ . We in fact claim much more:

**4.4 Theorem (De Giorgi).** *Suppose  $E$  has locally finite perimeter in  $U$ . Then*

$$\mu_E = \mathcal{H}^n \llcorner \partial^* E,$$

*$\partial^* E$  is countably  $n$ -rectifiable, and at each point  $x \in \partial^* E$  the approximate tangent space  $T_x \partial^* E$  of  $\partial^* E$  exists (in accordance with Definition 1.4) and is given by*

$$(\ddagger) \quad T_x \partial^* E = \{y \in \mathbb{R}^{n+1} : y \cdot v_E(x) = 0\},$$

*where  $v_E$  is as in 4.3 (so that  $|v_E(x)| = 1$  by the definition 4.2). Furthermore  $v_E(x)$  is the “inward pointing unit normal for  $E$ ” in the sense that*

$$(\ddagger\ddagger) \quad E_{x,\lambda} \equiv \{\lambda^{-1}(y-x) : y \in E\} \rightarrow \{y \in \mathbb{R}^{n+1} : y \cdot v_E(x) > 0\}$$

*in the  $L^1_{\text{loc}}(\mathbb{R}^{n+1})$  sense for each  $x \in \partial^* E$ .*

**Proof:** Take any  $y \in \partial^* E$ . For convenience of notation we suppose that  $y = 0$  and  $v(0) = (0, \dots, 0, 1)$ . Then we have

$$(1) \quad \lim_{\rho \downarrow 0} \frac{\int_{B_\rho(0)} v_{n+1} \, d\mu_E}{\mu_E(B_\rho(0))} = 1.$$

Since  $v_{n+1} \leq |v_{n+1}| \leq 1$  we have also  $\lim_{\rho \downarrow 0} \frac{\int_{B_\rho(0)} |v_{n+1}| \, d\mu_E}{\mu_E(B_\rho(0))} = 1$  and hence

$$(2) \quad \lim_{\rho \downarrow 0} \frac{\int_{B_\rho(0)} |v_i| \, d\mu_E}{\mu_E(B_\rho(0))} = 0, \quad i = 1, \dots, n,$$

because  $|v_i| \leq \sqrt{1 - v_{n+1}^2} \leq \sqrt{2} \sqrt{1 - |v_{n+1}|}$ . Further if  $\zeta \in C_0^1(U)$  has support in  $B_\rho(0) \subset U$ , then by 4.1

$$(3) \quad \begin{aligned} \int_U v_{n+1} \zeta \, d\mu_E &= - \int_U \chi_E D_{n+1} \zeta \, d\mathcal{L}^{n+1} \\ &\leq \int_E |D\zeta| \, d\mathcal{L}^{n+1}. \end{aligned}$$

Now replace  $\zeta$  by a decreasing sequence  $\{\zeta_k\}$  converging pointwise to the characteristic function of  $B_\rho(0)$  and satisfying

$$(4) \quad \lim_{k \rightarrow \infty} \int_E |D\zeta_k| = \frac{d}{d\rho} \mathcal{L}^{n+1}(E \cap B_\rho(0)).$$

(Notice that this can be done whenever the right side exists, which is  $\mathcal{L}^1$ -a.e.  $\rho$ , because  $\mathcal{L}^1(E \cap B_\rho(0))$  is an increasing function of  $\rho$ .) Then (3) gives

$$(5) \quad \int_{B_\rho(0)} v_{n+1} d\mu_E \leq \frac{d}{d\rho} \mathcal{L}^{n+1}(E \cap B_\rho(0))$$

for  $\mathcal{L}^1$ -a.e.  $\rho \in (0, \rho_0)$ ,  $\rho_0 = \text{dist}(0, \partial U)$ . Then by (1) we have, for suitable  $\rho_1 \in (0, \rho_0)$ ,

$$(6) \quad \begin{aligned} \mu_E(B_\rho(0)) &\leq 2 \frac{d}{d\rho} \mathcal{L}^{n+1}(E \cap B_\rho(0)) = 2\mathcal{H}^n(E \cap \partial B_\rho(0)) \\ &\leq 2(n+1)\omega_{n+1}\rho^n \end{aligned}$$

for  $\mathcal{L}^1$ -a.e.  $\rho \in (0, \rho_1)$ .

Then by the Compactness Theorem 2.6 of Ch. 2, it follows that we can select a sequence  $\rho_k \downarrow 0$  so that  $\chi_{\eta_0, \rho_k}(E) \rightarrow \chi_F$  in  $L^1_{\text{loc}}(\mathbb{R}^{n+1})$ , where  $F$  is a set of locally finite perimeter in  $\mathbb{R}^{n+1}$ . Hence in particular for any non-negative  $\zeta \in C_0^1(\mathbb{R}^{n+1})$

$$(7) \quad \lim_{k \rightarrow \infty} \int_{\eta_0, \rho_k}(E) D_i \zeta d\mathcal{L}^{n+1} = \int_F D_i \zeta d\mathcal{L}^{n+1}.$$

Now write  $\zeta_k(x) = \zeta(\rho_k^{-1}x)$  and change variable  $x \rightarrow \rho_k x$ ; then

$$(8) \quad \int_{\eta_0, \rho_k}(E) D_i \zeta d\mathcal{L}^{n+1} = \rho_k^{-n} \int_E D_i \zeta_k d\mathcal{L}^{n+1} \equiv -\rho_k^{-n} \int_U \zeta_k v_i d\mu_E$$

(by 4.1), so that  $\int_{\eta_0, \rho_k}(E) D_i \zeta d\mathcal{L}^{n+1} \rightarrow 0$  by (2) for  $i = 1, \dots, n$ . Thus (7) gives

$$\int_F D_i \zeta d\mathcal{L}^{n+1} = 0 \quad \forall \zeta \in C_0^1(\mathbb{R}^{n+1}), \quad i = 1, \dots, n,$$

and it follows that  $F = \mathbb{R}^n \times H$  for some  $\mathcal{L}^1$ -measurable subset  $H$  of  $\mathbb{R}$ .

On the other hand by 4.1 with  $g = \zeta_k e_{n+1}$  and by (1) we have, for  $k$  sufficiently large and  $\zeta \geq 0$ ,

$$\begin{aligned} 0 \leq \rho_k^{-n} \int_U \zeta_k v_{n+1} d\mu_E &= \int_{\eta_0, \rho_k}(E) D_{n+1} \zeta \\ &\rightarrow \int_F D_{n+1} \zeta \equiv \int_{\mathbb{R}^n} \left( \int_H \frac{\partial \zeta}{\partial x^{n+1}}(x', x^{n+1}) dx^{n+1} \right) dx' \end{aligned}$$

as  $k \rightarrow \infty$ , so that  $\chi_H$  is *non-decreasing* on  $\mathbb{R}$ , hence

$$(9) \quad F = \{x \in \mathbb{R}^{n+1} : x^{n+1} < \lambda\}$$

for some  $\lambda$ . We have next to show that  $\lambda = 0$ . To check this we use the Sobolev inequality (see e.g. [GT01]) to deduce that, if  $\zeta \geq 0$ ,  $\text{spt } \zeta \subset U$  and  $\sigma < \text{dist}(\text{spt } \zeta, \partial U)$ , then

$$\begin{aligned} & \left( \int_U (\zeta \varphi_\sigma * \chi_E)^{\frac{n+1}{n}} d\mathcal{L}^{n+1} \right)^{\frac{n}{n+1}} \\ & \leq C \int_U |D(\zeta \varphi_\sigma * \chi_E)| d\mathcal{L}^{n+1} \\ & \leq C \left( \int_U \zeta |D(\varphi_\sigma * \chi_E)| d\mathcal{L}^{n+1} + \int_U \varphi_\sigma * \chi_E |D\zeta| d\mathcal{L}^{n+1} \right). \end{aligned}$$

By 2.5 of Ch.2 it follows that

$$\left( \int_E \zeta^{\frac{n+1}{n}} d\mathcal{L}^{n+1} \right)^{\frac{n}{n+1}} \leq C \left( \int_U \zeta d\mu_E + \int_E |D\zeta| d\mathcal{L}^{n+1} \right),$$

and replacing  $\zeta$  by as sequence  $\zeta_k$  as in (4), we get for a.e.  $\rho \in (0, \rho_1)$

$$(\mathcal{L}^{n+1}(E \cap B_\rho(0)))^{\frac{n}{n+1}} \leq C(\mu_E(B_\rho(0)) + \frac{d}{d\rho} \mathcal{L}^{n+1}(E \cap B_\rho(0))),$$

which by (6) gives

$$(\mathcal{L}^{n+1}(E \cap B_\rho(0)))^{\frac{n}{n+1}} \leq C \frac{d}{d\rho} \mathcal{L}^{n+1}(E \cap B_\rho(0)) \quad \text{a.e. } \rho \in (0, \rho_1).$$

$$1 \leq C \frac{d}{d\rho} \mathcal{L}^{n+1}(E \cap B_\rho(0))^{1/(n+1)} \quad \text{a.e. } \rho \in (0, \rho_1).$$

Integration (using the fact that  $\mathcal{L}^{n+1}(E \cap B_\rho(0))^{1/(n+1)}$  is an increasing function of  $\rho$  and hence  $\int_0^\rho \frac{d}{d\sigma} \mathcal{L}^{n+1}(E \cap B_\sigma(0))^{1/(n+1)} d\sigma \leq \mathcal{L}^{n+1}(E \cap B_\rho(0))^{1/(n+1)}$ ) then implies

$$(10) \quad \mathcal{L}^{n+1}(E \cap B_\rho(0)) \geq C\rho^{n+1}$$

for all sufficiently small  $\rho$ . Repeating the same argument with  $U \setminus E$  in place of  $E$ , we also deduce

$$(11) \quad \mathcal{L}^{n+1}(B_\rho(0) \setminus E) \geq C\rho^{n+1}$$

for all sufficiently small  $\rho$ . (10) and (11) evidently tell us that  $\lambda = 0$  in (9).

The argument above guarantees  $\chi_{\eta_{0,\rho}(E)} \rightarrow \chi_{\{x \in \mathbb{R}^{n+1}; x^{n+1} < 0\}}$  as  $\rho \downarrow 0$ . Then by 4.1, (1) and (3) we have

$$\mu_{\eta_{0,\rho}(E)} \rightarrow \mu_{\{x \in \mathbb{R}^{n+1}; x^{n+1} < 0\}} = \mathcal{H}^n \llcorner \{x \in \mathbb{R}^{n+1} : x^{n+1} = 0\} \text{ as } \rho \downarrow 0.$$

Of course (since we can reduce to that above case  $y = 0$  and  $\nu_E(y) = e_{n+1}$  via an orthogonal transformation) this implies in general that

$$(12) \quad \mu_{\eta_{y,\rho}(E)} \rightarrow \mu_{\{x \in \mathbb{R}^{n+1}; (x-y) \cdot \nu_E(y) < 0\}} = \mathcal{H}^n \llcorner \{x \in \mathbb{R}^{n+1} : (x-y) \cdot \nu_E(y) = 0\}$$

as  $\rho \downarrow 0$  for each  $y \in \partial^* E$ . In particular

$$(13) \quad (\omega_n \rho^n)^{-1} \mu_E(B_\rho(y)) \rightarrow 1 \quad \forall y \in \partial^* E,$$

and by the comparison theorem 4.6 (with  $\mu_E$  in place of  $\mu$ ) we have

$$\mathcal{H}^n \llcorner \partial^* E \leq \mu_E \text{ in } U,$$

so in particular  $\mathcal{H}^n \llcorner \partial^* E$  is absolutely continuous with respect to  $\mu_E$  and  $\partial^* E$  is  $\mathcal{H}^n$ -measurable with locally finite  $\mathcal{H}^n$ -measure in  $U$ . Now in view of (12) we can repeat exactly the argument of the proof of 1.6 “ $\Leftarrow$ ” with  $\mu_E$  in place of the measure  $\mu = \mathcal{H}^n \llcorner (M \cap B_R(0))$  used in that proof, in order to prove that there are Lipschitz maps  $F_j : \mathbb{R}^n \rightarrow \mathbb{R}^{n+1}$  with  $\mu(\partial^* E \setminus (\cup_j F_j(\mathbb{R}^n))) = 0$ , hence in particular  $\partial^* E$  is countably  $n$ -rectifiable.

Next let  $A \subset \partial^* E$  be arbitrary and for each  $i = 1, 2, \dots$  let  $A_i$  be the set of  $y \in A$  with  $\mu_E(B_\rho(y)) \leq 2\omega_n \rho^n$  for all  $\rho < 1/i$ . Then  $A = \cup_{i=1}^\infty A_i$  by (13), and, by definition of  $\mathcal{H}_\delta^n$  with  $\delta = 1/i$ , we can choose a family  $C_{i1}, C_{i2}, \dots$  with  $A_i \subset \cup_j C_{ij}$ ,  $\text{diam } C_{ij} < 1/i$  and  $C_{ij} \cap A_i \neq \emptyset$  for each  $j$ , and  $\sum_j \omega_n (\text{diam } C_{ij}/2)^n \leq \mathcal{H}_{1/i}^n(A_i) + 1/i$ . Then, with  $y_{ij} \in C_{ij} \cap A_i$  and  $\rho_{ij} = \text{diam } C_{ij}$ , we have

$$\mu(A_i) \leq \sum_j \mu(B_{\rho_{ij}}(y_{ij})) \leq 2^{n+1} \mathcal{H}_{1/i}^n(A_i) + 2^{n+1}/i.$$

Hence letting  $i \rightarrow \infty$  we have  $\mu \leq 2^{n+1} \mathcal{H}^n \llcorner \partial^* E$  in  $U$ , so in particular  $\mu$  is absolutely continuous with respect to  $\mathcal{H}^n \llcorner \partial^* E$ .

Since  $\partial^* E$  is countably  $n$ -rectifiable we can write it as the disjoint union  $\cup_{j=0}^\infty M_j$ , where  $\mathcal{H}^n(M_0) = 0$ ,  $M_j \subset N_j$ ,  $N_j$  being  $n$ -dimensional embedded  $C^1$  submanifolds of  $\mathbb{R}^{n+1}$  for  $j \geq 1$ . Then by (13) and the Upper Density Theorem we have  $\lim_{\rho \downarrow 0} \frac{\mu(B_\rho(x))}{\mathcal{H}^n(B_\rho(x) \cap N_j)} = 1$ ,  $\mathcal{H}^n$ -a.e.  $x \in M_j$  and hence by the Radon Nikodym Theorem 4.18 of Ch.1 we have  $\mu = \mathcal{H}^n \llcorner \partial^* E$  as required.  $\square$

## CHAPTER 3 PROBLEMS

3.1 Suppose  $a, b \in \mathbb{R}$ ,  $a < b$ , and  $\gamma : [a, b] \rightarrow \mathbb{R}^n$  is absolutely continuous. The length  $L$  of  $\gamma$  is defined as usual by  $L = \int_a^b |\gamma'(t)| dt$ . If  $\tilde{\gamma} : [0, L] \rightarrow \mathbb{R}^n$  is defined by  $\tilde{\gamma}(\tau) = \gamma(t(\tau))$ , where  $t(\tau) = \sup\{t \in [a, b] : \int_a^t |\gamma'(x)| dx \leq \tau\}$  for  $\tau \in [0, L]$ , prove (i)  $\tilde{\gamma}$  is Lipschitz with  $\text{Lip } \tilde{\gamma} \leq 1$ , (ii)  $|\tilde{\gamma}'(s)| = 1$  for a.e.  $s \in (0, L)$ , (iii)  $\tilde{\gamma}([0, L]) = \gamma([a, b])$ .

3.2 If  $C \subset \mathbb{R}^2$  is the purely 1-unrectifiable subset constructed in Example 3.5 of Ch. 3, prove that  $C \times [0, 1]$  has positive  $\mathcal{H}^2$ -measure and is purely 2-unrectifiable.

3.3 (i) If  $v_1, \dots, v_{n+\ell}$  is a basis for  $\mathbb{R}^{n+\ell}$  and if  $L$  is an  $n$ -dimensional subspace of  $\mathbb{R}^{n+\ell}$ , prove that there exist  $1 \leq j_1 < j_2 < \dots < j_n \leq n + \ell$  such that the orthogonal projection  $p_{v_{j_1}, \dots, v_{j_n}}$  of  $\mathbb{R}^{n+\ell}$  onto  $\text{span}\{v_{j_1}, \dots, v_{j_n}\}$  has the property that  $p_{v_{j_1}, \dots, v_{j_n}}|_L$  is an isomorphism of  $L$  onto  $\text{span}\{v_{j_1}, \dots, v_{j_n}\}$ .

Hint: You can of course assume without loss of generality that  $L = \mathbb{R}^n \times \{0\}$ . Observe

$$\begin{aligned} \text{rank } p_{v_{j_1}, \dots, v_{j_n}}|_{\mathbb{R}^n \times \{0\}} &= \text{rank } p_{v_{j_1}, \dots, v_{j_n}} \circ p_{\mathbb{R}^n \times \{0\}} \\ &= \text{rank } p_{\mathbb{R}^n \times \{0\}} \circ p_{v_{j_1}, \dots, v_{j_n}} \\ &= \text{rank } p_{\mathbb{R}^n \times \{0\}}|_{\text{span}\{v_{j_1}, \dots, v_{j_n}\}}. \end{aligned}$$

(ii) Using (i), check the claim made in Remark 3.4 of Ch. 3 of the text.

3.4 Justify the claim made in Remark 1.5(3) of Ch. 3 of the text, that if  $M$  is  $\mathcal{H}^n$ -measurable with  $\mathcal{H}^n(M \cap K) < \infty$  for each compact  $K$  and if  $x \in \mathbb{R}^{n+\ell}$  is such that the approximate tangent space  $T_x M$  exists, then

$$\lim_{\rho \downarrow 0} \rho^{-n} \mathcal{H}^n(M \cap X_{1/2}((T_x M)^\perp, x) \cap B_\rho(x)) = 0.$$

3.5 If  $M$  is an  $n$ -dimensional  $C^1$  submanifold of  $\mathbb{R}^{n+\ell}$ , if  $x \in M$ , and if  $T_x M$  is the tangent space of  $M$  at  $x$ , prove that  $T_x M$  is also the approximate tangent space of  $M$  at  $x$ ; i.e.

$$\lim_{\lambda \downarrow 0} \int_{\eta_{x, \lambda} M} f d\mathcal{H}^n = \int_{T_x M} f d\mathcal{H}^n$$

for every  $f \in C_c^0(\mathbb{R}^{n+\ell})$ .

Hint: Suppose without loss of generality that  $T_x M = \mathbb{R}^n \times \{0\}$  and  $x = 0$ , and use a local graphical representation for  $M$  near  $0$  as discussed in Remark 4.4 of Ch. 2 of the text.



# Chapter 4

## Theory of Rectifiable $n$ -Varifolds

§ 1	Basic Definitions and Properties .....	104
§ 2	First Variation .....	105
§ 3	Monotonicity Formulae in the Stationary Case .....	107
§ 4	Monotonicity Formulae for $L^p$ Mean Curvature .....	113
§ 5	Local Conical Approximation .....	116
§ 6	Poincaré and Sobolev Inequalities .....	118
§ 7	Miscellaneous Additional Consequences .....	123

Let  $U \subset \mathbb{R}^{n+k}$  be open and  $M \subset U$  a countably  $n$ -rectifiable,  $\mathcal{H}^n$ -measurable subset of  $\mathbb{R}^{n+k}$ , and let  $\theta$  be a positive  $\mathcal{H}^n$ -measurable function on  $M$  with  $\int_{M \cap K} \theta \, d\mathcal{H}^n < \infty$  for each compact  $K \subset U$ . Corresponding to such a pair  $(M, \theta)$  we define the rectifiable  $n$ -varifold  $\underline{v}(M, \theta)$  to be simply the equivalence class of all pairs  $(\widetilde{M}, \widetilde{\theta})$ , where  $\widetilde{M} \subset U$  is countably  $n$ -rectifiable with  $\mathcal{H}^n((M \setminus \widetilde{M}) \cup (\widetilde{M} \setminus M)) = 0$  and where  $\widetilde{\theta} = \theta$   $\mathcal{H}^n$ -a.e. on  $M \cap \widetilde{M}$ .<sup>1</sup>  $V = \underline{v}(M, \theta)$  is referred to as *a rectifiable  $n$ -varifold in  $U$* , and  $\theta$  is called *the multiplicity function of  $\underline{v}(M, \theta)$* .  $\underline{v}(M, \theta)$  is called an *integer multiplicity rectifiable varifold* if this multiplicity function is positive integer-valued  $\mathcal{H}^n$ -a.e.

In this chapter and in Ch. 5 we develop the theory of  $n$ -rectifiable varifolds in  $U$  as introduced above, particularly concentrating on *stationary* (see §2 below) rectifiable  $n$ -varifolds, which generalize the notion of classical minimal submanifolds of  $\mathbb{R}^{n+k}$  discussed in §5 of Ch. 2. Since we now consider rectifiable  $M$  (which are not necessarily smooth—indeed only have approximate tangent planes  $\mathcal{H}^n$ -a.e.), it no longer makes much sense to take a  $C^1$  vector field  $X$  with support in a compact subset of  $M$ , which was the natural approach adopted in Ch. 2 when  $M$  was a  $C^2$  submanifold. So instead in this

---

<sup>1</sup>We shall see later, in Ch. 8, that this is essentially equivalent to Allard's ([All72]) notion of  $n$ -dimensional rectifiable varifold.

chapter we work with  $C^1$  (or sometimes just Lipschitz) vector functions  $X : U \rightarrow \mathbb{R}^{n+k}$  with support of  $X$  a compact subset of  $U$ , but still making deformations of  $M$  with initial velocity given by  $X$ .

The key section is §3, in which we obtain the monotonicity formula; much of the subsequent theory is based on this and closely related formulae.

## 1 Basic Definitions and Properties

Associated to a rectifiable  $n$ -varifold  $V = \underline{v}(M, \theta)$  in the open set  $U \subset \mathbb{R}^{n+k}$  (as described in the above introduction) there is a Radon measure  $\mu$  (called the weight measure of  $V$ ) on  $U$  defined by

$$1.1 \quad \mu_V = \mathcal{H}^n \llcorner \theta,$$

where we adopt the convention that  $\theta \equiv 0$  on  $U \setminus M$ . Thus for an  $\mathcal{H}^n$ -measurable set  $A \subset U$ ,

$$\mu_V(A) = \int_{A \cap M} \theta \, d\mathcal{H}^n,$$

the mass (or weight) of the varifold  $V$ ,  $\mathbb{M}(V)$ , is defined by

$$1.2 \quad \mathbb{M}(V) = \mu_V(U) = \int_M \theta \, d\mathcal{H}^n.$$

**1.3 Definition:** We define the tangent space  $T_x V$  of  $V = \underline{v}(M, \theta)$  to be the approximate tangent space of  $M$  (as defined in the statement of 1.9 of Ch.3) whenever this exists; notice that this is independent of the choice of representative  $(M, \theta)$  for the equivalence class  $\underline{v}(M, \theta)$ .

We also define

$$1.4 \quad \text{spt } V = \text{spt } \mu_V,$$

which is the (relatively closed) set of points  $y \in U$  such that  $\mu_V(B_\rho(y)) > 0$  for each  $\rho > 0$ , or, equivalently,  $U \setminus Y$  where  $Y$  is the union of all open subsets  $W$  of  $U$  with  $\mu_V(W) = 0$ .

For any  $\mathcal{H}^n$ -measurable subset  $A \subset \mathbb{R}^{n+k}$ ,  $v \llcorner A$  is the rectifiable  $n$ -varifold in  $U$  defined by

$$1.5 \quad V \llcorner A = \underline{v}(M \cap A, \theta|_A).$$

Given a sequence  $V_k = \underline{v}(M_k, \theta_k)$  of rectifiable  $n$ -varifolds in  $U$ , we say that  $V_k \rightarrow V$  provided  $\mu_{V_k} \rightarrow \mu_V$  in the usual sense of Radon measures in  $U$ . (Notice that this is *not* varifold convergence in the sense of Ch.8.)

Next we want to discuss the notion of mapping a rectifiable  $n$ -varifold relative to a Lipschitz map. Suppose  $V = \underline{v}(M, \theta)$ ,  $M \subset U$ ,  $U$  open in  $\mathbb{R}^{n+k}$ ,  $W$  open in  $\mathbb{R}^{n+k}$  and suppose  $f : \text{spt } V \cap U \rightarrow W$  is proper<sup>2</sup>, Lipschitz and 1:1. Then we define the image varifold  $f_{\#}V$  by

$$1.6 \quad f_{\#}V = \underline{v}(f(M), \theta \circ f^{-1}).$$

Since  $K$  compact  $\Rightarrow f^{-1}(K)$  compact and  $f(M) \cap K = f(M \cap f^{-1}(K))$ , the area formula 4.20 of Ch.2 gives

$$1.7 \quad \int_{f(M) \cap K} \theta \circ f^{-1} d\mathcal{H}^n = \int_{M \cap f^{-1}(K)} J_f^M \theta d\mathcal{H}^n,$$

so in particular  $\theta \circ f^{-1}$  is locally  $\mathcal{H}^n$ -integrable in  $W$ , and therefore 1.6 does indeed define a rectifiable  $n$ -varifold in  $W$ . More generally if  $f$  satisfies the conditions above, except that  $f$  is not necessarily 1:1, then we define  $f_{\#}V$  by

$$f_{\#}V = \underline{v}(f(M), \tilde{\theta}),$$

where  $\tilde{\theta}$  is defined on  $f(M)$  by  $\sum_{x \in f^{-1}(y) \cap M} \theta(x)$  ( $= \int_{f^{-1}(y) \cap M} \theta d\mathcal{H}^0$ ). Notice that  $\tilde{\theta}$  is locally  $\mathcal{H}^n$ -integrable in  $W$  by virtue of the area formula (see §2 of Ch.3), and in fact

$$1.8 \quad \mathbb{M}(f_{\#}V) = \int_{f(M)} \tilde{\theta} d\mathcal{H}^n \equiv \int_M J_f^M \theta d\mathcal{H}^n,$$

where  $J_f^M$  is the Jacobian of  $f$  relative to  $M$  as defined in §2 of Ch.3. Thus, assuming  $m \geq n$ , we define

$$1.9 \quad J_f^M(x) = \sqrt{\det \mathcal{J}(x)}.$$

where  $\mathcal{J}(x)$  is the matrix with  $D_{\tau_\ell} f(x) \cdot D_{\tau_m} f(x)$  in the  $\ell$ -th row and  $m$ -th column ( $\tau_1, \dots, \tau_n$  any orthonormal basis for  $T_x M$ ).

## 2 First Variation

We continue to assume that  $V = \underline{v}(M, \theta)$  is a rectifiable  $n$ -varifold in  $U$ ,  $U$  open in  $\mathbb{R}^{n+k}$ , and we assume  $\varphi : (-\varepsilon, \varepsilon) \times U \rightarrow \mathbb{R}^{n+k}$  (where  $\varepsilon > 0$ ) is a  $C^1$  map such that

$$2.1 \quad \begin{cases} \varphi(0, x) = x \text{ for each } x \in U, \text{ and } \exists \text{ compact } K \subset U \text{ with} \\ \varphi(t, x) = x \text{ for all } (t, x) \in (-\varepsilon, \varepsilon) \times U, \end{cases}$$

---

<sup>2</sup>i.e.  $f^{-1}(K) \cap \text{spt } V$  is compact whenever  $K$  is a compact subset of  $W$

and such that the velocity  $\partial\varphi(t, x)/\partial t$  is also  $C^1$ . Then the initial velocity  $X = \partial\varphi(t, x)/\partial t|_{t=0}$  is a  $C^1$  vector field with compact support in  $U$ . Of course given *any*  $C^1$  vector field  $X$  with compact support in  $U$ , we can construct a function  $\varphi$  as in 2.1, with initial velocity  $X$ , simply taking  $\varphi(t, x) = x + tX(x)$  and  $\varepsilon > 0$  sufficiently small.

According to 1.8,

$$\mathbb{M}(\varphi_{t\#}(V \llcorner K)) = \int_{M \cap K} J_M \varphi_t \theta \, d\mathcal{H}^n,$$

and we can compute the *first variation*  $\left. \frac{d}{dt} \mathbb{M}(\varphi_{t\#}(V \llcorner K)) \right|_{t=0}$  exactly as in §3 of Ch.2. We thus deduce

$$2.2 \quad \left. \frac{d}{dt} \mathbb{M}(\varphi_{t\#}(V \llcorner K)) \right|_{t=0} = \int_M \operatorname{div}_M X \, d\mu_V,$$

where  $X|_x = \left. \frac{\partial}{\partial t} \varphi(t, x) \right|_{t=0}$  is the initial velocity vector for the family  $\{\varphi_t\}$  and where  $\operatorname{div}_M X$  is as in §4 of Ch.2:

$$2.3 \quad \operatorname{div}_M X = \sum_{j=1}^{n+k} \nabla_j^M X^j (= \sum_{j=1}^{n+k} e_j \cdot (\nabla^M X^j)).$$

( $\nabla^M X^j$  as in §2 of Ch.3)

We say that  $V$  is *stationary in  $U$*  if the first variation vanishes in  $U$ . That is, by 2.2, the definition is as follows:

**2.4 Definition:**  $V = \underline{v}(M, \theta)$  is stationary in  $U$  if  $\left. \frac{d}{dt} \mathbb{M}(\varphi_{t\#}(V \llcorner K)) \right|_{t=0} = 0$  for every family  $\{\varphi_t\}$  as in 2.1; of course by 2.2 this is equivalent to the requirement  $\int \operatorname{div}_M X \, d\mu_V = 0$  for any  $C^1$  vector field  $X$  on  $U$  having compact support in  $U$ .

More generally let  $N$  be an  $(n + \ell)$ -dimensional  $C^2$  embedded submanifold of  $\mathbb{R}^{n+k}$  ( $\ell \leq k$ ), let  $M \subset N$  be  $\mathcal{H}^n$ -measurable and let  $\theta > 0$  on  $M$  be such that  $\int_{M \cap K} \theta \, d\mathcal{H}^n < \infty$  for each compact  $K \subset N$ . We call such  $V = \underline{v}(M, \theta)$  a *rectifiable varifold in  $N$* .

Observe that each local representation for  $N$  provides a homeomorphism between an open subset of Euclidean space  $\mathbb{R}^{n+\ell}$  and an open subset of  $N$ , so for each  $y \in N$  there is  $\rho_y > 0$  such that  $B_{\rho_y}(y) \cap N$  is a compact subset of  $N$ . Hence the set  $U = \bigcup_{y \in N} \check{B}_{\rho_y}(y)$  is open in  $\mathbb{R}^{n+k}$  and has the property that if  $K \subset U$  is compact then  $K \cap N$  is a compact subset of  $N$ . Thus  $V = \underline{v}(M, \theta)$  can in fact also be viewed as a rectifiable varifold in the open set  $U \subset \mathbb{R}^{n+k}$  and  $\operatorname{spt} \mu_V \subset N$ .

Now let  $\varphi : (\varepsilon, \varepsilon) \times N \rightarrow N$  (where  $\varepsilon > 0$ ) be a  $C^1$  map such that

$$2.5 \quad \begin{cases} \varphi_0(x) = x \quad \forall x \in N, \quad \exists \text{ compact } K \subset N \text{ with} \\ \varphi(t, x) = x \quad \forall (t, x) \in (-\varepsilon, \varepsilon) \times (N \setminus K), \end{cases}$$

and such that the initial velocity  $X(x) = \partial\varphi(t, x)/\partial t|_{t=0}$  is  $C^1$ ; note that  $X$  will of course automatically have the property that  $X(x) \in T_x N$  for each  $x \in N$  (so  $X$  is a *tangent* vector field on  $N$ ), because, for fixed  $x \in N$ ,  $\varphi(t, x)$ ,  $t \in (-\varepsilon, \varepsilon)$ , is a  $C^1$  curve in  $N$  which passes through  $x$  at time  $t = 0$ .

The quantity  $\left. \frac{d}{dt} \mathbb{M}(\varphi_{t\#}(V \lfloor K)) \right|_{t=0}$  with  $\varphi$  as in 2.5 is called the first variation of  $V$  in  $N$ , and of course we still have the identity 2.2.

**2.6 Definition:**  $V$  is stationary in  $N$  if  $\left. \frac{d}{dt} \mathbb{M}(\varphi_{t\#}(V \lfloor K)) \right|_{t=0} = 0$  for every  $\varphi$  as in 2.5, where  $\varphi_t(x) = \varphi(t, x)$ .

As already mentioned in the discussion preceding 5.10 of Ch.2, for each  $C^1$  vector field  $X$  on  $N$  with  $X|_x \in T_x N \forall x \in N$ , there is always  $\varphi$  as in 2.5 with initial velocity  $\left. \frac{\partial}{\partial t} \varphi_t(x) \right|_{t=0} = X|_x$  for each  $x$  in  $x$ . Thus, by  $V = \underline{v}(M, \theta)$  is stationary in  $N$  that  $V = \underline{v}(M, \theta)$  is stationary in  $N$  if and only if

$$\int_M \operatorname{div}_M X \, d\mu_V = 0$$

for each  $C^1$  vector field  $X$  on  $N$  with  $\{x \in N : X(x) \neq 0\}$  contained in a compact subset of  $N$  and  $X$  tangent to  $N$  at each point of  $N$ ; that is,  $X|_x \in T_x N \forall x \in N$ . On the other hand, by exactly the computation of 5.11 of Ch.2 (which did not depend on smoothness of  $M$ ), we can start with *any*  $C^1$  vector field  $X$  on  $N$  and compute (as in 5.11 of Ch.2) that

$$\operatorname{div}_M X = \operatorname{div}_M X^T - \sum_{i=1}^n \underline{H}_M^N \cdot X$$

at all points  $x \in M$  where  $M$  has an approximate tangent space  $T_x M$ , where  $X^T$  is  $C^1$  with compact support in  $N$  and tangent to  $N$  at each point of  $N$  and, as in 5.12 of Ch.2,

$$2.7 \quad \underline{H}_M^N(x) = \sum_{i=1}^n B_x^N(\tau_i, \tau_i),$$

with  $B^N$  the second fundamental form of  $N$  and  $\tau_1, \dots, \tau_n$  any orthonormal basis for  $T_x M$ . Thus in fact we conclude that  $V$  is stationary in  $N \iff$

$$2.8 \quad \int_M \operatorname{div}_M X \, d\mu_V = - \int_M X \cdot \underline{H}_M^N \, d\mu_V \text{ for each } C^1 \text{ map } X : N \rightarrow \mathbb{R}^{n+k}$$

with  $\{x \in N : X(x) \neq 0\}$  contained in a compact subset of  $N$ .

### 3 Monotonicity Formulae in the Stationary Case

In this section we assume that  $U$  is open in  $\mathbb{R}^{n+k}$ ,  $V = \underline{v}(M, \theta)$  is stationary in  $U$ , which means Definition 2.4 holds, i.e.

$$3.1 \quad \int_M \operatorname{div}_M X \, d\mu_V = 0$$

whenever  $X$  is a  $C^1$  vector field on  $U$  with compact support in  $U$ . We proceed to extract important information from this identity by taking specific choices of the vector function  $X = (X_1, \dots, X_{n+k})$ . In fact we begin by choosing  $X_x = \gamma(r)(x - \xi)$ , where  $\xi \in U$  is fixed,  $r = |x - \xi|$ , and  $\gamma : \mathbb{R} \rightarrow [0, 1]$  is a  $C^1(\mathbb{R})$  function with

$$\gamma'(t) \leq 0 \quad \forall t, \quad \gamma(t) = 1 \text{ for } t \leq \rho, \quad \gamma(t) = 0 \text{ for } t \geq R,$$

where  $R > \rho > 0$  and  $B_R(\xi) \subset U$ .

For any  $f \in C^1(U)$  and any  $x \in M$  such that  $T_x M$  exists (see 1.6, 1.9 of Ch.3) we have (by 2.1 of Ch.3)  $\nabla^M f(x) = \sum_{j,\ell=1}^{n+k} e^{j\ell} D_\ell f(x) e_j$ , where  $D_\ell f$  denotes the partial derivative  $\frac{\partial f}{\partial x^\ell}$  of  $f$  taken in  $U$  and where  $(e^{j\ell})$  is the matrix of the orthogonal projection of  $\mathbb{R}^{n+k}$  onto  $T_x M$ , viewed as a map  $\mathbb{R}^{n+k} \rightarrow \mathbb{R}^{n+k}$ . Thus, writing  $\nabla_j^M = e_j \cdot \nabla^M$  (as in §2), with the above choice of  $X$  we deduce

$$3.2 \quad \operatorname{div}_M X = \sum_{j=1}^{n+k} \nabla_j^M X^j = \gamma(r) \sum_{j=1}^{n+k} e^{jj} + r \gamma'(r) \sum_{j,k=1}^{n+k} e^{jk} \frac{(x^j - \xi^j)}{r} \frac{(x^k - \xi^k)}{r}.$$

Since  $(e^{j\ell})$  represents orthogonal projection onto  $T_x M$  we have  $\sum_{j=1}^{n+k} e^{jj} = n$  and

$$\sum_{j,k=1}^{n+k} e^{jk} \frac{(x^j - \xi^j)}{r} \frac{(x^k - \xi^k)}{r} = \left| p_{T_x V} \left( \frac{x - \xi}{r} \right) \right|^2$$

and, writing  $\mu = \mu_V$ , the formula 3.1 thus yields

$$3.3 \quad n \int \gamma(r) d\mu + \int r \gamma'(r) |\nabla^M r|^2 d\mu = 0.$$

A useful variant of this procedure is obtained by more generally taking any non-negative  $C^1$  functions  $h : U \rightarrow \mathbb{R}$  and  $\gamma : \mathbb{R} \rightarrow \mathbb{R}$  with support of  $h(x)\gamma(r)$  a compact subset of  $U$ ; then using the computations above and keeping track of the additional terms involving derivatives of  $h$ , we see that in place of 3.3 we get

$$3.4 \quad n \int \gamma(r) h d\mu + \int r \gamma'(r) |\nabla^M r|^2 h d\mu = - \int \gamma(r) (x - \xi) \cdot \nabla^M h d\mu.$$

For the moment we work with the identity 3.3 (which is 3.4 with  $h = 1$ ). Take  $\varepsilon \in (0, 1)$  and a  $C^1$  function  $\varphi : \mathbb{R} \rightarrow [0, 1]$  such that  $\varphi(t) \equiv 1$  for  $t \leq 1$ ,  $\varphi(t) = 0$  for  $t \geq 1 + \varepsilon$  and  $\varphi'(t) \leq 0$  for all  $t$ . Then we take  $\gamma(r) = \varphi(r/\rho)$  in the above identity, provided  $(1 + \varepsilon)\rho < R$ . Since

$$r \gamma'(r) = r \rho^{-1} \varphi'(r/\rho) = -\rho \frac{\partial}{\partial \rho} [\varphi(r/\rho)],$$

this gives

$$3.5 \quad n \int_M \varphi(r/\rho) d\mu - \rho \frac{d}{d\rho} \int_M |\nabla^M r|^2 \varphi(r/\rho) d\mu = 0, \quad \rho < R/(1 + \varepsilon),$$

provided  $B_R(\xi) \subset U$  and  $(1 + \varepsilon)\rho \in (0, R]$ , which we subsequently assume.

On the other hand  $p_{T_x V} v = v - p_{(T_x V)^\perp} v$  for  $v \in \mathbb{R}^{n+k}$ , so  $|p_{T_x V}(\frac{x-\xi}{r})|^2 = 1 - |D^\perp r|^2$ , where  $D^\perp r$  denote the orthogonal projection of  $r^{-1}(x-\xi) = Dr = (D_1 r, \dots, D_{n+k} r)$  (which is a vector of length = 1) onto  $(T_x V)^\perp$ , so 3.5 can be written

$$nI(\rho) - \rho I'(\rho) = -\rho \int_M \frac{\partial}{\partial \rho} [\varphi(r/\rho)] |D^\perp r|^2 d\mu, \quad \rho < R/(1 + \varepsilon),$$

where

$$I(\rho) = \int_M \varphi(r/\rho) d\mu.$$

Thus, multiplying by  $\rho^{-n-1}$  and rearranging, we have

$$\frac{d}{d\rho} (\rho^{-n} I(\rho)) = \int_M \rho^{-n} \frac{\partial}{\partial \rho} [\varphi(r/\rho)] |D^\perp r|^2 d\mu.$$

Since  $(1 + \varepsilon)^{-n} r^{-n} \frac{\partial}{\partial \rho} [\varphi(r/\rho)] \leq \rho^{-n} \frac{\partial}{\partial \rho} [\varphi(r/\rho)] \leq r^{-n} \frac{\partial}{\partial \rho} [\varphi(r/\rho)]$ , this gives

$$3.6 \quad (1 + \varepsilon)^{-n} J'(\rho) \leq \frac{d}{d\rho} (\rho^{-n} I(\rho)) \leq J'(\rho), \quad J(\rho) = \int_M r^{-n} |D^\perp r|^2 \varphi(r/\rho) d\mu_V.$$

By integration in 3.6 over the interval  $[\sigma, \rho]$  we thus get

$$3.7 \quad \begin{aligned} \int_M (1 + \varepsilon)^{-n} r^{-n} (\varphi(r/\rho) - \varphi(r/\sigma)) |D^\perp r|^2 d\mu_V &\leq \rho^{-n} I(\rho) - \sigma^{-n} I(\sigma) \\ &\leq \int_M r^{-n} (\varphi(r/\rho) - \varphi(r/\sigma)) |D^\perp r|^2 d\mu_V. \end{aligned}$$

Now we let  $\varepsilon \downarrow 0$ . Then  $\varphi$  decreases to the indicator function of the interval  $(-\infty, 1]$  and hence  $\varphi(r/\rho)$  decreasing pointwise to the indicator function of the closed unit ball, so we obtain

$$3.8 \quad \frac{\mu_V(B_\rho(\xi))}{\omega_n \rho^n} - \frac{\mu_V(B_\sigma(\xi))}{\omega_n \sigma^n} = \omega_n^{-1} \int_{B_\rho(\xi) \setminus B_\sigma(\xi)} \frac{|D^\perp r|^2}{r^n} d\mu_V, \quad 0 < \sigma \leq \rho < R,$$

provided  $B_R(\xi) \subset U$ .

3.8 is the fundamental monotonicity identity. In particular 3.8 tells us that the ratio

$$3.9 \quad (\omega_n \rho^n)^{-1} \mu(B_\rho(\xi)) \text{ is increasing in } \rho, \quad 0 < \rho < R,$$

and hence the density

$$3.10 \quad \Theta^n(\mu_V, \xi) = \lim_{\rho \downarrow 0} \frac{\mu_V(B_\rho(\xi))}{\omega_n \rho^n} \text{ exists and is real for every } \xi \in U,$$

and by letting  $\sigma \downarrow 0$  in 3.8 we also have

$$3.11 \quad (\omega_n \rho^n)^{-1} \mu_V(B_\rho(\xi)) - \Theta^n(\mu_V, \xi) = \omega_n^{-1} \int_{B_\rho(\xi)} \frac{|D^\perp r|^2}{r^n} d\mu_V, \quad 0 < \rho < R,$$

and in particular  $\int_{B_\rho(\xi)} \frac{|D^\perp r|^2}{r^n} d\mu_V < \infty$ . We also claim the upper semi-continuity

$$3.12 \quad \Theta^n(\mu_V, \xi) \geq \limsup_{x \rightarrow \xi} \Theta^n(\mu_V, x), \quad \xi \in U.$$

To check this take and  $\rho > 0$  and  $\varepsilon \in (0, 1)$  with  $B_\rho(\xi) \subset U$  and any sequence  $\xi_j \rightarrow \xi$ . Then  $B_{(1-\varepsilon)\rho}(\xi_j) \subset B_\rho(\xi)$  for all sufficiently large  $j$ , and hence using the monotonicity 3.9 we have

$$(1 - \varepsilon)^n \Theta^n(\mu_V, \xi_j) \leq (\omega_n \rho^n)^{-1} \mu_V(B_{(1-\varepsilon)\rho}(\xi_j)) \leq (\omega_n \rho^n)^{-1} \mu_V(B_\rho(\xi))$$

for all sufficiently large  $j$ , and hence

$$(1 - \varepsilon)^n \limsup_{j \rightarrow \infty} \Theta^n(\mu_V, \xi_j) \leq (\omega_n \rho^n)^{-1} \mu_V(B_\rho(\xi)).$$

Letting  $\varepsilon \downarrow 0$  we then have  $\limsup_{j \rightarrow \infty} \Theta^n(\mu_V, \xi_j) \leq (\omega_n \rho^n)^{-1} \mu_V(B_\rho(\xi))$ , and finally, by letting  $\rho \downarrow 0$ , we obtain 3.12 as claimed.

Since  $V = \underline{v}(M, \theta)$  and  $\Theta^n(\mu_V, x) = \theta(x)$  for  $\mathcal{H}^n$ -a.e.  $x \in M$  (by Remark 1.8 of Ch. 3), 3.12 enables us to choose “canonical representatives”  $M_V, \Theta_V$  for  $V$ , so that  $V = \underline{v}(M_V, \Theta_V)$ , where

$$3.13 \quad M_V = \{x \in U : \Theta^n(\mu_V, x) > 0\} \text{ and } \Theta_V(x) = \Theta^n(\mu_V, x) \quad \forall x \in U.$$

Since  $\Theta_V$  is then upper semi-continuous in  $U$  by 3.12 we then have

$$3.14 \quad \{x \in M_V : \Theta_V(x) \geq \alpha\} \text{ is relatively closed in } U \text{ for each } \alpha > 0$$

and in particular  $M_V$  itself is relatively closed in  $U$  (and in fact equal to  $\text{spt } V \cap U$ ) in case there exists  $\alpha > 0$  with  $\theta(x) \geq \alpha$  for  $\mathcal{H}^n$ -a.e.  $x \in M$  (and then of course  $\Theta_V(x) \geq \alpha$  for every  $x \in M_V$  by 3.12).

We now want to generalize this discussion to a context which includes varifolds which are stationary in an  $(n + k)$ -dimensional  $C^2$  embedded submanifold  $N \subset \mathbb{R}^P$  (for some  $P > n + k$ ) rather than in  $\mathbb{R}^{n+k}$ , as discussed in §2 above. We in fact introduce the concept of *generalized mean curvature vector* for the varifold  $V = \underline{v}(M, \theta)$  as follows:

**3.15 Definition:** Let  $V = \underline{v}(M, \theta)$  be a rectifiable varifold in the open set  $U \subset \mathbb{R}^{n+k}$ . Then we say that  $V$  has generalized mean curvature vector  $\underline{H}$  in  $U$  if

$$(\ddagger) \quad \int_M \text{div}_M X d\mu_V = - \int_M X \cdot \underline{H} d\mu_V \quad \forall X \in C_c^1(U, \mathbb{R}^{n+k}),$$

where  $\underline{H} \in L^1_{\text{loc}}(\mu_V)$  in  $U$ . Thus  $V$  is stationary in  $U$  if and only if it has generalized mean curvature zero.

Notice also  $V$  is stationary in  $N$ , where  $N$  is a  $C^2$   $(n+k)$ -dimensional embedded submanifold of  $\mathbb{R}^{n+\ell}$ , if and only if  $V$  has generalized mean curvature  $\underline{H}_M^N$  in  $U$ , with  $\underline{H}_M^N$  as in 2.7 of the previous section.

We want to show that the above monotonicity discussion generalizes to the case when  $V = \underline{v}(M, \theta)$  has when we assume suitable bounds on generalized mean curvature  $\underline{H}$ . For this purpose, first proceed on the left side of 3.15 (§) exactly as in the case  $\underline{H} = 0$  with the same choice  $X = h\gamma(r)(x - \xi)$  ( $h$  non-negative  $C^1$ ), thus giving, in place of 3.4, the general identity

$$3.16 \quad \int (n\gamma(r) + r\gamma'(r)|\nabla^M r|^2) h d\mu_V = - \int (x - \xi) \cdot (h\underline{H} + \nabla^M h) \gamma(r) d\mu_V.$$

Replacing  $\gamma(r)$  by  $\varphi(r/\rho)$  (as in the argument leading to 3.5), then we obtain

$$3.17 \quad n \int h \varphi(r/\rho) d\mu_V - \rho \frac{d}{d\rho} \int h |\nabla^M r|^2 \varphi(r/\rho) d\mu_V = E_h(\rho),$$

with  $E_h(\rho) = \rho^{-n-1} \int (h\underline{H} + \nabla^M h) \cdot (x - \xi) \varphi(r/\rho) d\mu_V$ .

Now suppose  $B_R(\xi) \subset U$  and that there is constant  $\Lambda$  such that

$$3.18 \quad R \sup_{B_R(\xi)} |\underline{H}| \leq \Lambda \text{ on } M.$$

Then, by the identity 3.17 with  $h = 1$ , writing  $|\nabla^M r|^2 = 1 - |D^\perp r|^2$  we obtain

$$3.19 \quad (1 + \varepsilon)^{-n} J'(\rho) \leq \frac{d}{d\rho} (\rho^{-n} I(\rho)) + E_1(\rho) \leq J'(\rho),$$

where  $I, J$  are as in 3.6 and where the extra term  $E_1$  is equal to  $\rho^{-n} \int_U \rho^{-1} (x - \xi) \cdot \underline{H} \varphi(r/\rho) d\mu_V$ . Thus, since  $\varphi(r/\rho) = 0$  for  $r > (1 + \varepsilon)\rho$ ,

$$-(1 + \varepsilon) R^{-1} \Lambda \rho^{-n} I(\rho) \leq E_1(\rho) \leq (1 + \varepsilon) R^{-1} \Lambda \rho^{-n} I(\rho),$$

and hence  $E_1(\rho) = E(\rho) \rho^{-n} I(\rho)$ , with  $E(\rho) \in [-(1 + \varepsilon) R^{-1} \Lambda, (1 + \varepsilon) R^{-1} \Lambda]$  for each  $\rho \in (0, R)$ . Thus, after multiplying 3.19 by the integrating factor  $e^{F(\rho)}$ , where  $F(\rho) = \int_0^\rho E(t) dt \in [-(1 + \varepsilon) \Lambda \rho / R, (1 + \varepsilon) \Lambda \rho / R]$ , we obtain (analogously to 3.6)

$$(1 + \varepsilon)^{-n} e^{-(1+\varepsilon)\Lambda} J'(\rho) \leq \frac{d}{d\rho} (e^{F(\rho)} \rho^{-n} I(\rho)) \leq e^{(1+\varepsilon)\Lambda} J'(\rho),$$

where  $J(\rho) = \int_M r^{-n} |D^\perp r|^2 \varphi(r/\rho) d\mu$ . Then integrating from  $\sigma$  to  $\rho$  as in the case  $\underline{H} = 0$  and then letting  $\varepsilon \downarrow 0$  as we did before, we obtain (analogous to 3.8)

$$e^{F(\rho)} \frac{\mu_V(B_\rho(\xi))}{\omega_n \rho^n} - e^{F(\sigma)} \frac{\mu_V(B_\sigma(\xi))}{\omega_n \sigma^n} = \omega_n^{-1} G(\sigma, \rho) \int_{B_\rho(\xi) \setminus B_\sigma(\xi)} r^{-n} |D^\perp r|^2 d\mu_V,$$

with  $G(\sigma, \rho) \in [e^{-\Lambda}, e^{\Lambda}]$ . Thus we have proved the following:

**3.20 Theorem.** *If  $U$  is open in  $\mathbb{R}^{n+k}$ , if  $B_R(\xi) \subset U$  and  $V$  has generalized mean curvature vector  $\underline{H}$  in  $U$  with  $|\underline{H}| \leq \Lambda$ , then*

$$e^{F(\rho)} \frac{\mu_V(B_\rho(\xi))}{\omega_n \rho^n} - e^{F(\sigma)} \frac{\mu_V(B_\sigma(\xi))}{\omega_n \sigma^n} = \omega_n^{-1} G(\sigma, \rho) \int_{B_\rho(\xi) \setminus B_\sigma(\xi)} r^{-n} |D^\perp r|^2 d\mu_V,$$

for all  $0 < \sigma \leq \rho < R$ , where  $F(\rho) \in [-\frac{\Lambda\rho}{R}, \frac{\Lambda\rho}{R}]$  and  $G(\sigma, \rho) \in [e^{-\Lambda}, e^{\Lambda}]$  for all  $0 < \sigma \leq \rho < R$ .

**3.21 Remark.** Since  $|F(\rho)| \leq \Lambda\rho/R$  in the above theorem, we again conclude that  $\Theta^n(\mu_V, \xi)$  exists for all  $\xi \in U$  and is an upper semi-continuous function on  $U$  by merely notational modifications to the previous argument for  $\underline{H} = 0$ .

We conclude this section with some variants of the above computations which will be important in our later discussion of local conical approximation and elsewhere.

Let  $B_R(\xi) \subset U$  and  $V = \underline{v}(M, \theta)$  have generalized mean curvature  $\underline{H} \in L^1(\mu_V)$  on  $\check{B}_R(\xi)$ . We let  $0 < \sigma < \rho \leq R$  and we observe that if we let  $\varepsilon \downarrow 0$  in 3.17, then we obtain, in the distribution sense,

$$\begin{aligned} 3.22 \quad \rho \frac{d}{d\rho} \int_{B_\rho(\xi)} h |\nabla^M r|^2 d\mu_V = \\ n \int_{B_\rho(\xi)} h d\mu_V + \int_{B_\rho(\xi)} (h \underline{H} + \nabla^M h) \cdot (x - \xi) d\mu_V. \end{aligned}$$

We also observe that the Co-area Formula 2.9 of Ch.3 with  $f(x) = |x - \xi| (= r)$  (in which case  $J_f^M = |\nabla^M r|$ ) and with  $M \cap B_\rho(\xi)$  in place of  $M$  implies

$$3.23 \quad \int_{B_\rho(\xi)} g |\nabla^M r| d\mu_V = \int_0^\rho \int_{\partial B_t(\xi)} g dv dt$$

for any non-negative  $\mathcal{H}^n$ -measurable function  $g$ , where  $dv = \theta d\mathcal{H}^{n-1}$ , so the left side is an absolutely continuous function of  $\rho$  and

$$3.24 \quad \frac{d}{d\rho} \int_{B_\rho(\xi)} g |\nabla^M r| d\mu_V = \int_{\partial B_\rho(\xi)} g dv,$$

for  $\mathcal{L}^1$ -a.e.  $\rho \in (0, R)$ . Hence, taking  $g = h |\nabla^M r|$ , 3.22 implies

$$3.25 \quad \int_{\partial B_\rho(\xi)} r h |\nabla^M r| dv = n \int_{B_\rho(\xi)} h d\mu_V + \int_{B_\rho(\xi)} (h \underline{H} + \nabla^M h) \cdot (x - \xi) d\mu_V$$

for  $\mathcal{L}^1$ -a.e.  $\rho \in (0, R)$ .

If we now subtract the inequality 3.25 with  $\sigma < \rho$  in place of  $\rho$  from the same inequality 3.25, then we get

$$\begin{aligned} 3.26 \quad & \int_{\partial B_\rho(\xi)} rh|\nabla^M r| dv - \int_{\partial B_\sigma(\xi)} rh|\nabla^M r| dv \\ &= n \int_{B_\rho(\xi) \setminus B_\sigma(\xi)} h d\mu_V + \int_{B_\rho(\xi) \setminus B_\sigma(\xi)} (h\underline{H} + \nabla^M h) \cdot (x - \xi) d\mu_V, \end{aligned}$$

and then by replacing  $h$  by  $r^{-n}h$  and observing that  $(x - \xi) \cdot \nabla^M r^{-n} = -nr^{-n}|\nabla^M r|^2 = -nr^{-n} + nr^{-n}|D^\perp r|^2$ , we obtain

$$\begin{aligned} 3.27 \quad & \int_{\partial B_\rho(\xi)} r^{1-n}h|\nabla^M r| dv - \int_{\partial B_\sigma(\xi)} r^{1-n}h|\nabla^M r| dv - n \int_{B_\rho(\xi) \setminus B_\sigma(\xi)} r^{-n}|D^\perp r|^2 h d\mu_V \\ &= \int_{B_\rho(\xi) \setminus B_\sigma(\xi)} r^{-n}(h\underline{H} + \nabla^M h) \cdot (x - \xi) d\mu_V. \end{aligned}$$

Now we take  $h : \mathbb{R}^{n+k} \setminus \{\xi\} \rightarrow [0, 1]$  to be a homogeneous degree zero  $C^1$  function of the variable  $x - \xi$  and let  $b \geq 1$  be such that

$$3.28 \quad |Dh(x - \xi)| \leq br^{-1}.$$

Observe that then  $(x - \xi) \cdot Dh = 0$ , and so  $(x - \xi) \cdot \nabla^M h = (x - \xi) \cdot p_{T_x M}(Dh) = p_{T_x M}(x - \xi) \cdot Dh = ((x - \xi) - p_{(T_x M)^\perp}(x - \xi)) \cdot Dh = -p_{(T_x M)^\perp}(x - \xi) \cdot Dh = -rD^\perp r \cdot Dh$ , so in fact

$$3.29 \quad |(x - \xi) \cdot \nabla^M h| \leq b|D^\perp r|,$$

with  $b$  as in 3.28. Then, since  $r = \rho$  on  $\partial B_\rho(\xi)$ , 3.27 implies

$$\begin{aligned} 3.30 \quad & \left| \rho^{1-n} \int_{\partial B_\rho(\xi)} h|\nabla^M r| dv - \sigma^{1-n} \int_{\partial B_\sigma(\xi)} h|\nabla^M r| dv - n \int_{B_\rho(\xi) \setminus B_\sigma(\xi)} r^{-n}|D^\perp r|^2 h d\mu_V \right| \\ & \leq \int_{B_\rho(\xi) \setminus B_\sigma(\xi)} (r^{1-n}|\underline{H}| + br^{-n}|D^\perp r|) d\mu_V. \end{aligned}$$

## 4 Monotonicity Formulae for $L^p$ Mean Curvature

Here we continue to assume that  $V = \underline{v}(M, \theta)$  is a rectifiable varifold in  $U$  ( $U$  open in  $\mathbb{R}^{n+k}$ ) with generalized mean curvature vector  $\underline{H}$  in  $U$ , but now we assume  $\underline{H}$  is merely in  $L^p$  function rather than  $L^\infty$  as in the previous section.

We begin with the inequalities 3.19, but now we assume only an  $L^p$  condition with  $p > n$  instead of a bound on  $H$ . Specifically we assume

$$4.1 \quad p > n \text{ and } \left( R^{p-n} \int_{B_R(\xi)} |H|^p d\mu_V \right)^{1/p} \leq \Lambda,$$

where  $\Lambda \geq 0$ . Observe that then by using Hölder's inequality we have (since  $\varphi(r/\rho) = 0$  for  $r < (1 + \varepsilon)\rho$ )

$$\begin{aligned}
 & \left| \rho^{-n} \int_U \rho^{-1}(x - \xi) \cdot \underline{H} \varphi(r/\rho) d\mu_V \right| \\
 & \leq (1 + \varepsilon) \rho^{-n} \|H\|_{L^p(\mu_V \llcorner B_R(\xi))} (I(\rho))^{1-1/p} \\
 & = (1 + \varepsilon) R^{-1} (\rho/R)^{-n/p} \left( R^{p-n} \int_{B_R(\xi)} |H|^p d\mu_V \right)^{1/p} (\rho^{-n} I(\rho))^{1-1/p} \\
 & \leq (1 + \varepsilon) \Lambda R^{-1} (\rho/R)^{-n/p} \left( \frac{1}{p} + \rho^{-n} I(\rho) \right), \quad \rho < R/(1 + \varepsilon),
 \end{aligned}$$

where at the last step we used  $a^{1-1/p} \leq \frac{1}{p} + a$ , valid for  $a \geq 0$ , as one checks by observing that the function  $f(a) = a^{1-1/p} - (\frac{1}{p} + \frac{p-1}{p}a)$ ,  $a \geq 0$ , attains a maximum value of zero at  $a = 1$ . Thus 3.19 implies

$$4.2 \quad (1 + \varepsilon)^{-n} J'(\rho) \leq \frac{d}{d\rho} (\rho^{-n} I(\rho)) + F_0(\rho) \left( \frac{1}{p} + \rho^{-n} I(\rho) \right) \leq J'(\rho),$$

where (as in 3.6)  $J(\rho) = \int_M r^{-n} |D^\perp r|^2 \varphi(r/\rho) d\mu_V$  and

$$|F_0(\rho)| \leq (1 + \varepsilon) \Lambda R^{-1} (\rho/R)^{-n/p}.$$

Observe that then  $F(\rho) = \int_0^\rho F_0(t) dt$  satisfies

$$|F(\rho)| \leq (1 + \varepsilon) \Lambda \kappa (\rho/R)^{1-n/p} \leq (1 + \varepsilon) \Lambda \kappa, \quad \kappa = p/(p - n),$$

so, after multiplying in 4.2 by the integrating factor  $e^{F(\rho)}$ , we obtain

$$(1 + \varepsilon)^{-n} e^{-(1+\varepsilon)\kappa\Lambda} J'(\rho) \leq \frac{d}{d\rho} (e^{F(\rho)} \rho^{-n} I(\rho) + \frac{1}{p} e^{F(\rho)}) \leq e^{(1+\varepsilon)\kappa\Lambda} J'(\rho).$$

Hence, after integrating over the interval  $[\sigma, \rho]$  and letting  $\varepsilon \downarrow 0$ ,

$$\begin{aligned}
 4.3 \quad & \left( e^{F(\rho)} \frac{\mu_V(B_\rho(\xi))}{\omega_n \rho^n} + \frac{1}{p} (e^{F(\rho)} - 1) \right) - \left( e^{F(\sigma)} \frac{\mu_V(B_\sigma(\xi))}{\omega_n \sigma^n} + \frac{1}{p} (e^{F(\sigma)} - 1) \right) \\
 & = G(\sigma, \rho) \int_{B_\rho(\xi) \setminus B_\sigma(\xi)} r^{-n} |D^\perp r|^2 d\mu_V
 \end{aligned}$$

with

$$4.4 \quad e^{-\kappa\Lambda} \leq G(\sigma, \rho) \leq e^{\kappa\Lambda}, \quad |F(\rho)| \leq \Lambda \kappa (\rho/R)^{1-n/p}, \quad \kappa = p/(p - n),$$

for all  $0 < \sigma < \rho < R$ , provided  $B_R(\xi) \subset U$ . In particular

$$4.5 \quad e^{F(\rho)} \frac{\mu_V(B_\rho(\xi))}{\omega_n \rho^n} + \frac{1}{p} (e^{F(\rho)} - 1) \text{ is increasing in } \rho, \quad 0 < \rho \leq R,$$

Since  $e^{F(\rho)} \rightarrow 1$  as  $\rho \downarrow 0$ , 4.3, 4.4 and 4.5 enable us to argue precisely as in §3, to conclude that  $\Theta^n(\mu_V, \xi) = \lim_{\sigma \downarrow 0} (\omega_n \sigma^n)^{-1} \mu_V(B_\sigma(\xi))$  exists  $\forall \xi \in U$  and

$$\begin{aligned} 4.6 \quad \left( e^{F(\rho)} \frac{\mu_V(B_\rho(\xi))}{\omega_n \rho^n} + \frac{1}{\rho} (e^{F(\rho)} - 1) \right) - \Theta^n(\mu_V, \xi) \\ = G(\rho) \int_{B_\rho(\xi)} r^{-n} |D^\perp r|^2 d\mu_V, \end{aligned}$$

with  $F(\rho), G(\rho)$  as in 4.4, so in particular  $\int_{B_\rho(\xi)} r^{-n} |D^\perp r|^2 d\mu_V < \infty$ . Also, again following the argument of §3,

$$4.7 \quad \Theta^n(\mu_V, \xi) \text{ is an upper semi-continuous function of } \xi \in U,$$

and, as in 3.13,  $V = \underline{v}(M, \theta)$  has a “canonical representative”  $(M_V, \Theta_V)$ :

$$4.8 \quad V = \underline{v}(M_V, \Theta_V), \quad \Theta_V(x) = \Theta^n(\mu_V, x), \quad M_V = \{x \in U : \Theta^n(\mu_V, x) > 0\}.$$

**4.9 Remarks: (1)** In the case of  $\underline{H} \in L^p_{\text{loc}}(\mu_V)$  with  $p > n$ , if  $\theta \geq 1$   $\mu_V$ -a.e. in  $U$ , then, by upper semi-continuity,  $\Theta^n(\mu_V, x) \geq 1$  at *each* point of  $\text{spt } \mu_V \cap U$ , and hence in this case the canonical representative  $M_V$  in 4.8 is just the *closed* set  $\text{spt } V$ , because  $M_V = \{x \in U : \Theta_V(x) > 0\} = \{x \in U : \Theta_V(x) \geq 1\} = \text{spt } \mu_V$ .

**(2)** Notice that as  $p \rightarrow \infty$ , 4.3, 4.4 yield 3.20.

**(3)** If  $\Gamma > 0$  and  $\rho^{-n} \mu_V(B_\rho(\xi)) \leq \Gamma$ , then 4.3 gives bounds of the form  $\mu(B_\sigma(\xi)) \leq \beta \sigma^n$  for  $0 < \sigma \leq R$ , with  $\beta = \beta(\Lambda, \Gamma)$ . It follows that

$$\int_{B_\rho(\xi) \setminus B_\sigma(\xi)} |x - \xi|^{-\alpha} d\mu \leq \begin{cases} n\beta(n-\alpha)^{-1}(\rho^{n-\alpha} - \sigma^{n-\alpha}), & 0 < \alpha < n \\ \beta \log(\rho/\sigma), & \alpha = n. \end{cases}$$

for any  $0 < \sigma < \rho < R$ . This is proved by using the following general fact with  $f(x) = |x - \xi|^{-1}$ ,  $t_0 = \rho^{-1}$ , and with  $n - \alpha$  in place of  $\alpha$ .

**4.10 Lemma.** *If  $X$  is an abstract space,  $\mu$  is a measure on  $X$  with  $\mu(X) < \infty$ ,  $f \geq 0$ ,  $f$   $\mu$ -measurable, and  $A_t = \{x \in X : f(x) > t\}$ , then*

$$\begin{aligned} \int_{t_0}^{\infty} t^{\alpha-1} \mu(A_t) dt &= \alpha^{-1} \int_{A_{t_0}} (f^\alpha - t_0^\alpha) d\mu, \quad 0 < \alpha < n \\ \int_{t_0}^{\infty} t^{-1} \mu(A_t) dt &= \int_{A_{t_0}} \log(f/t_0) d\mu \end{aligned}$$

for each  $t_0 \geq 0$ .

**Proof:** Since  $\int_{t_0}^{\infty} t^{\alpha-1} \mu(A_t) dt = \int_{t_0}^{\infty} \int_{A_{t_0}} t^{\alpha-1} \chi_{A_t}(x) d\mu(x) dt$ , this is proved simply by applying Fubini's theorem on the product space  $A_{t_0} \times [t_0, \infty)$ .  $\square$

## 5 Local Conical Approximation

Here we want to derive an important theorem concerning conical approximation of  $V$  near an arbitrary point of  $\text{spt } V$ , assuming  $V = \underline{v}(M, \theta)$  has generalized mean curvature in  $L^p$  with  $p > n$ .

**5.1 Theorem (Conical Approximation.)** *Suppose  $\Lambda > 0$ ,  $p > n$ ,  $\lambda \in (0, \frac{1}{4}]$ ,  $\delta \in [0, \frac{1}{16}]$ , and  $V = \underline{v}(M, \theta)$  has generalized mean curvature  $\underline{H}$  in a neighborhood of  $B_1(0)$  and satisfies the hypotheses*

$$0 \in \text{spt } \mu_V, \quad \Theta^n(\mu_V, 0) \leq \Gamma,$$

$$\omega_n^{-1} \mu_V(B_1(0)) \leq \Theta^n(\mu_V, 0) + \delta, \quad \left( \int_{B_1(0)} |\underline{H}|^p d\mu_V \right)^{1/p} \leq \delta.$$

Then

$$\frac{\mu_V(B_{(1-\delta^{\frac{1}{4}})\rho}(\xi))}{\omega_n \rho^n} - C\kappa(\delta, \lambda) \leq \frac{\mu_V(B_{\tau\rho}(\tau\xi))}{\omega_n (\tau\rho)^n} \leq \frac{\mu_V(B_{(1+\delta^{\frac{1}{4}})\rho}(\xi))}{\omega_n \rho^n} + C\kappa(\delta, \lambda)$$

for all  $\xi \in \check{B}_1(0) \setminus B_{2\lambda}(0)$  ( $\xi$  does not need to be in  $\text{spt } \mu_V$ ), all  $\tau \in [\lambda, 1]$ , and all  $\rho$  with  $\lambda \leq \rho \leq \min\{|\xi| - \lambda, (1 + \delta^{\frac{1}{4}})^{-1}(1 - |\xi|)\}$ , where  $C = C(n, p, \Gamma)$  and where

$$\kappa(\delta, \lambda) = \lambda^{-n} \delta + \delta^{\frac{1}{4}} |\log \lambda|.$$

**5.2 Remarks:** (1) In the special case when  $\underline{H} = 0$  and  $\omega_n^{-1} \mu_V(\check{B}_1(0)) = \Theta^n(\mu_V, 0)$  (which implies  $(\omega_n \rho^n)^{-1} \mu_V(\check{B}_\rho(0)) = \Theta^n(\mu_V, 0) \forall \rho > 0$  by monotonicity 3.9) we can apply the above with  $\delta = 0$  and with arbitrarily small  $\lambda$ , so we can let  $\rho \downarrow 0$  in the conclusion to infer that  $\Theta^n(\mu_V, \tau\xi) = \Theta^n(\mu_V, \xi)$  for all  $\tau \in (0, 1]$ . Thus, by 4.8, in this case  $V \llcorner B_1(0)$  is a cone:  $(\eta_{0,\tau\#} V) \llcorner B_1(0) = V \llcorner B_1(0) \forall \tau \in (0, 1)$ .

(2) If  $V = \underline{v}(M, \theta)$  is any rectifiable  $n$ -varifold with generalized mean curvature  $\underline{H} \in L^p$ ,  $p > n$ , and if  $\xi \in \text{spt } \mu_V$ , then, for  $\sigma \in (0, 1)$ ,  $\eta_{\xi, \sigma\#} V$  has 0 in its support and generalized mean curvature with  $L^p$ -norm  $\rightarrow 0$  in  $B_1(0)$  as  $\sigma \downarrow 0$  and also  $\omega_n^{-1} \mu_{\eta_{\xi, \sigma\#} V}(B_1(0)) - \Theta(\mu_{\eta_{\xi, \sigma\#} V}, 0) = (\omega_n \sigma^n)^{-1} \mu_V(B_\sigma(\xi)) - \Theta^n(\mu_V, \xi) \rightarrow 0$  as  $\sigma \downarrow 0$ , so the above theorem is applicable to  $\eta_{\xi, \sigma\#} V$  with arbitrarily small  $\delta$  by taking  $\sigma > 0$  small enough.

Indeed if  $\Theta^n(\mu_V, \xi) \geq 1$  on  $\text{spt } \mu_V$ , the Allard compactness theorem (which will be proved in Theorem 5.8 of Ch. 8) guarantees that for each sequence  $\sigma_j \downarrow 0$  there is a subsequence  $\sigma_{j'}$  with  $\mu_{\eta_{\xi, \sigma_{j'}\#} V} \rightarrow \mu_T$  for some rectifiable  $n$ -varifold  $T$  which is stationary in  $\mathbb{R}^{n+k}$  and has multiplicity  $\geq 1$  on  $\text{spt } T$ . Of course by construction this  $T$  satisfies the hypotheses on the theorem with  $\delta = 0$  and  $\lambda \in (0, \frac{1}{2}]$  arbitrary, so  $T$  is a cone by the first remark above. Such a  $T$  is called a tangent cone of  $V$  at  $\xi$ . It is still an open (and important) question as to whether  $T$  is *unique* (i.e. whether or not  $T$  is independent of the choice of the sequence  $\sigma_j$  and the subsequence  $\sigma_{j'}$ ). Tangent cones will be more

systematically discussed in §5 of Ch. 8.

**Proof of 5.1:** We can assume  $\delta > 0$  and obtain  $\delta = 0$  as a limit. In view of the stated hypotheses we can apply 4.6 (with  $\Lambda = \delta$ ) to give

$$(1) \quad \int_{B_1(0)} r^{-n} |D^\perp r|^2 d\mu_V \leq C\delta, \quad C = C(n, p, \Gamma).$$

Let  $h : \mathbb{R}^{n+k} \setminus \{0\} \rightarrow [0, 1]$  be a homogeneous degree zero  $C^1$  function  $\mathbb{R}^{n+k} \setminus \{0\} \rightarrow [0, 1]$  with

$$(2) \quad |Dh(x)| \leq b/r$$

(as in 3.28 with  $\xi = 0$ ), where for the moment  $b \geq 1$  is arbitrary.

Let  $t \in [\lambda, 1]$ ,  $\tau \in [\lambda, 1)$ , and apply 3.30 with  $t, \tau t, 0$  in place of  $\rho, \sigma, \xi$  respectively. This gives

$$(3) \quad \left| t^{1-n} \int_{\partial B_t(0)} h |\nabla^M r| d\nu - (t\tau)^{1-n} \int_{\partial B_{\tau t}(0)} h |\nabla^M r| d\nu - n \int_{B_t(0) \setminus B_{\tau t}(0)} r^{-n} |D^\perp r|^2 h d\mu_V \right| \\ \leq \int_{B_t(0) \setminus B_{\tau t}(0)} r^{-n} (rh|\underline{H}| + b|D^\perp r|) d\mu_V \leq \int_{B_1(0) \setminus B_{\lambda^2}(0)} r^{-n} (rh|\underline{H}| + b|D^\perp r|) d\mu_V.$$

Take any  $t \in [|\xi| - \rho, |\xi| + \rho]$  (i.e. any  $t$  such that  $\partial B_t(0) \cap B_\rho(\xi) \neq \emptyset$ ), and observe that we can select the homogeneous degree zero  $C^1$  function  $h : \mathbb{R}^{n+k} \setminus \{0\} \rightarrow [0, 1]$  (depending on  $t$ ) such that  $h$  is identically 1 on  $\partial B_t(0) \cap B_\rho(\xi)$ ,  $h = 0$  on  $\partial B_t(0) \setminus B_{(1+\delta^{\frac{1}{4}})\rho}(\xi)$  and  $|Dh(x)| \leq 2/(r\delta^{\frac{1}{4}}\rho)$ , so (2) holds with  $b = 2/(\delta^{\frac{1}{4}}\rho)$ . Thus (3) implies

$$(4) \quad \tau^{1-n} \int_{\partial B_{\tau t}(0) \cap B_{\tau\rho}(\tau\xi)} |\nabla^M r| d\nu \leq \int_{\partial B_t(0) \cap B_{(1+\delta^{1/4})\rho}(\xi)} |\nabla^M r| d\nu \\ + t^{n-1} \int_{B_1(0) \setminus B_{\lambda^2}(0)} r^{-n} \left( r|\underline{H}| + 2\rho^{-1}\delta^{-1/4}|D^\perp r| \right) d\mu_V.$$

Similarly, considering  $t$  with  $\partial B_t(0) \cap B_{(1-\delta^{\frac{1}{4}})\rho}(\xi) \neq \emptyset$  and choosing another homogeneous degree zero  $C^1$  function  $h : \mathbb{R}^{n+k} \setminus \{0\} \rightarrow [0, 1]$  such that  $h$  is identically 1 on  $\partial B_t(0) \cap B_{(1-\delta^{\frac{1}{4}})\rho}(\xi)$ ,  $h = 0$  on  $\partial B_t(0) \setminus B_\rho(\xi)$  and again  $|Dh(x)| \leq 2/(r\delta^{\frac{1}{4}}\rho)$ , we obtain

$$(5) \quad \int_{\partial B_t(0) \cap B_{(1-\delta^{1/4})\rho}(\xi)} |\nabla^M r| d\nu \leq \tau^{1-n} \int_{\partial B_{\tau t}(0) \cap B_{\tau\rho}(\tau\xi)} |\nabla^M r| d\nu \\ + n t^{n-1} \int_{B_1(0)} r^{-n} |D^\perp r|^2 d\mu_V + t^{n-1} \int_{B_1(0) \setminus B_{\lambda^2}(0)} r^{-n} \left( r|\underline{H}| + 2\rho^{-1}\delta^{-\frac{1}{4}}|D^\perp r| \right) d\mu_V.$$

Observe that the term involving  $\underline{H}$  on the right of (4) and (5) can be estimated by the Hölder inequality and the fact that  $\|\underline{H}\|_{L^p(\mu_V \llcorner B_1(0))} \leq \delta$ :

$$(6) \quad \int_{B_1(0) \setminus B_{\lambda^2}(0)} r^{1-n} |\underline{H}| d\mu_V \leq \|r^{1-n}\|_{L^{\frac{p}{p-1}}(\mu_V \llcorner B_1(0) \setminus B_{\lambda^2}(0))} \|\underline{H}\|_{L^p(\mu_V \llcorner B_1(0))} \leq C\delta,$$

where  $C = C(n, p, \Gamma)$ , and where we used 4.9(3) with  $\alpha = (n-1)p/(p-1)(< n)$ .

Also to handle the term  $\int_{B_1(0) \setminus B_{\lambda^2}(0)} 2r^{-n} \rho^{-1} \delta^{-\frac{1}{4}} |D^\perp r| d\mu_V$  on the right of (4) and (5) we use the Cauchy inequality  $a \leq \frac{1}{2}\varepsilon + \frac{1}{2}a^2/\varepsilon$  with  $\varepsilon = \delta^{\frac{1}{4}}$ , so

$$(7) \quad \begin{aligned} & \int_{B_1(0) \setminus B_{\lambda^2}(0)} 2r^{-n} \rho^{-1} \delta^{-1/4} |D^\perp r| d\mu_V \\ & \leq \delta^{\frac{1}{4}} \int_{B_1(0) \setminus B_{\lambda^2}(0)} r^{-n} d\mu_V + \delta^{-\frac{3}{4}} \int_{B_1(0)} r^{-n} |D^\perp r|^2 d\mu_V \leq C\delta^{\frac{1}{4}} |\log \lambda|, \end{aligned}$$

with  $C = C(n, p, \Gamma)$ , where we again used 4.9(3) and also (1).

Now we integrate in the inequality (4) with respect to  $t$  over the interval  $[|\xi| - \rho, |\xi| + \rho]$  (which is integration over  $[\tau\xi - \tau\rho, \tau\xi + \tau\rho]$  with respect to  $\tau t$ ), and use the bounds (6), (7) together with the Coarea Formula 3.24. This gives

$$\frac{\int_{B_{\tau\rho}(\tau\xi)} |\nabla^M r|^2 d\mu_V}{\omega_n(\tau\rho)^n} \leq \frac{\int_{B_{(1+\delta^{1/4})\rho}(\xi)} |\nabla^M r|^2 d\mu_V}{\omega_n \rho^n} + C\delta^{\frac{1}{4}} |\log \lambda|, \quad C = C(n, p, \Gamma).$$

Hence (since  $|\nabla^M r|^2 = 1 - |D^\perp r|^2$ ), after another application of (1), noting that  $r \leq \tau$  in  $B_{\tau\rho}(\tau\xi)$ , so  $(\tau\rho)^{-n} \int_{B_{\tau\rho}(\tau\xi)} |D^\perp r|^2 d\mu_V \leq \lambda^{-n} \int_{B_1(0)} r^{-n} |D^\perp r|^2 d\mu_V$ ,

$$\frac{\mu_V(B_{\tau\rho}(\tau\xi))}{\omega_n(\tau\rho)^n} \leq \frac{\mu_V(B_{(1+\delta^{\frac{1}{4}})\rho}(\xi))}{\omega_n \rho^n} + C(\delta^{\frac{1}{4}} |\log \lambda| + \delta \lambda^{-n}), \quad C = C(n, p, \Gamma).$$

Similarly, integrating with respect to  $t$  over  $[|\xi| - (1 - \delta^{\frac{1}{4}})\rho, |\xi| + (1 - \delta^{\frac{1}{4}})\rho]$  in the inequality (5), and again using the bounds (6), (7) and (1), we obtain

$$\frac{\mu_V(B_{(1-\delta^{\frac{1}{4}})\rho}(\xi))}{\omega_n \rho^n} \leq \frac{\mu_V(B_{\tau\rho}(\tau\xi))}{\omega_n(\tau\rho)^n} + C(\delta^{\frac{1}{4}} |\log \lambda| + \delta \lambda^{-n}). \quad \square$$

## 6 Poincaré and Sobolev Inequalities

In this section<sup>3</sup> we continue to assume that  $V = \underline{v}(M, \theta)$  has generalized mean curvature  $\underline{H}$  in  $U$ , and we again write  $\mu$  for  $\mu_V$ . We shall also assume  $\theta \geq 1$   $\mu$ -a.e.  $x \in U$ , so that

<sup>3</sup>Note: The results of this section are not needed in the sequel

(by the comments in 4.9)  $\Theta^n(\mu, x) \geq 1$  everywhere in  $\text{spt } \mu \cap U$  if  $\underline{H} \in L^p_{\text{loc}}(\mu)$  for some  $p > n$ .

We start with the identity 3.17, which since  $|\nabla^M r|^2 = 1 - |D^\perp r|^2$ , can be written in the form

$$\begin{aligned} 6.1 \quad \frac{\partial}{\partial \rho}(\rho^{-n} \tilde{I}(\rho)) &= \rho^{-n} \frac{\partial}{\partial \rho} \int |(Dr)^\perp|^2 h \varphi(r/\rho) d\mu \\ &\quad + \rho^{-n-1} \int (x - \xi) \cdot [\nabla^M h + \underline{H}h] \varphi(r/\rho) d\mu \end{aligned}$$

where now  $\tilde{I}(\rho) = \int \varphi(r/\rho) h d\mu$ .

Thus

$$\frac{\partial}{\partial \rho} [\rho^{-n} \tilde{I}(\rho)] \geq \rho^{-n-1} \int (x - \xi) \cdot (\nabla^M h + \underline{H}h) \varphi(r/\rho) d\mu.$$

We can estimate the right-side  $R$  here in two ways: if  $|\underline{H}| \leq \Gamma$  we have

$$6.2 \quad R \geq -\rho^{-n-1} \int r |\nabla^M h| \varphi(r/\rho) d\mu - (\Gamma \rho) \rho^{-n} \tilde{I}(\rho).$$

Alternatively, without any assumption on  $\underline{H}$  we can clearly estimate

$$6.3 \quad R \geq -\rho^{-n-1} \int r (|\nabla^M h| + h |\underline{H}|) \varphi(r/\rho) d\mu.$$

If we use 6.2 in 6.1 and integrate (making use of 4.10) we obtain (after letting  $\varphi$  increase to the indicator function of  $(-\infty, 1)$  as before)

$$6.4 \quad \frac{1}{\omega_n \sigma^n} \int_{B_\sigma(\xi)} h d\mu \leq e^{\Lambda \rho} \left( \frac{1}{\omega_n \rho^n} \int_{B_\rho(\xi)} h d\mu + \frac{1}{n \omega_n} \int_{B_\rho(\xi)} \frac{|\nabla^M h|}{|x - \xi|^{n-1}} d\mu \right),$$

provided  $B_\rho(\xi) \subset U$  and  $0 < \sigma < \rho$ .

If instead we use 6.3 then we similarly get

$$\frac{1}{\omega_n \sigma^n} \int_{B_\sigma(\xi)} h d\mu \leq \frac{1}{\omega_n \rho^n} \int_{B_\rho(\xi)} h d\mu + \frac{1}{\omega_n} \int_\sigma^\rho \tau^{-n-1} \int_{B_\tau(\xi)} r (|\nabla^M h| + h |\underline{H}|) d\mu d\tau.$$

and hence (by 4.10 again)

$$6.5 \quad \frac{1}{\omega_n \sigma^n} \int_{B_\sigma(\xi)} h d\mu \leq \frac{1}{\omega_n \rho^n} \int_{B_\rho(\xi)} h d\mu + \frac{1}{n \omega_n} \int_{B_\rho(\xi)} \frac{(|\nabla^M h| + h |\underline{H}|)}{|x - \xi|^{n-1}} d\mu$$

provided  $B_\rho(\xi) \subset U$  and  $0 < \sigma < \rho$ .

If we let  $\sigma \downarrow 0$  in 6.4 then we get (since  $\Theta(\mu, \xi) \geq 1$  for  $\xi \in \text{spt } \mu$ )

$$h(\xi) \leq e^{\Lambda \rho} \left( \frac{1}{\omega_n \rho^n} \int_{B_\rho(\xi)} h d\mu + \frac{1}{n \omega_n} \int_{B_\rho(\xi)} \frac{|\nabla^M h|}{|x - \xi|^{n-1}} \right), \quad \xi \in \text{spt } \mu, \quad B_\rho(\xi) \subset U.$$

We now state our Poincaré-type inequality.

**6.6 Theorem.** *Suppose  $h \in C^1(U)$ ,  $h \geq 0$ ,  $B_{2\rho}(\xi) \subset U$ ,  $|\underline{H}| \leq \Lambda$ ,  $\theta \geq 1$   $\mu$ -a.e. in  $U$  and*

$$\mu\{x \in B_\rho(\xi) : h(x) > 0\} \leq (1 - \alpha)\omega_n \rho^n, \quad e^{\Lambda\rho} \leq 1 + \alpha$$

*for some  $\alpha \in (0, 1)$ . Suppose also that*

$$(\ddagger) \quad \mu(B_{2\rho}(\xi)) \leq \Gamma \rho^n, \quad \Gamma > 0.$$

*Then there are constants  $\beta = \beta(n, \alpha, \Gamma) \in (0, 1/2)$  and  $c = c(n, \alpha, \Gamma) > 0$  such that*

$$\int_{B_{\beta\rho}(\xi)} h \, d\mu \leq c\rho \int_{B_\rho(\xi)} |\nabla^M h| \, d\mu.$$

**Proof:** To begin we take  $\beta$  to be an arbitrary parameter in  $(0, \frac{1}{2})$  and apply 6.5 with  $\eta \in B_{\beta\rho}(\xi) \cap \text{spt } \mu$  in place of  $\xi$ . This gives

$$\begin{aligned} (1) \quad h(\eta) &\leq e^{\Lambda(1-\beta)\rho} \left( \frac{1}{\omega_n((1-\beta)\rho)^n} \int_{B_{(1-\beta)\rho}(\eta)} h \, d\mu + \frac{1}{n\omega_n} \int_{B_{(1-\beta)\rho}(\eta)} \frac{|\nabla^M h|}{|x - \eta|^{n-1}} \, d\mu \right) \\ &\leq e^{\Lambda\rho} \left( \frac{1}{\omega_n((1-\beta)\rho)^n} \int_{B_\rho(\xi)} h \, d\mu + \frac{1}{n\omega_n} \int_{B_\rho(\xi)} \frac{|\nabla^M h|}{|x - \eta|^{n-1}} \, d\mu \right). \end{aligned}$$

Now let  $\gamma$  be a fixed  $C^1$  non-decreasing function on  $\mathbb{R}$  with  $\gamma(t) = 0$  for  $t \leq 0$  and  $\gamma(t) \leq 1$  everywhere, and apply (1) with  $\gamma(h - t)$  in place of  $h$ , where  $t \geq 0$  is fixed. Then by (1)

$$\gamma(h(\eta) - t) \leq \frac{1 + \alpha}{n\omega_n} \int_{B_\rho(\xi)} \frac{\omega'(h - t) |\nabla^M h|}{|x - \eta|^{n-1}} \, d\mu + (1 - \alpha^2)(1 - \beta)^{-n}.$$

Selecting  $\beta$  small enough so that  $(1 - \beta)^{-n}(1 - \alpha^2) \leq 1 - \alpha^2/2$ , we thus get

$$(2) \quad \frac{\alpha^2}{2} \leq \frac{1 + \alpha}{n\omega_n} \int_{B_\rho(\xi)} \frac{\gamma'(h - t) |\nabla^M h|}{|x - \eta|^{n-1}} \, d\mu$$

for any  $\eta \in B_{\beta\rho}(\xi) \cap \text{spt } \mu$  such that  $\gamma(h(\eta) - t) \geq 1$ . Now let  $\varepsilon > 0$  and choose  $\gamma$  such that  $\gamma(t) \equiv 1$  for  $t \geq 1 + \varepsilon$ . Then (2) implies

$$1 \leq C \int_{B_\rho(\xi)} \frac{\gamma'(h - t) |\nabla^M h|}{|x - \eta|^{n-1}} \, d\mu, \quad \eta \in B_{\beta\rho}(\xi) \cap A_{t+\varepsilon},$$

where  $A_\tau = \{y \in \text{spt } \mu : h(y) > \tau\}$ . Integrating over  $A_{t+\varepsilon} \cap B_{\beta\rho}(\xi)$  we thus get (after interchanging the order of integration on the right)

$$\begin{aligned} \mu(A_{t+\varepsilon} \cap B_{\beta\rho}(\xi)) &\leq C \int_{B_\rho(\xi)} \gamma'(h(x) - t) |\nabla^M h(x)| \left( \int_{B_{\beta\rho}(\xi)} \frac{1}{|x - \eta|^{n-1}} \, d\mu(\eta) \right) \, d\mu(x) \\ &\leq C\Gamma\rho \int_{B_\rho(\xi)} \gamma'(h - t) |\nabla^M h| \, d\mu \end{aligned}$$

by hypothesis 6.6(‡) and Remark 4.9(3). Since  $\gamma'(h(x) - t) = -\frac{\partial}{\partial t}\gamma(h(x) - t)$  we can now integrate over  $t \in (0, \infty)$  to obtain (from 4.10) that

$$\int_{A_\varepsilon \cap B_{\beta\rho}(\xi)} (h - \varepsilon) \leq C \Gamma \rho \int_{B_\rho(\xi)} |\nabla^M h| d\mu.$$

Letting  $\varepsilon \downarrow 0$ , we have the required inequality.  $\square$

**Remark:** If we drop the assumption that  $\theta \geq 1$ , then the above argument still yields

$$\int_{\{x: \theta(x) \geq 1\} \cap B_{\beta\rho}(\xi)} h d\mu \leq C \rho \int_{B_\rho(\xi)} |\nabla^M h| d\mu.$$

We can also prove a Sobolev inequality as follows.

**6.7 Theorem.** *Suppose  $h \in C_0^1(U)$ ,  $h \geq 0$ , and  $\theta \geq 1$   $\mu$ -a.e. in  $U$ . Then*

$$(\ddagger) \quad \left( \int h^{\frac{n}{n-1}} d\mu \right)^{\frac{n-1}{n}} \leq C \int (|\nabla^M h| + h|\underline{H}|) d\mu, \quad c = c(n).$$

**Note:**  $C$  does not depend on  $k$ .

In the proof we shall need the following simple calculus lemma.

**6.8 Lemma.** *Suppose  $f, g$  are bounded and increasing on  $(0, \infty)$  and*

$$1 \leq \sigma^{-n} f(\sigma) \leq \rho^{-n} f(\rho) + \int_0^\rho \tau^{-n} g(\tau) d\tau, \quad 0 < \sigma < \rho < \infty.$$

*then  $\exists \rho$  with  $0 < \rho < \rho_0 = 2(f(\infty))^{1/n}$  ( $f(\infty) = \lim_{\rho \uparrow \infty} f(\rho)$ ) such that*

$$(\ddagger) \quad f(5\rho) \leq \frac{1}{2} 5^n \rho_0 g(\rho).$$

**Proof of Lemma:** Suppose  $(\ddagger)$  is false for each  $\rho \in (0, \rho_0)$ . Then

$$\begin{aligned} 1 &\leq \sup_{0 < \sigma < \rho_0} \sigma^{-n} f(\sigma) \leq \rho_0^{-n} f(\rho_0) + \frac{2 \cdot 5^{-n}}{\rho_0} \int_0^{\rho_0} \rho^{-n} f(5\rho) d\rho \\ &= \rho_0^{-n} f(\rho_0) + \frac{2}{5\rho_0} \int_0^{\tilde{5}\rho_0} \rho^{-n} f(\rho) d\rho \\ &= \rho_0^{-n} f(\rho_0) + \frac{2}{5\rho_0} \left( \int_0^{\rho_0} \rho^{-n} f(\rho) d\rho + \int_{\rho_0}^{\tilde{5}\rho_0} \rho^{-n} f(\rho) d\rho \right) \\ &\leq \rho_0^{-n} f(\infty) + \frac{2}{5} \sup_{0 < \rho < \rho_0} \rho^{-n} f(\rho) + \frac{2}{5(n-1)} \rho_0^{-n} f(\infty). \end{aligned}$$

Thus

$$\frac{1}{2} \leq \frac{1}{2} \sup_{0 < \sigma < \rho_0} \sigma^{-n} f(\sigma) < 2\rho_0^{-n} f(\infty) = 2^{-n},$$

which is a contradiction since  $n \geq 2$ .  $\square$

**Continuation of the proof of 6.7:** First note that because  $h$  has compact support in  $U$ , the formula 6.5 is actually valid here for all  $0 < \sigma < \rho < \infty$ . Hence we can apply the above lemma with the choices

$$\begin{aligned} f(\rho) &= \omega_n^{-1} \int_{B_\rho(\xi)} h \, d\mu, \\ g(\rho) &= \omega_n^{-1} \int_{B_\rho(\xi)} (|\nabla^M h| + h|\underline{H}|) \, d\mu, \end{aligned}$$

provided that  $\xi \in \text{spt } \mu$  and  $h(\xi) \geq 1$ .

Thus for each  $\xi \in \{x \in \text{spt } \mu : h(x) \geq 1\}$  we have  $\rho < 2(\omega_n^{-1} \int_M h \, d\mu)^{1/n}$  such that

$$(1) \quad \int_{B_{5\rho}(\xi)} h \, d\mu \leq 5^n (\omega_n^{-1} \int_M h \, d\mu)^{1/n} \int_{B_\rho(\xi)} (|\nabla^M h| + h|\underline{H}|) \, d\mu.$$

Using the covering Lemma (3.4 of Ch. 1) we can select disjoint balls  $B_{\rho_1}(\xi_1)$ ,  $B_{\rho_2}(\xi_2)$ ,  $\dots$ ,  $\xi_1 \in \{\xi \in \text{spt } \mu : h(\xi) \geq 1\}$  such that  $\{\xi \in M : h(\xi) \geq 1\} \subset \cup_{j=1}^\infty B_{5\rho_j}(\xi_j)$ . Then applying (1) and summing over  $j$  we have

$$\int_{\{x \in \text{spt } \mu : h(x) \geq 1\}} h \, d\mu \leq 5^n (\omega_n^{-1} \int_M h \, d\mu)^{1/n} \int_M (|\nabla^M h| + h|\underline{H}|) \, d\mu.$$

Next let  $\gamma$  be a non-decreasing  $C^1(\mathbb{R})$  function such that  $\gamma(t) \equiv 1$  for  $t > \varepsilon$  and  $\gamma(t) \equiv 0$  for  $t < 0$ , and use this with  $\gamma(h-t)$ ,  $t \geq 0$ , in place of  $h$ . This gives

$$\mu(M_{t+\varepsilon}) \leq 5^n \omega_n (\mu(M_t))^{1/n} \int_M (\gamma'(h-t)|\nabla^M h| + \gamma(h-t)|\underline{H}|) \, d\mu,$$

where

$$M_\alpha = \{x \in M : h(x) > \alpha\}, \quad \alpha \geq 0.$$

Multiplying this inequality by  $(t+\varepsilon)^{\frac{1}{n-1}}$  and using the trivial inequality  $(t+\varepsilon)^{\frac{1}{n-1}} \mu(M_t) \leq \int_{M_t} (h+\varepsilon)^{\frac{1}{n-1}} \, d\mu$  on the right, we then get

$$\begin{aligned} (t+\varepsilon)^{\frac{1}{n-1}} \mu(M_{t+\varepsilon}) &\leq \\ &5^n \omega_n^{-1/n} \left( \int_M (h+\varepsilon)^{\frac{n}{n-1}} \, d\mu \right)^{\frac{1}{n}} \left( -\frac{d}{dt} \int_M \gamma(\xi-t) |\nabla^M h| + \int_{M_t} |\underline{H}| \, d\mu \right). \end{aligned}$$

Now integrate of  $t \in (0, \infty)$  and use 4.10. This then gives

$$\int_{M_\varepsilon} (h^{\frac{n}{n-1}} - \varepsilon^{\frac{n}{n-1}}) \, d\mu \leq 5^{n+1} \omega_n^{-1/n} \left( \int_M (h+\varepsilon)^{\frac{n}{n-1}} \right)^{\frac{1}{n}} \int_M (|\nabla^M h| + h|\underline{H}|) \, d\mu.$$

The theorem (with  $C = 5^{n+1} \omega_n^{-1/n}$ ) now follows by letting  $\varepsilon \downarrow 0$ .  $\square$

**6.9 Remark:** Note that the inequality of 6.7 is valid without any boundedness hypothesis on  $\underline{H}$ : it suffices that  $\underline{H}$  is merely in  $L^1_{\text{loc}}(\mu)$ .

## 7 Miscellaneous Additional Consequences

Here  $V = \underline{v}(M, \theta)$  is a rectifiable  $n$ -varifold in  $\mathbb{R}^{n+k}$  with generalized mean curvature  $\underline{H}$  in  $U$ ,  $U \subset \mathbb{R}^{n+k}$  open, as in Definition 3.15 of the present chapter. We first derive a preliminary property for  $V$  in case  $\underline{H}$  is bounded.

**7.1 Lemma.** *Suppose  $U = \mathbb{R}^{n+k} \setminus B_R(\xi)$  and  $V \llcorner U$  has  $L^1_{\text{loc}}(\mu_V)$  generalized mean curvature  $\underline{H}$  in  $U$  with  $n^{-1}|\underline{H}(x) \cdot (x - \xi)| < 1$   $\mu_V$ -a.e. in  $U$ , and suppose also that  $\text{spt } V$  is compact. Then*

$$\text{spt } V \subset B_R(\xi).$$

(i.e.  $V \llcorner U = 0$ .)

**Proof:** Since  $\text{spt } V$  is compact it is easily checked that the identity (see §3)

$$n \int \gamma(r) d\mu_V + \int r \gamma'(r) (1 - |D^\perp r|^2) d\mu_V = - \int \underline{H}(x) \cdot (x - \xi) \gamma(r) d\mu_V(x)$$

(where  $r = |x - \xi|$ ) actually holds for any non-negative increasing  $C^1(\mathbb{R})$  function  $\gamma$  with  $\gamma(t) = 0$  for  $t \leq R + \varepsilon$ . ( $\varepsilon > 0$  arbitrary.) We see this as in §3, by substituting  $X(x) = \psi(x)\gamma(r)(x - \xi)$ , where  $\psi \in C_c^1(\mathbb{R}^{n+k})$  with  $\psi \equiv 1$  in a neighborhood of  $\text{spt } V$ . Since  $1 - |D^\perp r|^2 \geq 0$  and  $|\underline{H} \cdot (x - \xi)| < n \mu_V$ -a.e., we thus deduce  $\int \gamma(r) d\mu_V = 0$  for any such  $\gamma$ . Since we may select  $\gamma$  so that  $\gamma(t) > 0$  for  $t > R + \varepsilon$ , we thus conclude  $\text{spt } V (\equiv \text{spt } \mu_V) \subset B_{R+\varepsilon}(\xi)$ . Because  $\varepsilon > 0$  was arbitrary, this proves the lemma.

**7.2 Theorem (Convex hull property for stationary varifolds.)** *Suppose  $K \subset \mathbb{R}^{n+k}$  is compact, let  $U = \mathbb{R}^{n+k} \setminus K$ , and  $V = \underline{v}(M, \theta)$  is a stationary rectifiable  $n$ -varifold in  $U$  with  $\text{spt } V$  is bounded. Then*

$$\text{spt } V \subset \text{convex hull of } K.$$

**Proof:** The convex hull of  $K$  can be written as the intersection of all balls  $B_R(\xi)$  with  $K \subset B_R(\xi)$ . Hence the result follows immediately from 7.1.  $\square$

Next we want to discuss local Hausdorff distance sense convergence of the support of a sequence of stationary rectifiable varifolds or more generally a sequence of rectifiable varifolds with mean curvature in  $L^p$  for  $p > n$ .

We first recall the definition of local Hausdorff distance sense convergence.

**7.3 Definition:** If  $F, F_1, F_2, \dots$  are subsets of the open set  $U$  then we say  $F_j$  converges to  $F$  locally in the Hausdorff distance sense in  $U$  if for each compact  $K \subset U$  and each  $\delta > 0$  we have  $j_0 = j_0(\delta, K)$  such that

$$\begin{cases} F_j \cap K \subset \{x \in U : \text{dist}(x, F) < \delta\} \\ F \cap K \subset \{x \in U : \text{dist}(x, F_j) < \delta\} \end{cases}$$

for all  $j \geq j_0$ .

**7.4 Theorem (Distance Sense Cvce. of Supports.)** *Let  $p > n$  and let  $V_j = \underline{v}(M_j, \theta_j)$ ,  $j = 1, 2, \dots$ , be rectifiable varifolds in  $U$  with generalized mean curvature vector  $\underline{H}_j \in L^p_{\text{loc}}(\mu_{V_j})$  in  $U$  with  $\Theta^n(\mu_{V_j}, x) \geq 1$  for all  $x \in \text{spt } \mu_{V_j}$  and*

$$\sup_j \mu_{V_j}(K) < \infty, \quad \sup_j \|\underline{H}_j\|_{L^p(\mu_{V_j} \llcorner K)} < \infty$$

for each compact  $K \subset U$ .

*Then there is a Borel regular measure  $\mu$  on  $U$  and a subsequence  $V_{j_\ell}$  with  $\mu_{V_{j_\ell}} \rightarrow \mu$  and  $\text{spt } \mu_{V_{j_\ell}}$  converging to  $\text{spt } \mu$  in the Hausdorff distance sense 7.3.*

**7.5 Remark:** We will show in Chapter 8 (in the Allard compactness theorem) that the limiting measure  $\mu$  in the above statement is in fact the weight measure  $\mu_V$  of a rectifiable varifold  $V = \underline{v}(M, \theta)$  with  $\theta \geq 1$   $\mu_V$ -a.e. and with generalized mean curvature  $\underline{H} \in L^p_{\text{loc}}(\mu_V)$ .

**Proof of Theorem 7.5:** First note that the existence of a subsequence  $\mu_{V_{j_\ell}}$  converging to a limiting measure  $\mu$  is a consequence of the general convergence theorem 5.15, and indeed the inclusion  $K \cap \text{spt } \mu \subset \{x \in U : \text{dist}(x, \text{spt } \mu_{V_{j_\ell}}) < \delta\}$  for all sufficiently large  $\ell$  is a general property of convergent sequences of measures, and easily checked using the definition of  $\text{spt } \mu_{V_{j_\ell}}$  and  $\text{spt } \mu$ .

So only the inclusion  $K \cap \text{spt } \mu_{V_{j_\ell}} \subset \{x \in U : \text{dist}(x, \text{spt } \mu) < \delta\}$  for all sufficiently large  $\ell$  needs to be checked. Supposing the contrary, we would have compact  $K \subset U$  and  $\delta > 0$  and  $z_\ell \in K \cap \text{spt } \mu_{V_{j_{q_\ell}}}$  with  $\text{dist}(z_\ell, \text{spt } \mu) \geq \delta$  and  $q_\ell \geq \ell$ . Since  $K$  is compact  $z_\ell$  has a subsequence (still denoted  $z_\ell$ ) which converges to  $z \in K$  with  $\text{dist}(z, \text{spt } \mu) \geq \delta$ . By convergence of  $\mu_{V_{j_{q_\ell}}}$  to  $\mu$  we have  $\limsup_{\ell \rightarrow \infty} \mu_{V_{j_{q_\ell}}}(B_{\delta/2}(z)) \leq \mu(B_{3\delta/4}(z)) = 0$ , i.e.  $\lim_{\ell \rightarrow \infty} \mu_{V_{j_{q_\ell}}}(B_{\delta/2}(z)) = 0$ . But  $B_{\delta/2}(z) \supset B_{\delta/4}(z_\ell)$  for sufficiently large  $\ell$ , hence, by the monotonicity 4.5,  $\mu_{V_{j_{q_\ell}}}(B_{\delta/2}(z)) \geq \mu_{V_{j_{q_\ell}}}(B_{\delta/4}(z_\ell)) \geq C$  for a fixed positive constant  $C$  because  $z_\ell \in \text{spt } \mu_{V_{q_\ell}}$ .  $\square$

We note the following corollary of the above Theorem:

**7.6 Corollary.** *Suppose  $\theta \geq 1$   $\mu$ -a.e. in  $U$ ,  $\underline{H} \in L^p_{\text{loc}}(\mu)$  in  $U$  for some  $p > n$ . If the approximate tangent space  $T_x V$  (see §1) exists at a given point  $x \in U$ , then  $T_x V$  is a “classical” tangent plane for  $\text{spt } \mu_V$  in the sense that  $\eta_{x,\rho}(\text{spt } \mu_V)$  converges, as  $\rho \downarrow 0$ , locally in the Hausdorff distance sense in  $\mathbb{R}^{n+k}$  to the subspace  $T_x V$ .*

## CHAPTER 4 PROBLEMS

4.1 Let  $M$  be a smooth  $n$ -dimensional minimal surface in  $\mathbb{R}^{n+\ell}$  with  $\overline{M} \setminus M = \emptyset, 0 \in M$  and  $\lim_{\rho \rightarrow \infty} (\omega_n \rho^n)^{-1} \mathcal{H}^n(M \cap B_\rho(0)) = 1$ . Prove that  $M$  is an  $n$ -dimensional subspace of  $\mathbb{R}^{n+\ell}$ .

Hint: Monotonicity identity.

4.2 Suppose  $M$  is a bounded,  $\mathcal{H}^n$ -measurable, countably  $n$ -rectifiable subset of  $\mathbb{R}^{n+\ell}$  with  $\mathcal{H}^n(M \cap K) < \infty$  for each compact  $K \subset \mathbb{R}^{n+\ell}$  and with  $M$  stationary in  $\mathbb{R}^{n+\ell}$  (i.e.  $\int_M \operatorname{div}_M X \, d\mathcal{H}^n = 0$  for each  $C^1$  vector field  $X$  on  $\mathbb{R}^{n+\ell}$  with compact support in  $\mathbb{R}^{n+\ell}$ ). Prove  $\mathcal{H}^n(M) = 0$  (so in particular there are no smooth compact minimal surfaces without boundary in  $\mathbb{R}^{n+\ell}$ ).

Hint: Use monotonicity.

4.3 Let  $U$  be open in  $\mathbb{R}^{n+\ell}$ , let  $V = \underline{v}(M, \theta)$  be a rectifiable  $n$ -varifold which is stationary in  $U$ , let  $\mu_V$  be the weight measure (i.e.  $d\mu_V = \theta d\mathcal{H}^n \llcorner M$ ), and assume  $\theta \geq 1$   $\mu_V$ -a.e.

(i) Prove  $V = \underline{v}(S, \Theta)$ , where  $S = \operatorname{spt} \mu_V$  (closed), and  $\Theta(x) = \Theta^n(\mu_V, x)$ ,  $x \in S$  (i.e. prove that  $\mu_V((S \setminus M) \cup (M \setminus S)) = 0$  and  $\Theta = \theta \mu_V$ -a.e. on  $M \cap S$ ).

(Recall that by definition  $\operatorname{spt} \mu_V = \{x \in U : \mu_V(B_\sigma(x) \cap U) > 0 \, \forall \sigma > 0\}$ .)

(ii) If  $B_\rho(y) \subset U$  and  $(\omega_n \rho^n)^{-1} \mu_V(B_\rho(y)) < 1$ , prove that  $y \notin S$ .

4.4 If  $U, V, S$  are as in 4.3 above, if  $x \in U$ ,  $q \in \{1, 2, \dots\}$ , and if  $V$  has multiplicity  $q$  approximate tangent space at  $x$  (i.e. there is an  $n$ -dimensional subspace  $L \subset \mathbb{R}^{n+\ell}$  such that  $\int_{\eta_{x,\lambda} S} f \theta \circ \eta_{x,\lambda}^{-1} d\mathcal{H}^n \rightarrow q \int_L f d\mathcal{H}^n$  as  $\lambda \downarrow 0 \, \forall f \in C_c^0(U)$ ), prove that  $\eta_{x,\lambda} S \rightarrow L$  locally in the Hausdorff distance sense.

Note:  $\eta_{x,\lambda} S \rightarrow L$  locally in the Hausdorff distance sense means that for each  $R > 0$ ,  $\varepsilon > 0$  there is  $\delta > 0$  with  $\eta_{x,\lambda} S \cap B_R(0) \subset \{x : \operatorname{dist}(x, L) < \varepsilon\}$  and  $L \cap B_R(0) \subset \{x : \operatorname{dist}(x, \eta_{x,\lambda} S) < \varepsilon\}$  for all  $0 < \lambda < \delta$ .



# Chapter 5

## The Allard Regularity Theorem

§ 1	Harmonic Approximation in the Smooth Case .....	128
§ 2	Preliminaries, Lipschitz Approximation .....	130
§ 3	Approximation by Harmonic Functions .....	139
§ 4	The Tilt-Excess Decay Lemma .....	140
§ 5	Main Regularity Theorem .....	145
§ 6	Some Initial Applications of the Allard Theorem .....	150

Here we discuss Allard's ([All72]) regularity theorem, which says roughly that if the generalized mean curvature of a rectifiable  $n$ -varifold  $V = \underline{v}(M, \theta)$  is in  $L^p_{\text{loc}}(\mu_V)$  in  $U$ ,  $p > n$ , if  $\theta \geq 1$   $\mu_V$ -a.e. in  $U$ , if  $\xi \in \text{spt } V \cap U$ , and if  $\omega_n^{-1} \rho^{-n} \mu_V(B_\rho(\xi))$  is sufficiently close to 1 for *some* sufficiently small<sup>1</sup>  $\rho$ , then  $V$  is regular near  $\xi$  in the sense that  $\text{spt } V$  is a  $C^{1,1-n/p}$   $n$ -dimensional embedded submanifold near  $\xi$ .

A key idea of the proof is to show that  $V$  is well-approximated by the graph of a harmonic function near  $\xi$ . We begin in the first section with a motivating discussion, where we consider smooth minimal surfaces with small  $C^1$  norm, and discuss the fact that in such a classical setting harmonic functions do indeed give a very good approximation.

The rest of the chapter is devoted to Allard's theorem, beginning in §2 with a discussion of the fact that a stationary  $n$ -dimensional rectifiable varifold  $V$  in a ball  $B_R(\xi) \subset \mathbb{R}^{n+k}$  which has mass density ratio  $(\omega_n R^n)^{-1} \mu_V(B_R(\xi))$  close to 1 has nice affine approximation properties near every point in the support, and can be very well approximated by a Lipschitz graph with small Lipschitz constant. We in fact do this under the assumption that the generalized mean curvature has small  $L^p$  norm with  $p > n$ .

In §4 we show that the harmonic approximation lemma of §3 can be applied to the

---

<sup>1</sup>Depending on  $\|\underline{H}\|_{L^p(\mu_V)}$

Lipschitz approximation of §2, leading to the “tilt-excess decay” theorem, which is the main step in the proof of the Allard theorem.

The idea of approximating by harmonic functions (in roughly the sense used here) goes back to De Giorgi [DG61] who proved a special case of the above theorem (when  $k = 1$  and when  $V$  corresponds to the reduced boundary of a set of least perimeter—see the previous discussion in §4 of Ch. 3 and the discussion in §5 below). Almgren used analogous approximations in his work [Alm68] for arbitrary  $k \geq 1$ . Reifenberg [Rei60, Rei64] used approximation by harmonic functions in a rather different way in his work on regularity of minimal surfaces.

## 1 Harmonic Approximation in the Smooth Case

Suppose  $M$  is an  $n$ -dimensional  $C^2$  embedded submanifold of  $\mathbb{R}^{n+k}$ . We say that  $M$  is a minimal submanifold if its mean curvature vector  $\underline{H}$  is identically zero. From the discussion in Ch. 2 we have seen that this is exactly equivalent to the volume  $\mathcal{H}^n(M)$  being stationary with respect to compactly supported perturbations of the identity. Thus, in the notation of §6 of Ch. 2,  $M$  is minimal if and only if  $\frac{d}{dt} \mathcal{H}^n(\varphi_t(M))|_{t=0} = 0$ . We showed that this in turn is equivalent to the first variation identity  $\int_M \operatorname{div}_M X \, d\mathcal{H}^n = 0$ .

In the present smooth case we can use the local graphical representations discussed in Remark 4.2(3) of Ch. 2. Thus, modulo an orthogonal transformation of  $\mathbb{R}^{n+k}$  we can locally write  $M$  as a graph of a  $C^2$  function with values in  $\mathbb{R}^k$  over a domain in  $\mathbb{R}^n$ . Thus for each  $\xi \in M$  we can assume there are open sets  $W \subset \mathbb{R}^{n+k}$  and a ball  $B_\rho(\eta) \subset \mathbb{R}^n$  and a  $C^2$  map  $u : B_\rho(\eta) \rightarrow \mathbb{R}^k$  such that  $u(\eta) = \xi$ ,  $Du(\eta) = 0$  and  $\operatorname{graph} u = \{(x, u(x)) : x \in \check{B}_\rho(\eta)\} = M \cap W$ . Then stationarity of  $M$  implies in particular that the area functional

$$\mathcal{A}(u) = \int_{B_\rho(\eta)} J_u \, d\mathcal{H}^n$$

must be stationary with respect to compactly supported perturbations of  $u$  in  $B_\rho(\eta)$ , where  $J_u$  is the Jacobian of the graph map  $x \in B_\rho(\eta) \mapsto (x, u(x)) \in \operatorname{graph} u = M \cap W$ . Thus  $J_u = \sqrt{\det \mathcal{J}}$ , where  $\mathcal{J}(x)$  is the  $n \times n$  matrix  $(D_i(x, u(x)) \cdot D_j(x, u(x)))$ ; i.e., the  $n \times n$  matrix with entry  $(e_i, D_i u(x)) \cdot (e_j, D_j u(x)) = \delta_{ij} + D_i u(x) \cdot D_j u(x)$  in the  $i$ -th row and  $j$ -th column. Thus

$$\mathcal{A}(u) = \int_{B_\rho(\eta)} \sqrt{\det(\delta_{ij} + D_i u \cdot D_j u)} \, d\mathcal{L}^n$$

and for  $|Du| < \varepsilon_0$  (for suitably small  $\varepsilon_0 = \varepsilon_0(n, k) \in (0, 1)$ ) we can use a Taylor series

expansion to give

$$1.1 \quad \mathcal{A}(u) = \int_{B_\rho(\eta)} \left(1 + \frac{1}{2}|Du|^2 + F(Du)\right) d\mathcal{L}^n,$$

where  $F = F(P)$  is a real analytic map of  $n \times k$  matrices  $P = (p_{ij})_{i=1,\dots,n, j=1,\dots,k}$  with  $|P| < \varepsilon_0$  such that

$$1.2 \quad |F(P)| \leq C|P|^4, \quad |D_{p_{ij}}F(P)| \leq C|P|^3, \quad |P| \leq 1,$$

where  $C$  is a fixed constant depending only on  $n, k$ .

Since  $\mathcal{A}(u)$  is stationary with respect to compactly supported perturbations of  $u$  we have

$$\frac{d}{dt}\mathcal{A}(u + t\zeta)|_{t=0} = 0, \quad \zeta = (\zeta_1, \dots, \zeta_k) \in C_0^1(B_\rho(\eta), \mathbb{R}^k),$$

where  $C_0^1(B_\rho(\eta), \mathbb{R}^k)$  denotes the  $C^1$  maps  $\zeta : B_\rho(\eta) \rightarrow \mathbb{R}^k$  with  $\zeta = 0$  on  $\partial B_\rho(\eta)$ .

In view of 1.1, if  $|Du| < \varepsilon_0$  this takes the form

$$1.3 \quad \int_{B_\rho(\eta)} \sum_{i=1}^n D_i u \cdot D_i \zeta d\mathcal{L}^n = \int_{B_\rho(\eta)} \sum_{i=1}^n \sum_{j=1}^k A_{ij}(Du) D_i \zeta_j,$$

for all  $\zeta \in C_0^1(B_\rho(\eta), \mathbb{R}^k)$ , where  $A_{ij}(P) = D_{p_{ij}}F(P)$ , so  $|A_{ij}(P)| \leq C|P|^3$ . Integrating by parts, we get

$$\Delta u = \sum_{i,j=1}^n a_{ij}(Du) D_i D_j u, \quad a_{ij}(Du) = O(|Du|^2).$$

It is therefore reasonable, so long as  $|Du|$  is small, to expect that  $u$  is well approximated by a harmonic function. Indeed let us check this rigorously: Assume  $|Du| < \varepsilon_0$  ( $\varepsilon_0$  as above), and let  $v$  be the harmonic function on the ball  $B_\rho(\eta)$  with  $v = u$  on  $\partial B_\rho(\eta)$ —it is standard that such a harmonic function  $v$  exists and it is  $C^1$  on  $B_\rho(\eta)$  and  $C^\infty(\check{B}_\rho(\eta))$ . Multiplying the equation  $\Delta v = 0$  by  $\zeta$  and integrating by parts over the ball  $B_\rho(\eta)$ , we obtain

$$1.4 \quad \int_{B_\rho(\eta)} \sum_{i=1}^n D_i v \cdot D_i \zeta d\mathcal{L}^n = 0, \quad \zeta \in C_0^1(B_\rho(\eta)).$$

Taking the difference between 1.3 and 1.4, we see then that

$$\int_{B_\rho(\eta)} \sum_{i=1}^n D_i (u-v) \cdot D_i \zeta d\mathcal{L}^n = \int_{B_\rho(\eta)} \sum_{i=1}^n \sum_{j=1}^k A_{ij}(Du) D_i \zeta_j, \quad \zeta \in C_0^1(B_\rho(\eta), \mathbb{R}).$$

In this identity we take  $\zeta = u - v$ , so that

$$\int_{B_\rho(\eta)} |D(u-v)|^2 d\mathcal{L}^n = \int_{B_\rho(\eta)} \sum_{i=1}^n \sum_{j=1}^k A_{ij}(Du) D_i (u_j - v_j),$$

and using the Cauchy-Schwarz inequality  $ab \leq \frac{1}{2}a^2 + \frac{1}{2}b^2$  on the right side we get finally

$$\int_{B_\rho(\eta)} |D(u-v)|^2 d\mathcal{L}^n \leq \frac{1}{2} \int_{B_\rho(\eta)} \sum_{i,j} (A_{ij}(Du))^2 + \frac{1}{2} \int |D(u-v)|^2,$$

so

$$\int_{B_\rho(\eta)} |D(u-v)|^2 d\mathcal{L}^n \leq \int_{B_\rho(\eta)} \sum_{i,j} (A_{ij}(Du))^2.$$

That is, since  $|\sum_{ij} (A_{ij}(P))^2| \leq C|P|^6$  for  $|P| < \varepsilon_0$ , we obtain

$$1.5 \quad \int_{B_\rho(\eta)} |D(u-v)|^2 d\mathcal{L}^n \leq C \int_{B_\rho(\eta)} |Du|^6.$$

This shows that indeed  $v$  is a very good approximation of  $u$  for  $|Du|$  small: For example if  $\sup_{B_\rho(\eta)} |Du| = \varepsilon < \varepsilon_0$ , then 1.5 shows

$$\int_{B_\rho(\eta)} |D(u-v)|^2 d\mathcal{L}^n \leq C\varepsilon^4 \int_{B_\rho(\eta)} |Du|^2 d\mathcal{L}^n,$$

where (as in 1.2)  $C$  is a fixed constant depending only on  $n, k$ , so that  $\int_{B_\rho(\eta)} |D(u-v)|^2$  is much smaller than  $\int_{B_\rho(\eta)} |Du|^2$  for  $\varepsilon$  small.

So there is good motivation to think that harmonic approximation could be relevant in the study of the regularity of stationary rectifiable varifolds; indeed, as mentioned in the introduction to this chapter, we will show that such approximations are appropriate even in the more general context of rectifiable varifolds with generalized mean curvature in  $L^p$ ,  $p > n$ .

## 2 Preliminaries, Lipschitz Approximation

In this section  $U$  is an open subset of  $\mathbb{R}^{n+k}$ ,  $V = \underline{v}(M, \theta)$  is a rectifiable  $n$ -varifold with generalized mean curvature  $\underline{H}$  in  $U$  (as in Definition 3.15 of Ch. 4).

A key quantity which will appear in the computations to follow is the *tilt excess*  $E(\xi, \sigma, T)$  of  $V$  over a ball  $B_\sigma(\xi) \subset U$  relative to a given  $n$ -dimensional subspace  $T \subset \mathbb{R}^{n+k}$ ; this is defined by

$$2.1 \quad E(\xi, \sigma, T) = \sigma^{-n} \int_{B_\sigma(\xi)} |p_{T_x} M - p_T|^2 d\mu_V(x)^2.$$

Notice that this could be roughly described as “the mean square deviation of  $T_x M$  away from  $T$  in  $B_\sigma(\xi)$ .”

---

<sup>2</sup> $|p_{T_x} M - p|^2$  denotes the inner product norm trace  $(p_{T_x} - p)^2$ ; this differs from  $\|p_{T_x} M - p\|^2$  by at most a constant factor depending on  $n+k$ —see Remark 2.2 below.

**2.2 Remark (Operator norm v. inner product norm):** In the above definition 2.1 of tilt excess we use the inner product norm for  $p_{T_x} - p_T$ , but we could equivalently use the operator norm: If  $L : \mathbb{R}^P \rightarrow \mathbb{R}^Q$  is linear with matrix  $\ell = (\ell_i^j)$  (so that  $L(x) = \sum_{j=1}^Q \sum_{i=1}^P \ell_i^j x^i e_j$ ) then the operator norm is  $\|L\| = \sup_{|x|=1} |L(x)|$ , whereas the inner product norm is  $|L| = \sqrt{\sum_{i,j} (\ell_i^j)^2}$ . Observe  $|L(x)|^2 = x^T \ell^T \ell x$  and  $\ell^T \ell$  is a symmetric positive semi-definite  $P \times P$  matrix with non-negative eigenvalues  $0 \leq \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_P$  and  $|L|^2 = \text{trace } \ell^T \ell = \sum_{j=1}^P \lambda_j$ , while  $\|L\|^2 = \lambda_P = \max\{\lambda_1, \dots, \lambda_P\}$ , so

$$P^{-1}|L|^2 \leq \|L\|^2 \leq |L|^2.$$

In particular  $(n+k)^{-1} \int_{B_r(\xi)} |p_{T_x M} - p_{\mathbb{R}^n \times \{0\}}|^2 \leq \int_{B_\sigma(\xi)} \|p_{T_x M} - p_{\mathbb{R}^n \times \{0\}}\|^2 \leq \int_{B_\sigma(\xi)} |p_{T_x M} - p_{\mathbb{R}^n \times \{0\}}|^2$ —i.e.  $\int_{B_\sigma(\xi)} \|p_{T_x M} - p_{\mathbb{R}^n \times \{0\}}\|^2$  and  $\int_{B_\sigma(\xi)} |p_{T_x M} - p_{\mathbb{R}^n \times \{0\}}|^2$  differ by at most fixed factor depending on  $n+k$ .

If  $T = \mathbb{R}^n \times \{0\}$  then, in terms of the  $(n+k) \times (n+k)$  matrices  $(e^{ij})$  and  $(\varepsilon^{ij})$  for the orthogonal projections  $p_{T_x M}$  and  $p_T$  respectively,  $|p_{T_x M} - p_{\mathbb{R}^n \times \{0\}}|^2$  is just  $\sum_{i,j} (e^{ij} - \varepsilon^{ij})^2 = \sum_{i,j} ((e^{ij})^2 + (\varepsilon^{ij})^2 - 2e^{ij}\varepsilon^{ij}) = 2(n - \sum_{j=1}^n e^{jj}) = 2\sum_{j=n+1}^{n+k} e^{jj} = 2\sum_{j=1}^k |\nabla^M x^{n+j}|^2$ , where we used the facts that  $(e^{ij})^2 = (e^{ij})$  and  $\text{trace}(e^{ij}) = n$ . Thus

$$2.3 \quad \frac{1}{2} |p_{T_x M} - p_T|^2 = \sum_{j=n+1}^{n+k} e^{jj} = \sum_{j=1}^k |\nabla^M x^{n+j}|^2,$$

so, still assuming  $T = \mathbb{R}^n \times \{0\}$ ,

$$2.4 \quad E(\xi, \rho, T) = 2\sigma^{-n} \int_{B_\sigma(\xi)} \sum_{j=1}^k |\nabla^M x^{n+j}|^2 d\mu_V$$

( $\nabla^M$  = gradient operator on  $M$  as defined in §2 of Ch.3).

We begin with the following lemma relating tilt-excess and  $L^2$  distance from the relevant affine plane.

**2.5 Lemma.** *Suppose  $B_\rho(y) \subset U$ . Then for any  $n$ -dimensional subspace  $T \subset \mathbb{R}^{n+k}$  and any  $\gamma \in (0, 1)$  we have*

$$\begin{aligned} & \rho^{-n} \int_{B_{\gamma\rho}(y)} |p_{T_x M} - p_T|^2 d\mu_V(x) \\ & \leq C\rho^{-n} \int_{B_\rho(y)} \left( \frac{\text{dist}(x, \xi + T)}{\rho} \right)^2 d\mu + C\rho^{2-n} \int_{B_\rho(y)} |\underline{H}|^2 d\mu_V, \end{aligned}$$

where  $C = C(n, \gamma)$ .

**Proof of 2.5:** It evidently suffices to prove the result with  $y = 0$  and  $T = \mathbb{R}^n \times \{0\}$ . The proof simply involves making a suitable choice of  $X$  in the first variation formula of 3.15

( $\ddagger$ ) of Ch. 4. In fact we take

$$X = \zeta^2(x)x', \quad x' = (0, x^{n+1}, \dots, x^{n+k})$$

for  $x = (x^1, \dots, x^{n+k}) \in U$ , where  $\zeta \in C_c^1(U)$  with  $\zeta \geq 0$  will be chosen below.

By the definition of  $\operatorname{div}_M$  (see §2 of Ch. 3) we have

$$\operatorname{div}_M x' = \sum_{i=n+1}^{n+k} e^{ii}, \quad \mu\text{-a.e. } x \in M,$$

where  $(e^{ij})$  is the matrix of the projection  $p_{T_x M}$  (relative to the standard orthonormal basis for  $\mathbb{R}^{n+k}$ ), and where, here and subsequently, we write  $\mu = \mu_V$ . Thus by the definition 3.15 of Ch. 4 of  $\underline{H}$  we have

$$(1) \quad \int \sigma \zeta^2 d\mu = \int (-2\zeta \sum_{i=n+1}^{n+k} \sum_{j=1}^{n+k} x^i e^{ij} D_j \zeta - \zeta^2 x' \cdot \underline{H}) d\mu,$$

with

$$(2) \quad \sigma = \sum_{i=n+1}^{n+k} e^{ii} = \frac{1}{2} \sum_{i,j=1}^{n+k} (e^{ij} - \varepsilon^{ij})^2 = \frac{1}{2} |p_{T_x M} - p_{\mathbb{R}^n \times \{0\}}|^2,$$

by 2.3, where  $(\varepsilon^{ij}) = \text{matrix of } p_{\mathbb{R}^n \times \{0\}}$ . Also observe that  $\varepsilon^{ij} = 0$  if  $i > n$ , so (1) can be written

$$(3) \quad \int \sigma \zeta^2 d\mu = \int (-2\zeta \sum_{i=n+1}^{n+k} \sum_{j=1}^{n+k} x^i (e^{ij} - \varepsilon^{ij}) D_j \zeta - \zeta^2 (0, x') \cdot \underline{H}) d\mu,$$

so

$$\int \sigma \zeta^2 d\mu \leq \int (2\sqrt{2\sigma} |x'| |\nabla \zeta| \zeta + |x'| |\underline{H}| \zeta^2) d\mu.$$

Hence (using  $ab \leq \frac{1}{2}a^2 + \frac{1}{2}b^2$ )

$$\int \sigma \zeta^2 d\mu \leq 16 \int (|x'|^2 |\nabla \zeta|^2 + |x'| |\underline{H}| \zeta^2) d\mu.$$

The lemma now follows by choosing  $\zeta \equiv 1$  in  $B_{\gamma\rho}(0)$ ,  $\zeta \equiv 0$  outside  $B_\rho(0)$  and  $|\nabla \zeta| \leq 2/((1-\gamma)\rho)$ , and then noting that  $|x'| |\underline{H}| = (\rho^{-1}|x'|)(|\underline{H}|\rho) \leq \frac{1}{2}\rho^{-2}|x'|^2 + \frac{1}{2}(|\underline{H}|\rho)^2$ .  $\square$

For the remainder of this section we continue to assume that  $V = \underline{v}(M, \theta)$  has generalized mean curvature  $\underline{H}$  and now we additionally assume, with  $\delta \in (0, \frac{1}{4}]$  a constant to be specified below and  $\mu = \mu_V = \mathcal{H}^n \llcorner \theta$ , the following:

$$2.6 \quad \left\{ \begin{array}{l} 1 \leq \theta \text{ } \mu\text{-a.e.}, \quad 0 \in \operatorname{spt} V, \quad B_\rho(0) \subset U \\ \omega_n^{-1} \rho^{-n} \mu(B_\rho(0)) \leq 1 + \delta, \quad (\rho^{p-n} \int_{B_\rho(0)} |\underline{H}|^p d\mu)^{1/p} \leq \delta. \end{array} \right.$$

Notice that then by 3.13 of Ch. 4 we then have the “canonical representative”  $(\text{spt } \mu_V, \Theta^n(\mu_V, \cdot))$  for  $V$ , so we can, and we shall, assume  $\theta(y) = \Theta^n(\mu_V, y)$  for all  $y \in B_1(0)$ , and  $\Theta^n(\mu, x)$  is an upper semi-continuous function of  $x$  by 4.7 of Ch. 4, so

$$2.7 \quad \theta(x) = \Theta^n(\mu, x) \geq 1 \text{ at every point } x \in \text{spt } \mu \cap \check{B}_\rho(0).$$

Also, by the monotonicity 4.5 of Ch. 4 we have, subject to 2.6, that  $y \in \text{spt } V \cap B_{2\delta\rho}(0)$  and  $\sigma \in (0, (1 - 2\delta)\rho] \Rightarrow 1 - C\delta \leq (\omega_n \sigma^n)^{-1} \mu(B_\sigma(y)) \leq (1 + C\delta)(\omega_n(1 - 2\delta)^n \rho^n)^{-1} \mu(B_{(1-2\delta)\rho}(y)) \leq 1 + C\delta$ , because  $B_{(1-2\delta)\rho}(y) \subset B_\rho(0)$ . Thus for  $\delta \leq \delta_0$ , with  $\delta_0 = \delta_0(n, k, p)$  sufficiently small, we have

$$2.8 \quad \frac{1}{2} \leq 1 - C\delta \leq \frac{\mu(B_\sigma(y))}{\omega_n \sigma^n} \leq 1 + C\delta \leq 2, \quad \sigma \in (0, (1 - 2\delta)\rho], \quad y \in \text{spt } \mu \cap B_{2\delta\rho}(0),$$

where  $C = C(n, k, p)$ . In particular, letting  $\sigma \downarrow 0$ , we have

$$2.9 \quad \theta(y) \leq 1 + C\delta, \quad y \in \text{spt } \mu \cap B_{2\delta\rho}(0).$$

Also, by the monotonicity identity 4.3 of Ch. 4, we have, assuming 2.6,

$$2.10 \quad \int_{B_{(1-2\delta)\rho}(y)} |x - y|^{-n} \left| \left( \frac{x - y}{|x - y|} \right)^\perp \right|^2 d\mu \leq C\delta, \quad y \in \text{spt } \mu \cap B_{2\delta\rho}(0),$$

with  $C = C(n, k, p)$ .

We now establish a lemma which guarantees local affine approximations of the support at all points of  $\text{spt } \mu_V$  in the ball  $B_{2\delta\rho}(0)$  and at all radii  $\sigma \leq 4\delta\rho$ .

**2.11 Lemma (Affine Approximation Lemma.)** *If  $\delta \in (0, \frac{1}{16}]$  and 2.6 holds, then, for each  $y \in \text{spt } \mu_V \cap B_{2\delta\rho}(0)$  and each  $\sigma \in (0, 4\delta\rho]$ ,*

$$(\ddagger) \quad \sup_{x \in \text{spt } \mu_V \cap B_\sigma(y)} \text{dist}(x, y + T(y, \sigma)) \leq C\delta^{\frac{1}{2n+2}} \sigma, \quad C = C(n, k, p).$$

**Proof:** Take any fixed  $\sigma \in (0, 4\delta\rho]$  and  $y \in \text{spt } V \cap B_{2\delta\rho}(0)$ , and suppose for convenience of notation (by changing scale and translating the origin) that  $\sigma = 1$  and  $y = 0$ , so now, since  $\delta \leq \frac{1}{16}$  and hence  $(1 - 6\delta)\rho/(4\delta\rho) > 2$ , 2.8 in this rescaled setting ensures that

$$(1) \quad \frac{1}{2} \leq \frac{\mu(B_\sigma(y))}{\omega_n \sigma^n} \leq 2, \quad \forall \sigma \in (0, 2], \quad y \in \text{spt } \mu \cap B_1(0),$$

where, here and subsequently,  $\mu = \mu_V$ . Also, 2.10 now guarantees

$$(2) \quad \int_{B_2(y)} |p_{(T_x M)^\perp}(x - y)|^2 d\mu \leq \int_{B_2(y)} |p_{(T_x M)^\perp}(x - y)|^2 |x - y|^{-n-2} d\mu \leq C\delta$$

for  $y \in \text{spt } V \cap B_1(0)$ . Next take  $\alpha \in (0, 1)$  (to be chosen shortly, but for the moment arbitrary). Recall the general principle that if  $K$  is compact and  $\eta > 0$  then any maximal

pairwise disjoint collection  $B_{\eta/2}(y_j)$  of closed balls with  $y_j \in K$  will automatically have the property that  $K \subset \cup_j B_{\eta}(y_j)$ . Using this with  $\eta = \delta^\alpha$  we have pairwise disjoint balls  $B_{\delta^\alpha/2}(y_1), \dots, B_{\delta^\alpha/2}(y_N)$  with  $y_j \in \text{spt } \mu \cap B_1(0)$  such that

$$(3) \quad \text{spt } \mu \cap B_1(0) \subset \cup_{j=1}^N B_{\delta^\alpha/2}(y_j).$$

Notice that then, by (1),

$$(4) \quad \omega_n 2^{-n-1} \delta^{\alpha n} \leq \mu(B_{\delta^\alpha/2}(y_j)), \quad j = 1, \dots, N,$$

and hence, using (1) with  $\sigma = 2$ ,

$$\omega_n 2^{-n-1} N \delta^{\alpha n} \leq \sum_{j=1}^N \mu(B_{\delta^\alpha/2}(y_j)) = \mu(\cup_{j=1}^N B_{\delta^\alpha/2}(y_j)) \leq \mu(B_2(0)) \leq 2^{n+1} \omega_n.$$

Thus  $N \leq 4^{n+1} \delta^{-\alpha n}$ , and so, by using (2) with  $y = y_j$  and noting that  $B_2(y_j) \supset B_1(0)$  for each  $j$ , we have

$$\int_{B_1(0)} \sum_{j=1}^N |p_{T_x M^\perp}(x - y_j)|^2 d\mu \leq CN\delta = C\delta^{1-\alpha n}.$$

Thus for any given  $k \geq 1$  we have

$$(5) \quad \sum_{j=1}^N |p_{T_x M^\perp}(x - y_j)|^2 \leq Ck\delta^{1-\alpha n},$$

except possibly for a set of  $x \in B_1(0) \cap \text{spt } \mu$  of  $\mu$ -measure  $\leq 1/k$ . Since  $\mu(B_{\delta^\alpha}(0)) \geq C^{-1}\delta^{\alpha n}$  by 2.8, we can select  $k = C\delta^{-\alpha n}$ , thus ensuring that (5) holds for some  $x_0 \in \text{spt } \mu \cap B_{\delta^\alpha}(0)$ . So we have shown there is  $x_0 \in \text{spt } \mu \cap B_{\delta^\alpha}(0)$  with

$$(6) \quad \sum_{j=1}^N |p_{(T_{x_0} M)^\perp}(x_0 - y_j)|^2 \leq C\delta^{1-2\alpha n},$$

and hence

$$|p_{(T_{x_0} M)^\perp}(y_j - x_0)| \leq C\delta^{\frac{1}{2}-\alpha n}, \quad j = 1, \dots, N.$$

Since  $|x_0| < \delta^\alpha$ , we then have

$$(7) \quad |p_{(T_{x_0} M)^\perp} y_j| \leq C(\delta^{\frac{1}{2}-\alpha n} + \delta^\alpha), \quad j = 1, \dots, N.$$

Then, selecting  $\alpha$  such that  $\frac{1}{2} - \alpha n = \alpha$  (i.e.  $\alpha = \frac{1}{2n+2}$ ), we have shown that all the points  $y_1, \dots, y_N$  are in the  $C\delta^{1/(2n+2)}$  neighborhood of the subspace  $T_0 = T_{x_0} M$ , and hence by (3) we have

$$\text{dist}(y, T_0) \leq C\delta^{1/(2n+2)} \quad \forall y \in \text{spt } \mu \cap B_1(0),$$

so the inequality ( $\ddagger$ ) is proved with  $T = T_0 = T_{x_0} M$ .  $\square$

**2.12 Remark:** Note that if  $y \in B_{2\delta}(0)$  and  $\sigma \in (0, 4\delta]$  we can use the Hölder inequality and  $\mu(B_\sigma(y)) \leq 2\omega_n \sigma^n$  (by 2.6) to estimate

$$\rho^{2-n} \int_{B_\sigma(y)} |\underline{H}|^2 d\mu \leq C(\rho^{p-n} \int_{B_\sigma(y)} |\underline{H}|^p d\mu)^{2/p},$$

and then choose  $T = T(y, \sigma)$  (notation as in 2.11) in the conclusion of 2.5 to give

$$\begin{aligned} (\ddagger) \quad & \sigma^{-n} \int_{B_{\gamma\sigma}(y)} |p_{T_x M} - p_T|^2 d\mu(x) \\ & \leq C \sup_{x \in \text{spt } \mu_V \cap B_\sigma(y)} \left( \frac{\text{dist}(x, y + T)}{\sigma} \right)^2 + C (\sigma^{p-n} \int_{B_\sigma(y)} |\underline{H}|^p d\mu)^{2/p} \leq C \delta^{\frac{1}{n+1}} \end{aligned}$$

with  $C = C(n, k, p, \gamma)$ .

An important corollary of the above Lemma 2.11 is the following Lipschitz Approximation Theorem, which provides a key step in proving the Allard theorem. We assume in this (without loss of generality since we can rescale and rotate coordinates) that the hypotheses 2.6 hold with  $\rho = 1$  and that the subspace  $T(0, 4\delta)$ , which, according to Lemma 2.11 with  $\rho = 1$ , provides the affine approximation for  $\text{spt } \mu_V$  in the ball  $B_{4\delta}(0)$ , is just  $\mathbb{R}^n \times \{0\}$ . Thus we assume

$$(\ddagger\ddagger) \quad T(0, 4\delta) = \mathbb{R}^n \times \{0\}.$$

**2.13 Lemma (Lipschitz Approximation Theorem.)** *Let  $L \in (0, 1]$  be given. There is  $\beta = \beta(n, k, p) \in (0, \frac{1}{16}]$  such that if  $0 < \delta \leq (\beta L)^{2n+2}$ , if  $\rho = 1$ , if 2.6 holds, and if  $(\ddagger\ddagger)$  above holds, then there is a Lipschitz  $f : B_\delta^n(0) \rightarrow \mathbb{R}^k$  with*

$$\text{Lip } f \leq L, \quad \sup |f| \leq C \delta^{\frac{1}{2n+2}}$$

$$\begin{aligned} & \delta^{-n} (\mu_V(B_\delta(0) \cap (\text{spt } \mu_V \setminus \text{graph } f)) + \mathcal{H}^n(B_\delta(0) \cap (\text{graph } f \setminus \text{spt } \mu_V))) \\ & \leq C L^{-2} \delta^{-n} \int_{B_{3\delta}(0)} |p_{T_x M} - p_{\mathbb{R}^n \times \{0\}}|^2 d\mu_V \leq C L^{-2} \delta^{\frac{1}{n+1}}, \quad C = C(n, k, p). \end{aligned}$$

**Proof:** Let  $\beta \in (0, \frac{1}{2}]$  be for the moment arbitrary, but which we will choose eventually to depend only on  $n, k, p$ . Assume 2.6 holds, where for the moment  $\delta \in (0, \frac{1}{16}]$  is also arbitrary and let  $T_0 = T(0, 4\delta)$ , so  $T_0 = \mathbb{R}^n \times \{0\}$  in accordance with  $(\ddagger\ddagger)$  above. Throughout the proof  $C$  denotes any constant depending only on  $n, k, p$ , and we let  $\mu = \mu_V$ .

Let

$$G = \left\{ y \in \text{spt } \mu \cap B_{2\delta}(0) : \sup_{\sigma \in (0, \delta]} \sigma^{-n} \int_{B_\sigma(y)} |p_{T_x M} - p_{T_0}|^2 d\mu \leq \beta^2 L^2 \right\}.$$

Thus  $y \in \text{spt } \mu \cap B_{2\delta}(0) \setminus G \Rightarrow \exists \sigma \in (0, \delta]$  with

$$(1) \quad \beta^2 L^2 \sigma^n < \int_{B_\sigma(y)} |p_{T_x M} - p_{T_0}|^2 d\mu.$$

By the five-times covering lemma we can pick pairwise disjoint balls  $B_{\sigma_j}(y_j)$  such that (1) holds with  $\sigma = \sigma_j \in (0, \beta]$  and  $y = y_j \in \text{spt } \mu \cap B_{2\delta}(0) \setminus G$ , and such that

$$\text{spt } \mu \cap B_{2\delta}(0) \setminus G \subset \cup_j B_{5\sigma_j}(y_j).$$

Thus using (1) with  $\sigma = \sigma_j$ ,  $y = y_j$  and summing over  $j$  we obtain

$$\begin{aligned} \beta^2 L^2 \mu(B_{2\delta}(0) \setminus G) &\leq \beta^2 L^2 \sum_j \mu(B_{5\sigma_j}(y_j)) \leq C \beta^2 L^2 \sum_j \sigma_j^n \\ &\leq C \int_{\cup_j B_{\sigma_j}(y_j)} |p_{T_x M} - p_{T_0}|^2 d\mu \leq C \int_{B_{3\delta}(0)} |p_{T_x M} - p_{T_0}|^2 d\mu. \end{aligned}$$

Thus

$$(2) \quad \mu(B_{2\delta}(0) \setminus G) \leq C \beta^{-2} L^{-2} \int_{B_{3\delta}(0)} |p_{T_x M} - p_{T_0}|^2 d\mu.$$

We now claim that  $G$  is contained in the graph of a Lipschitz function. To check this, let  $y_1, y_2$  be distinct points of  $G$  and let  $\sigma = |y_1 - y_2|$ , so  $\sigma \leq 4\delta$ . Observe that

$$\begin{aligned} \sigma^{-n} \int_{B_{\sigma/4}(y_1)} |p_{T_x M} - p_{T_0}|^2 &\leq 4^n \beta^2 L^2, & (\text{because } y_1 \in G), \\ \sigma^{-n} \int_{B_{\sigma/4}(y_1)} |p_{T_x M} - p_{T(y_1, \sigma)}|^2 &\leq C \delta^{\frac{1}{n+1}} \leq C \beta^2 L^2 & (\text{by 2.12}(\frac{1}{2})). \end{aligned}$$

Since  $|p_{T_0} - p_{T(y_1, \sigma)}|^2 \leq 2|p_{T_x M} - p_{T_0}|^2 + 2|p_{T_x M} - p_{T(y_1, \sigma)}|^2$  and  $\mu(B_{\sigma/4}(y_1)) \geq \frac{1}{2} \omega_n (\sigma/4)^n$  by 2.8, we now have

$$(3) \quad |p_{T_0} - p_{T(y_1, \sigma)}| \leq C \beta L,$$

and so

$$\begin{aligned} (4) \quad |p_{T_0^\perp}(y_1 - y_2)| &= |(p_{T(y_1, \sigma)^\perp} + (p_{T_0^\perp} - p_{T(y_1, \sigma)^\perp}))(y_1 - y_2)| \\ &\leq \text{dist}(y_2, y_1 + T(y_1, \sigma)) + |p_{T_0} - p_{T(y_1, \sigma)}| \sigma \leq C \beta L \sigma, \end{aligned}$$

where at the last step we again used the Affine Approximation Lemma 2.11. Thus

$$|Q(y_2 - y_1)| \leq C \beta L |y_1 - y_2| \leq C \beta L (|Q(y_1 - y_2)| + |P(y_1 - y_2)|),$$

where  $P, Q$  denote the projections of  $y = (y^1, \dots, y^{n+k})$  onto its first  $n$  and last  $k$  coordinates respectively. Assuming  $C\beta \leq \frac{1}{2}$  we then have

$$|Q(y_1) - Q(y_2)| \leq C \beta L |P(y_1) - P(y_2)|.$$

In view of the arbitrariness of  $y_1, y_2 \in G$  this says that  $G$  is contained in the graph of a Lipschitz function  $f : G \cap B_\delta(0) \rightarrow \mathbb{R}^k$  with Lipschitz constant  $\leq C \beta L$ , provided we eventually choose  $\beta = \beta(n, k, p)$  to satisfy the above restriction  $C\beta \leq \frac{1}{2}$ . Also, by 2.11,  $\sup |f| \leq C \delta^{\frac{1}{2n+2}} \delta$ , so, by the Lipschitz Extension Theorem 1.2 of Ch.2,  $f$  extends to give a Lipschitz  $\tilde{f} : \mathbb{R}^n \rightarrow \mathbb{R}^k$  (henceforth denoted  $f$ ) with

$$(5) \quad G \cap B_{2\delta}(0) \subset \text{graph } f, \text{ with } \text{Lip } f \leq C \beta L \text{ and } \sup_{\mathbb{R}^n} |f| \leq C \beta \delta^{\frac{1}{2n+2}} \delta.$$

Thus we get  $\text{Lip } f \leq L$  as required by choosing  $\beta = \beta(n, k, p)$  such that  $C\beta \leq 1$ . Also, by (2), with  $F = \text{graph } f$  we have

$$(6) \quad \delta^{-n} \mu(B_{2\delta}(0) \setminus F) \leq C L^{-2} \delta^{-n} \int_{B_{3\delta}(0)} |p_{T_x M} - p_{T_0}|^2 d\mu \leq C L^{-2} \delta^{\frac{1}{n+1}}.$$

It thus remains only to prove

$$(7) \quad \mathcal{H}^n(B_\delta(0) \cap F \setminus \text{spt } \mu) \leq CL^{-2} \int_{B_{3\delta}(0)} |p_{T_x M} - p_{T_0}|^2 d\mu.$$

To check this, take any  $\eta \in B_\delta(0) \cap F \setminus \text{spt } \mu$  and let

$$\sigma = \frac{3}{2} \text{dist}(\eta, \text{spt } \mu) \ (\leq \frac{3}{2}\delta \text{ because } 0 \in \text{spt } \mu \text{ and } \eta \in B_\delta(0)).$$

So  $\check{B}_{2\sigma/3}(\eta) \cap \text{spt } \mu = \emptyset$  and  $B_{2\sigma/3}(\eta) \cap \text{spt } \mu \neq \emptyset$ , and the monotonicity identity 4.3 of Ch. 4 implies

$$(8) \quad \begin{aligned} \sigma^{-n} \mu(B_\sigma(\eta)) &= \sigma^{-n} \mu(B_\sigma(\eta)) - (\sigma/2)^{-n} \mu(B_{\sigma/2}(\eta)) \\ &\leq C \int_{B_\sigma(\eta) \setminus B_{\sigma/2}(\eta)} |x - \eta|^{-n} |p_{(T_x M)^\perp}(\frac{x-\eta}{|x-\eta|})|^2 d\mu + C\delta \\ &\leq C\sigma^{-n-2} \int_{B_\sigma(\eta)} |p_{(T_x M)^\perp}(x - \eta)|^2 d\mu + C\delta. \end{aligned}$$

Now  $\text{spt } \mu \cap B_{2\sigma/3}(\eta) \neq \emptyset$  so 2.8 implies

$$(9) \quad \mu(B_\sigma(\eta)) \geq 3^{-n-1} \omega_n \sigma^n,$$

and also  $C\delta \leq C(\beta L)^{2n+2} \leq C\beta^{2n+2}$ , hence, for small enough  $\beta = \beta(n, k, p)$ , (8) gives

$$(10) \quad \begin{aligned} \sigma^n &\leq C \int_{B_\sigma(\eta)} |p_{(T_x M)^\perp}(\frac{x-\eta}{\sigma})|^2 d\mu \\ &\leq C \left( \int_{B_\sigma(\eta)} |p_{T_0^\perp}(\frac{x-\eta}{\sigma})|^2 d\mu + \int_{B_\sigma(\eta)} |p_{T_x M} - p_{T_0}|^2 d\mu \right) \end{aligned}$$

for some  $\sigma \in (0, \frac{3}{2}\delta]$ . On the other hand if  $\sigma \in [\delta, \frac{3}{2}\delta]$  then, by 2.12 (§) with  $\sigma = 4\delta$ ,  $\gamma = \frac{3}{4}$  and  $y = 0$ , and by affine approximation 2.11 with  $\sigma = 4\delta$  and  $y = 0$ , the above inequality gives  $\sigma^n \leq C\delta^{\frac{1}{2n+2}} \sigma^n \leq C\beta L \sigma^n \leq C\beta \sigma^n$ , which is impossible assuming  $C\beta \leq \frac{1}{2}$ . Thus in fact, assuming we do so choose  $\beta = \beta(n, k, p)$ , (10) must hold for some  $\sigma \in (0, \delta]$  rather than  $\sigma \in (0, \frac{3}{2}\delta]$ . Also

$$(11) \quad \begin{aligned} &\int_{B_\sigma(\eta)} |p_{T_0^\perp}(\frac{x-\eta}{\sigma})|^2 d\mu + \int_{B_\sigma(\eta)} |p_{T_x M} - p_{T_0}|^2 d\mu \\ &\leq \int_{B_\sigma(\eta) \cap F} |p_{T_0^\perp}(\frac{x-\eta}{\sigma})|^2 d\mu + C\mu(B_\sigma(\eta) \setminus F) + \int_{B_\sigma(\eta)} |p_{T_x M} - p_{T_0}|^2 d\mu. \end{aligned}$$

Since  $|p_{T_0^\perp}(\frac{x-y}{\sigma})| \leq C\beta$  for  $x, y \in F \cap B_\sigma(\eta)$  (because  $\text{Lip } f \leq C\beta L \leq C\beta$ ), and  $\mu(B_\sigma(\eta)) \leq 2\omega_n \sigma^n$  by 2.8, the inequalities (10) and (11) imply

$$\sigma^n \leq C(\beta \sigma^n + \mu(B_\sigma(\eta) \setminus F) + \int_{B_\sigma(\eta)} |p_{T_x M} - p_{T_0}|^2 d\mu).$$

With  $\beta$  chosen appropriately (depending only on  $n, k, p$ ), we can arrange that  $C\beta \leq \frac{1}{2}$ ,

and hence we have proved

$$(12) \quad \sigma^n \leq C(\mu(B_\sigma(\eta) \setminus F) + \int_{B_\sigma(\eta)} |p_{T_x M} - p_{T_0}|^2 d\mu)$$

for some  $\sigma \in (0, \delta]$ .

Now observe that the collection of such balls  $B_\sigma(\eta)$  by definition cover all of  $B_\delta(0) \cap F \setminus \text{spt } \mu$ , so by the 5-times covering lemma we can find a pairwise disjoint collection  $B_{\sigma_j}(\eta_j)$  of such balls with

$$(13) \quad \sigma_j^n \leq C(\mu(B_{\sigma_j}(\eta_j) \setminus F) + \int_{B_{\sigma_j}(\eta_j)} |p_{T_x M} - p_{T_0}|^2 d\mu)$$

for each  $j$  and  $B_\delta(0) \cap F \setminus \text{spt } \mu \subset \cup_j B_{5\sigma_j}(\eta_j)$ .  $F$  is the graph of the Lipschitz function  $f$  with  $\text{Lip } f \leq 1$ , so we of course have  $\mathcal{H}^n(B_{5\sigma_j}(\eta_j) \cap F) \leq C\sigma_j^n$  for each  $j$  with  $C = C(n)$ , hence by (13)

$$(14) \quad \begin{aligned} \mathcal{H}^n(B_\delta(0) \cap F \setminus \text{spt } \mu) &\leq \mathcal{H}^n(F \cap (\cup_j B_{5\sigma_j}(\eta_j))) \leq \sum_j \mathcal{H}^n(F \cap B_{5\sigma_j}(\eta_j)) \\ &\leq C \sum_j \sigma_j^n \leq C \sum_j (\mu(B_{\sigma_j}(\eta_j) \setminus F) + \int_{B_{\sigma_j}(\eta_j)} |p_{T_x M} - p_{T_0}|^2 d\mu) \\ &\leq C(\mu(\cup_j B_{\sigma_j}(\eta_j) \setminus F) + \int_{\cup_j B_{\sigma_j}(\eta_j)} |p_{T_x M} - p_{T_0}|^2 d\mu) \text{ by disjointness of } \{B_{\sigma_j}(\eta_j)\} \\ &\leq C(\mu(B_{2\delta}(0) \setminus F) + \int_{B_{2\delta}(0)} |p_{T_x M} - p_{T_0}|^2 d\mu), \end{aligned}$$

and  $\mu(B_{2\delta}(0) \setminus F) \leq CL^{-2} \int_{B_{3\delta}(0)} |p_{T_x M} - p_{T_0}|^2 d\mu$  by (6), so (7) is established and the proof is complete.  $\square$

**2.14 Corollary.** *There is a choice of  $\beta = \beta(n, k, p) \in (0, \frac{1}{4}]$  such that if the notation and assumptions are as in Lemma 2.13 and if*

$$\sup_{\sigma \in (0, \delta]} \sigma^{-n} \int_{B_\sigma(y)} |p_{T_x M} - p_{T_0}|^2 d\mu \leq \beta^2 L^2$$

for every  $y \in \text{spt } \mu \cap B_{2\delta}(0)$ , then

$$\text{spt } \mu \cap B_\delta(0) = \text{graph } f \cap B_\delta(0)$$

for some Lipschitz map  $f : \mathbb{R}^n \rightarrow \mathbb{R}^k$  with  $\text{Lip } f \leq L$ ,  $\sup |f| \leq C\delta^{\frac{1}{2n+2}}$ .

**Proof:** The hypotheses ensure that the set  $G$  in the above proof is *all of*  $\text{spt } \mu_V \cap B_{2\delta}(0)$ , and if  $\eta \in B_\delta(0) \cap \text{graph } f \setminus \text{spt } \mu_V$  then the inequality (12) in the above proof gives  $\sigma^n \leq C \int_{B_\sigma(\eta)} |p_{T_x M} - p_{T_0}|^2 d\mu_V \leq C\beta^2 \sigma^n$  for some  $\sigma \in (0, \delta]$ , which is evidently impossible for  $\beta = \beta(n, k, p)$  sufficiently small.  $\square$

### 3 Approximation by Harmonic Functions

The main result we shall need is given in the following lemma, which is an almost trivial consequence of Rellich's theorem:

**3.1 Lemma (Harmonic Approximation Lemma.)** *Given any  $\varepsilon > 0$  there is a constant  $\delta = \delta(n, \varepsilon) > 0$  such that if  $f \in W^{1,2}(B)$ ,  $B \equiv \check{B}_1(0) =$  open unit ball in  $\mathbb{R}^n$ , satisfies*

$$\int_B |\nabla f|^2 \leq 1, \quad \left| \int_B \nabla f \cdot \nabla \zeta \, d\mathcal{L}^n \right| \leq \delta \sup |\nabla \zeta|$$

*for every  $\zeta \in C_c^\infty(B)$ , then there is a harmonic function  $u$  on  $B$  such that  $\int_B |\nabla u|^2 \leq 1$  and*

$$\int_B (u - f)^2 \leq \varepsilon.$$

**Proof:** Suppose the lemma is false. Then we can find  $\varepsilon > 0$  and a sequence  $\{f_k\} \in W^{1,2}(B)$  such that

$$(1) \quad \left| \int_B \nabla f_k \cdot \nabla \zeta \, d\mathcal{L}^n \right| \leq k^{-1} \sup |\nabla \zeta|$$

for each  $\zeta \in C_c^\infty(B)$ , and

$$\int_B |\nabla f_k|^2 \leq 1,$$

but so that

$$(2) \quad \int_B (f_k - u)^2 > \varepsilon$$

whenever  $u$  is a harmonic function on  $B$  with  $\int_B |\nabla u|^2 \leq 1$ . Let  $\lambda_k = \omega_n^{-1} \int_B f_k \, d\mathcal{L}^n$ . Then by the Poincaré inequality (see e.g. [GT01]) we have

$$\int_B |f_k - \lambda_k|^2 \leq C \int_B |\nabla f_k|^2 \leq C,$$

and hence, by Rellich's theorem (see [GT01]), we have a subsequence  $\{k'\} \subset \{k\}$  such that  $f_{k'} - \lambda_{k'} \rightarrow w$  with respect to the  $\mathcal{L}^2(B)$  norm and  $\nabla f_{k'} \rightharpoonup \nabla w$  weakly in  $L^2$ , where  $w \in W^{1,2}(B)$  with  $\int_B |\nabla w|^2 \leq 1$ . By the weak convergence of  $\nabla f_{k'}$  to  $\nabla w$  and by (1) we evidently have

$$\int_B \nabla w \cdot \nabla \zeta \, d\mathcal{L}^n = \lim \int_B \nabla f_{k'} \cdot \nabla \zeta \, d\mathcal{L}^n = 0$$

for each  $\zeta \in C_c^\infty(B)$ . Thus  $w$  is harmonic in  $B$  and  $\int_B |f_{k'} - w - \lambda_{k'}|^2 \rightarrow 0$ . Since  $w + \lambda_{k'}$  is harmonic, this contradicts (2).  $\square$

We also recall the following standard estimates for harmonic functions (which follow directly from the mean-value property—see e.g. [GT01]): If  $u$  is harmonic on  $B \equiv B_\sigma(0)$ , then

$$3.2 \quad \sup_{B_{\sigma/2}(0)} \sigma^q |D^q u| \leq C \sigma^{-n/2} \|u\|_{L^2(B)}$$

for each integer  $q \geq 0$ , where  $C = C(q, n)$ . Indeed applying this with  $Du$  in place of  $u$  we get

$$3.3 \quad \sup_{B_{\sigma/2}(0)} \sigma^{q-1} |D^q u| \leq C \|\nabla u\|_{L^2(B)}$$

for  $q \geq 1$ . Using 3.2, 3.3 and an order 2 Taylor polynomial expansion for  $u$ , we see that if  $\ell$  is the affine approximation to  $u$  given by  $\ell(x) = u(0) + x \cdot \nabla u(0)$  then

$$3.4 \quad \begin{cases} |\ell(0)| = |u(0)| \leq C \sigma^{-n/2} \|u\|_{L^2(B)}, & |\nabla \ell| = |\nabla u(0)| \leq C \sigma^{-n/2} \|\nabla u\|_{L^2(B)} \\ \sup_{B_{\eta\sigma}(0)} |u - \ell| \leq (\eta\sigma)^2 \sup_{B_{\eta\sigma}} |D^2 u| \leq (\eta\sigma)^2 \sup_{B_{\sigma/2}} |D^2 u| \leq C \eta^2 \sigma^{1-n/2} \|\nabla u\|_{L^2(B)} \end{cases}$$

for  $\eta \in (0, \frac{1}{4}]$ , where  $C = C(n)$  is independent of  $\eta$ .

**3.5 Remark:** We note particular that the first two inequalities above can be applied to the approximating harmonic function  $u$  of Lemma 3.1, thus giving

$$\begin{aligned} |u(0)| &\leq C \|u\|_{L^2(B)} \leq C (\|u - f\|_{L^2(B)} + \|f\|_{L^2(B)}) \leq C (\sqrt{\varepsilon} + \|f\|_{L^2(B)}), \\ |\nabla u(0)| &\leq C \|\nabla u\|_{L^2(B)} \leq C \end{aligned}$$

for the harmonic approximating function of Lemma 3.1.

## 4 The Tilt-Excess Decay Lemma

In this section we continue to assume  $V$  has generalized mean curvature  $\underline{H}$  in  $U$  (as in Definition 3.15 of Ch. 4), and we write  $\mu$  for  $\mu_V$ .

We are now ready to discuss the following Tilt-excess Decay Theorem, which is the main result concerning tilt-excess needed for the regularity theorem of the next section. In this theorem the tilt excess  $E(\xi, \sigma, T)$  is as defined in 2.1, and we also use the notation

$$E_*(\xi, \sigma, T) = \max \left\{ E(\xi, \sigma, T), \delta^{-1} \left( \sigma^{p-n} \int_{B_\sigma(\xi)} |\underline{H}|^p d\mu \right)^{2/p} \right\},$$

where  $\delta$  is as in 2.6.

**4.1 Theorem (Tilt-excess Decay Theorem.)** *There are constants  $\eta = \eta(n, k, p)$ ,  $\delta_0 = \delta_0(n, k, p) \in (0, \frac{1}{16}]$  such that if  $\delta \in (0, \delta_0]$  and if hypotheses 2.6 hold, if  $\sigma \in (0, \delta\rho]$ ,  $\xi \in \text{spt } \mu_V \cap B_{\delta\rho}(0)$ , and if  $T$  is any  $n$ -dimensional subspace of  $\mathbb{R}^{n+k}$ , then*

$$E_*(\xi, \eta\sigma, S) \leq \eta^{2(1-n/p)} E_*(\xi, \sigma, T)$$

for some  $n$ -dimensional subspace  $S \subset \mathbb{R}^{n+k}$ .

**4.2 Remark:** Notice that any such  $S$  automatically satisfies

$$(\ddagger) \quad |p_S - p_T|^2 \leq C\eta^{-n} E_*(\xi, \sigma, T).$$

Indeed we trivially have

$$(\eta\sigma)^{-n} \int_{B_{\eta\sigma}(\xi)} |p_{T_x M} - p_T|^2 d\mu \leq \eta^{-n} E(\xi, \sigma, T),$$

while by 4.1 we have

$$(\eta\sigma)^{-n} \int_{B_{\eta\sigma}(\xi)} |p_{T_x M} - p_S|^2 d\mu \leq E_*(\xi, \sigma, T),$$

and hence, since  $|p_S - p_T|^2 \leq 2|p_{T_x M} - p_T|^2 + 2|p_{T_x M} - p_S|^2$ ,  $(\ddagger)$  follows by adding these inequalities and using the fact that  $\mu(B_{\eta\sigma}(\xi)) \geq \frac{1}{2}(\omega_n \eta\sigma)^n$  (by 2.8).

**Proof of 4.1:** Throughout the proof,  $C = C(n, k, p)$ . We can suppose (via translation and rotation of coordinates) that

$$(1) \quad \xi = 0, \quad T = \mathbb{R}^n \times \{0\}.$$

Let  $T_0 = T(0, 2\sigma)$  (notation as in Lemma 2.11). By Remark 2.12  $(\ddagger)$ ,  $T_0$  satisfies

$$(2) \quad E(0, \sigma, T_0) \leq C\delta^{\frac{1}{n+1}},$$

and hence we can assume that  $T$  also satisfies

$$(3) \quad E(0, \sigma, T) \leq C\delta^{\frac{1}{n+1}},$$

because otherwise we just prove the lemma with  $T_0$  in place of  $T$  and this then trivially implies the lemma for the original  $T$ . Since  $|p_T - p_{T_0}|^2 \leq 2|p_T - p_{T_x M}|^2 + 2|p_{T_x M} - p_{T_0}|^2$  we see from (2) and (3) that

$$|p_T - p_{T_0}| \leq C\delta^{\frac{1}{2n+2}}$$

and hence, since  $\sup_{x \in \text{spt } \mu \cap B_\sigma(0)} \text{dist}(x, T_0) \leq C \delta^{\frac{1}{2n+2}} \sigma$  by Lemma 2.11, we must also have  $\sup_{x \in \text{spt } \mu \cap B_\sigma(0)} \text{dist}(x, T) \leq C \delta^{\frac{1}{2n+2}} \sigma$ . Since  $T = \mathbb{R}^n \times \{0\}$ , this ensures

$$(4) \quad \sup_{B_\sigma(0) \cap \text{spt } \mu} \sum_{j=1}^k |x^{n+j}| \leq C \delta^{\frac{1}{2n+2}} \sigma.$$

By the Lipschitz Approximation Lemma 2.13, with  $L = 1$  and with  $\sigma/3$  in place of  $\delta\rho$ , there is a Lipschitz function  $f : B_{\sigma/3}(0) \rightarrow \mathbb{R}^k$  with

$$(5) \quad \begin{cases} \text{Lip } f \leq 1, \sup |f| \leq C \delta^{\frac{1}{2n+2}} \sigma \\ \mu(\text{spt } \mu \cap B_{\sigma/3}(0) \setminus F) + \mathcal{H}^n(F \cap B_{\sigma/3}(0) \setminus \text{spt } \mu) \leq C E_0 \sigma^n, \end{cases}$$

where, here and subsequently,

$$F = \text{graph } f,$$

and  $E_0 = E_*(0, \sigma, T)$ , i.e.

$$E_0 = \max \left\{ \sigma^{-n} \int_{B_\sigma(0)} |p_{T_x M} - p_T|^2 d\mu, \delta^{-1} (\sigma^{p-n} \int_{B_\sigma(0)} |\underline{H}|^p d\mu)^{2/p} \right\}.$$

Let us agree that  $C \delta^{\frac{1}{2n+2}} \leq \frac{1}{4}$ ,  $C = C(n, k, p)$  as in (4), in which case (4) implies

$$(6) \quad \text{spt } \mu \cap B_\sigma(0) \cap (B_{\sigma/4}^n(0) \times \mathbb{R}^k) \subset B_{\sigma/4}^n(0) \times B_{\sigma/4}^k(0) \subset \check{B}_{\sigma/3}(0).$$

Our aim now is to prove that each component of the Lipschitz function  $f$  is well-approximated by a harmonic function. Preparatory to this, note that the defining identity for  $\underline{H}$  (see 3.15 of Ch.4), with  $X = \zeta e_{n+j}$ , implies

$$\int_M \nabla_{n+j}^M \zeta d\mu = - \int_M e_{n+j} \cdot \underline{H} \zeta d\mu, \quad \zeta \in C_0^1(\check{B}_{\sigma/3}(0)),$$

$j = 1, \dots, k$ , where  $\nabla_{n+j}^M = e_{n+j} \cdot \nabla^M = p_{T_x M}(e_{n+j}) \cdot \nabla^M = (\nabla^M x^{n+j}) \cdot \nabla^M$  ( $\nabla^M$  = gradient operator for  $M$  as in §2 of Ch.3). Thus we can write

$$(7) \quad \int_M (\nabla^M x^{n+j}) \cdot \nabla^M \zeta d\mu = - \int_M e_{n+j} \cdot \underline{H} \zeta d\mu.$$

Since  $x^{n+j} = \tilde{f}^j(x)$  on  $M \cap F$  (where  $\tilde{f}^j$  is defined on  $\mathbb{R}^{n+k}$  by  $\tilde{f}^j(x^1, \dots, x^{n+k}) = f^j(x^1, \dots, x^n)$  for  $x = (x^1, \dots, x^{n+k}) \in \mathbb{R}^{n+k}$ ), we have by the definition of  $\nabla^M$  (see §2 of Ch.3) that

$$(8) \quad \nabla^M x^{n+j} = \nabla^M \tilde{f}^j(x) \quad \mu\text{-a.e. } x \in M \cap F.$$

Hence by (7) can be written

$$\int_{M \cap F} \nabla^M \tilde{f}^j \cdot \nabla^M \zeta d\mu = - \int_{M \setminus F} \nabla^M x^{n+j} \cdot \nabla^M \zeta d\mu - \int_M e_{n+j} \cdot \underline{H} \zeta d\mu,$$

and hence by (5), together with the fact that (by 2.6)

$$\int_{B_\sigma(\xi)} |\underline{H}| d\mu \leq \left( \int_{B_\sigma(\xi)} |\underline{H}|^p d\mu \right)^{1/p} (\mu(B_\sigma(\xi)))^{1-1/p} \leq C \delta^{\frac{1}{2}} E_0^{\frac{1}{2}} \sigma^{n-1},$$

we obtain

$$(9) \quad \left| \sigma^{-n} \int_{M \cap F} (\nabla^M \tilde{f}^j) \cdot \nabla^M \zeta d\mu \right| \leq C (\sigma^{-1} \sup |\zeta| \delta^{\frac{1}{2}} E_0^{\frac{1}{2}} + \sup |\nabla \zeta| E_0) \\ \leq C \sup |\nabla \zeta| (\delta^{\frac{1}{2}} E_0^{\frac{1}{2}} + E_0),$$

for any smooth  $\zeta$  with  $\text{spt } \zeta \subset \check{B}_{\sigma/3}(0)$ .

Furthermore by (8) and 2.4 we evidently have

$$(10) \quad \sigma^{-n} \int_{M \cap F \cap B_{\sigma/3}(0)} |\nabla^M \tilde{f}^j|^2 d\mu \leq 2E_0.$$

Now suppose that  $\zeta$  is an arbitrary  $C_c^1(\check{B}_{\sigma/3}^n(0))$  function, and let  $\tilde{\zeta}(x^1, \dots, x^{n+k}) = \zeta(x^1, \dots, x^n)$ , so  $\text{spt } \tilde{\zeta} = \text{spt } \zeta \times \mathbb{R}^k \subset \check{B}_{\sigma/3}^n(0) \times \mathbb{R}^k$ . By (6) there is a function  $\hat{\zeta} \in C_c^1(\check{B}_{\sigma/3}(0))$  which agrees with  $\tilde{\zeta}$  in a neighborhood of  $\text{spt } \mu \cap \text{spt } \tilde{\zeta}$ , and hence it is legitimate to use  $\hat{\zeta}$  in place of  $\zeta$  in the above discussion. Also,

$$(11) \quad \nabla^M \tilde{f}^j \cdot \nabla^M \tilde{\zeta} = \sum_{i,\ell=1}^n e^{i\ell} D_i \tilde{f}^j D_\ell \tilde{\zeta} = \nabla \tilde{f}^j \cdot \nabla \tilde{\zeta} - \sum_{i,\ell}^n (\delta_{i\ell} - e^{i\ell}) D_\ell \tilde{f}^j D_m \tilde{\zeta}, \\ \text{where } (e^{i\ell})_{i,\ell=1,\dots,n+k} \text{ is the matrix of } p_{T_x M}, \text{ and the maximum eigenvalue of } ((\delta_{i\ell} - e^{i\ell})_{i,\ell=1,\dots,n}) \text{ is } \leq \text{trace}((\delta_{i\ell} - e^{i\ell})_{i,\ell=1,\dots,n}), \text{ so, by Cauchy's inequality and 2.3,} \\ |\sum_{i,\ell=1}^n (\delta_{i\ell} - e^{i\ell}) \nabla_i \tilde{f}^j \nabla_\ell \tilde{\zeta}| \leq (n - \sum_{i=1}^n e^{ii}) |\nabla_i \tilde{f}^j| |\nabla_i \tilde{\zeta}| \\ = (\sum_{i=n+1}^{n+k} e^{ii}) |\nabla_i \tilde{f}^j| |\nabla_i \tilde{\zeta}| = \frac{1}{2} |p_{T_x M} - p_T|^2 |\nabla_i \tilde{f}^j| |\nabla_i \tilde{\zeta}|.$$

So, since  $|\nabla \tilde{f}^j| \leq 1$ , (11) gives

$$(12) \quad |\nabla^M \tilde{f}^j \cdot \nabla^M \tilde{\zeta} - \nabla \tilde{f}^j \cdot \nabla \tilde{\zeta}| \leq \frac{1}{2} |p_T - p_{T_x M}|^2 \sup |\nabla \tilde{\zeta}|.$$

Thus (9) and (12) imply

$$(13) \quad \left| \sigma^{-n} \int_{M \cap F} \nabla \tilde{f}^j \cdot \nabla \tilde{\zeta} d\mu \right| \leq C (\delta^{\frac{1}{2}} E_0^{\frac{1}{2}} + E_0) \sup |\nabla \zeta|.$$

Also since (12) is valid with  $\zeta = f^j$ , we conclude from (10) that

$$(14) \quad \sigma^{-n} \int_{M \cap F \cap B_{\sigma/3}(0)} |\nabla \tilde{f}^j|^2 d\mu \leq C E_0.$$

From 2.9, (5), (13), (14) and the area formula ?? of Ch.2 we then have

$$(15) \quad \left| \sigma^{-n} \int_{B_{\sigma/4}^n(0)} \nabla f^j \cdot \nabla \zeta \theta \circ G J_G d\mathcal{L}^n \right| \leq C \delta^{\frac{1}{2}} E_0^{\frac{1}{2}} \sup |\nabla \zeta|, \\ \sigma^{-n} \int_{B_{\sigma/4}^n(0)} |\nabla f^j|^2 \theta \circ G J_G d\mathcal{L}^n \leq C E_0,$$

where  $G : \mathbb{R}^n \rightarrow \mathbb{R}^{n+k}$  is the graph map defined by  $x \in B_{\sigma/3}^n(0) \mapsto G(x) = (x, f(x)) \in F \subset \mathbb{R}^{n+k}$ ,  $x \in B_{\sigma/3}^n(0)$ , and where  $J_G$  is the Jacobian of  $G$  defined, as in §3 of Ch.2, by

$$J_G(x) = \sqrt{\det(D_i G(x) \cdot D_j G(x))} = \sqrt{\det(\delta_{ij} + D_i f(x) \cdot D_j f(x))}.$$

Then  $1 \leq J_G \leq 1 + C|\nabla f|^2$  on  $B_\sigma^n(0)$  and  $1 \leq \theta \leq 1 + C\delta$  (by 2.8), so we conclude

$$(16) \quad \left| \sigma^{-n} \int_{B_\sigma^n(0)} \nabla f^j \cdot \nabla \zeta \, d\mathcal{L}^n \right| \leq C(\delta^{\frac{1}{2}} E_0^{\frac{1}{2}} + \delta \sigma^{-n} \int_{B_{\sigma/2}^n(0)} |\nabla f^j| \, d\mathcal{L}^n) \sup |\nabla \zeta| \\ \leq C\delta^{\frac{1}{2}} E_0^{1/2} \sup |\nabla \zeta|$$

by (15), because by (15) (and the fact that  $\theta \geq 1$ ,  $J_F \geq 1$ ) we have

$$(17) \quad \sigma^{-n} \int_{B_{\sigma/4}^n(0)} |\nabla f^j|^2 \, d\mathcal{L}^n \leq CE_0.$$

Now (16), (17) and the Harmonic Approximation 3.1 (with  $(CE_0)^{-1/2} f^j$  in place of  $f$ ) we know that for any given  $\varepsilon \in (0, 1)$  there is  $\delta_0 = \delta_0(n)$  such that, if the hypotheses of 3.1 hold with  $\delta \leq \delta_0$ , there are harmonic functions  $u^1, \dots, u^k$  on  $B_{\sigma/4}(0)$  such that

$$(18) \quad \sigma^{-n} \int_{B_{\sigma/4}^n(0)} |Du|^2 \, d\mathcal{L}^n \leq CE_0, \quad \sigma^{-n-2} \int_{B_{\sigma/4}^n(0)} |f - u|^2 \, d\mathcal{L}^n \leq \varepsilon E_0,$$

By (5) and Remark 3.5

$$(19) \quad \sigma^{-1} |u(0)| \leq C(\varepsilon^{1/2} E_0^{1/2} + \delta^{\frac{1}{2n+2}}) \leq C\delta^{\frac{1}{2n+2}}, \quad |\nabla u(0)| \leq CE_0^{1/2}.$$

Now, defining  $\lambda(x) = (\lambda^1(x), \dots, \lambda^k(x))$  with  $\lambda^j(x) = u^j(0) + x \cdot \nabla u^j(0)$  for  $j = 1, \dots, k$ , and again using 3.4 with  $\eta \in (0, \frac{1}{4})$ , we have also

$$(20) \quad (\eta\sigma)^{-n-2} \int_{B_{\eta\sigma}^n(0)} |f - \lambda|^2 \, d\mathcal{L}^n \leq 2(\eta\sigma)^{-n-2} \int_{B_{\eta\sigma}^n(0)} (|f - u|^2 + |u - \lambda|^2) \, d\mathcal{L}^n \\ \leq 2\eta^{-n-2} \varepsilon E_0 + 2\omega_n \eta^{-2} \sigma^{-2} \sup_{B_{\eta\sigma}(0)} |u - \lambda|^2 \\ \leq 2\eta^{-n-2} \varepsilon E_0 + C\eta^2 \sigma^{-n} \int_{B_{\eta\sigma}^n(0)} |Du|^2 \, d\mathcal{L}^n \\ \leq 2\eta^{-n-2} \varepsilon E_0 + C\eta^2 E_0,$$

where at the last step we used (18). Now let  $S$  be the  $n$ -dimensional subspace graph $(\lambda - \lambda(0))$ , let  $\tau = (0, \lambda(0))$ , and observe that  $\text{dist}(x, \tau + S) \leq |f(x') - \lambda(x')|$  for any  $x = (x', f(x')) \in B_{\eta\sigma}(\tau) \cap F$ , so (20) implies

$$(\eta\sigma)^{-n-2} \int_{B_{\eta\sigma}(\tau) \cap F} \text{dist}(x - \tau, S)^2 \, d\mathcal{H}^n \leq C\eta^{-n-2} \varepsilon E_0 + C\eta^2 E_0.$$

Then by (5), (4), and (19), keeping in mind  $\theta(\xi) \leq 1 + C\delta \leq 2$  in  $B_\sigma(0)$ ,

$$(\eta\sigma)^{-n-2} \int_{B_{\eta\sigma}(\tau)} \text{dist}(x - \tau, S)^2 d\mu \leq C\eta^{-n-2}(\varepsilon + \delta^{\frac{1}{n+1}})E_0 + C\eta^2 E_0,$$

and then by Remark 2.12 (§) we have

$$(21) \quad E(\tau, \eta\sigma/2, S) \leq C\eta^{-n-2}(\varepsilon + \delta^{\frac{1}{n+1}})E_0 + C(\eta^2 + \delta)E_0.$$

Now (19) implies  $|\tau| \leq C\delta^{\frac{1}{2n+2}}\sigma$ , hence

$$(22) \quad C\delta^{\frac{1}{2n+2}} < \eta/4 \Rightarrow B_{\eta\sigma/4}(0) \subset B_{\eta\sigma/2}(\tau)$$

(for  $\delta$  small enough depending on  $n, k, p$  and  $\eta$ ), and then (21) gives

$$(23) \quad E(0, \eta\sigma/4, S) \leq C\eta^{-n-2}(\varepsilon + \delta^{\frac{1}{n+1}})E_0 + C(\eta^2 + \delta)E_0.$$

The proof is now completed as follows:

With  $C = C(n, k, p)$  as in (23), first select  $\eta = \eta(n, k, p)$  so that  $C\eta^2 \leq \frac{1}{4}(\eta/4)^{2(1-n/p)}$ , and then choose  $\varepsilon = \varepsilon(n, k, p)$  so that  $C\eta^{-n-2}\varepsilon \leq \frac{1}{4}(\eta/4)^{2(1-n/p)}$ , and finally choose  $\delta \leq \delta_0(n, k, p)$  with  $\delta_0$  small enough so that  $B_{\eta\sigma/4}(0) \subset B_{\eta\sigma/2}(\tau)$  as in (22) and so that the above harmonic approximation is valid with the choice of  $\varepsilon$  made above, and also so that  $C\eta^{-n-2}\delta^{\frac{1}{n+1}} \leq \frac{1}{4}(\eta/4)^{2(1-n/p)}$ . Then (23) implies

$$E(0, \tilde{\eta}\sigma, S) \leq \tilde{\eta}^{2(1-n/p)}E_0,$$

where  $\tilde{\eta} = \eta/4$ . Since

$$((\tilde{\eta}\sigma)^{p-n} \int_{B_{\tilde{\eta}\sigma}(0)} |\underline{H}|^p d\mu)^{1/p} \leq \tilde{\eta}^{1-n/p}(\sigma^{p-n} \int_{B_\sigma(0)} |\underline{H}|^p d\mu)^{1/p}$$

by virtue of the inclusion  $B_{\tilde{\eta}\sigma}(0) \subset B_\sigma(0)$ , we thus conclude that

$$E_*(0, \tilde{\eta}\sigma, S) \leq \tilde{\eta}^{2(1-n/p)}E_*(0, \sigma, T).$$

This completes the proof of 4.1 (with  $\tilde{\eta}$  in place of  $\eta$ ).  $\square$

## 5 Main Regularity Theorem

We recall the hypotheses of §2 on  $V$  (which is a rectifiable varifold  $V = \underline{v}(M, \theta)$  with generalized mean curvature  $\underline{H}$  in the open set  $U \subset \mathbb{R}^{n+k}$ ):

$$5.1 \quad \begin{cases} \theta \geq 1 \text{ } \mu\text{-a.e.}, 0 \in \text{spt } V, B_\rho(0) \subset U \\ \omega_n^{-1}\rho^{-n}\mu(B_\rho(0)) \leq 1 + \delta, (\rho^{p-n} \int_{B_\rho(0)} |\underline{H}|^p d\mu)^{1/p} \leq \delta. \end{cases}$$

Then we have the following:

**5.2 Theorem (Allard Regularity Theorem.)** *If  $p > n$  is arbitrary, then there are  $\gamma = \gamma(n, k, p)$ ,  $\delta_0 = \delta_0(n, k, p) \in (0, \frac{1}{16}]$  such that if  $\delta \in (0, \delta_0]$  and if the hypotheses 5.1 hold, then there is an orthogonal transformation  $Q$  of  $\mathbb{R}^{n+k}$  and a  $u = (u^1, \dots, u^k) \in C^{1,1-n/p}(B_{\gamma\rho}^n(0); \mathbb{R}^k)$  with  $Du(0) = 0$ ,  $\text{spt } V \cap B_{\gamma\rho}(0) = Q(\text{graph } u) \cap B_{\gamma\rho}(0)$ , and*

$$\rho^{-1} \sup |u| + \sup |Du| + \rho^{1-n/p} \sup_{x, y \in B_{\gamma\rho}^n(0), x \neq y} |x - y|^{-(1-n/p)} |Du(x) - Du(y)| \leq C \delta^{\frac{1}{2n+2}},$$

where  $C = C(n, k, p) > 0$  and  $\gamma = \gamma(n, k, p) \in (0, 1)$ .

**5.3 Remark:** At the conclusion of this section we shall prove a slight improvement on the above theorem, in that for every  $\gamma \in (0, 1)$  there is  $\delta_0 = \delta_0(\gamma, n, k, p) \in (0, \frac{1}{16}]$  such that the hypotheses 5.1 with some  $\delta \leq \delta_0$  imply the conclusion of the above theorem.

In the proof of 5.2, we shall need the following corollary of the Affine Approximation Lemma 2.11, which shows that if  $T_0 = T(0, 4\delta)$  (notation as in 2.11), and if 5.1 holds then the tilt excess is  $\leq C\delta^{\frac{1}{n+1}}$  on every ball centered at 0 with radius  $< \rho$ .

**5.4 Lemma.** *There is  $\delta_0 = \delta_0(n, k, p) \in (0, \frac{1}{16}]$  such that if  $\delta \in (0, \delta_0]$ , if 2.6 holds, and if, with the notation of Lemma 2.11,  $T_0 = T(0, 4\delta\rho)$ , then, for each  $\theta \in (0, 1)$ ,*

$$\rho^{-n} \int_{B_{\theta\rho}(0)} |p_{T_x M} - p_{T_0}|^2 d\mu_V \leq C\delta^{\frac{1}{n+1}}, \quad C = C(\theta, n, k, p).$$

**Proof:** Let  $t \in [\delta, 1)$ . The inequality 3.30 of Ch. 4 with  $\rho = t \in (0, 1)$ ,  $\sigma = \delta t$  and with  $h(x) = |p_{T_0^\perp}(x/|x|)|^2$  implies that

$$\begin{aligned} \int_{\partial B_t(0)} h |\nabla^M r| d\mu_V &\leq \delta^{1-n} \int_{\partial B_{\delta t}(0)} h |\nabla^M r| d\mu_V + n \int_{B_1(0)} r^{-n} |D^\perp r|^2 d\mu_V \\ &\quad + C \int_{B_1(0) \setminus B_{\delta^2}(0)} (r^{-n} |D^\perp r| + r^{1-n} h |\underline{H}|) d\mu_V. \end{aligned}$$

By the Cauchy inequality  $ab \leq \frac{1}{2}a^2 + \frac{1}{2}b^2$  we have  $|D^\perp r| \leq \delta^{\frac{1}{2}} + \delta^{-1/2} |D^\perp r|^2$ , and since  $\int_{B_1(0) \setminus B_{\delta^2}(0)} r^{-n} d\mu_V \leq C |\log \delta|$  by Remark 4.9 of Ch. 4, we then get

$$\begin{aligned} (1) \quad \int_{\partial B_t(0)} h |\nabla^M r| d\mu_V &\leq \delta^{1-n} \int_{\partial B_{\delta t}(0)} h |\nabla^M r| d\mu_V \\ &\quad + C\delta^{-1/2} \int_{B_1(0)} r^{-n} |D^\perp r|^2 d\mu_V + C\delta^{\frac{1}{2}} |\log \delta| + \int_{B_1(0)} r^{1-n} h |\underline{H}| d\mu_V. \end{aligned}$$

Also, by the Hölder inequality and Remark 4.9 of Ch. 4,  $\int_{B_1} r^{1-n} h |\underline{H}| d\mu_V \leq C \|\underline{H}\|_{L^p(\mu_V \llcorner B_1)} C\delta$ , so in fact (1) gives

$$(2) \quad \int_{\partial B_t(0)} h |\nabla^M r| d\mu_V \leq \delta^{1-n} \int_{\partial B_{\delta t}(0)} h |\nabla^M r| d\mu_V + C\delta^{\frac{1}{2}} |\log \delta|.$$

Now by integrating over  $t \in (\delta, 1)$  and using the coarea identity 3.24 of Ch. 4 we get

$$\int_{B_1(0) \setminus B_{\delta}(0)} h |\nabla^M r|^2 d\mu_V \leq \delta^{-n} \int_{B_{\delta}(0) \setminus B_{\delta^2}(0)} h |\nabla^M r|^2 d\mu_V + C\delta^{\frac{1}{2}} |\log \delta|,$$

so

$$\int_{B_1(0)} h |\nabla^M r|^2 d\mu_V \leq (1 + \delta^{-n}) \int_{B_{\delta}(0)} h |\nabla^M r|^2 d\mu_V + C\delta^{\frac{1}{2}} |\log \delta|.$$

Since  $|\nabla^M r|^2 = 1 - |D^\perp r|^2$ , by 2.10 this gives

$$(3) \quad \int_{B_1(0)} h d\mu_V \leq (1 + \delta^{-n}) \int_{B_{\delta}(0)} h d\mu_V + C\delta^{\frac{1}{2}} |\log \delta| + C\delta,$$

and  $h(x) = |p_{T_0^\perp}(x/|x|)|^2 \leq C\delta^{\frac{1}{n+1}}$  for  $x \in \text{spt } \mu_V \cap B_{\delta}(0)$  by the Affine Approximation Lemma 2.11 (because  $T_0 = T(0, 4\delta)$ ), so (3) gives

$$(4) \quad \int_{B_1(0)} |p_{T_0^\perp}(x)|^2 d\mu_V \leq C\delta^{\frac{1}{n+1}}.$$

The proof is now completed by using 2.12 (‡) with  $T = T_0$ .  $\square$

**Proof of 5.2:** The proof is based on the Tilt-excess Decay 4.1 of the previous section. Throughout the proof  $C = C(n, k, p) > 0$ .

Take  $\xi \in B_{\delta\rho/2}(0) \cap \text{spt } V$  and  $\sigma \in (0, \delta\rho/2]$  and let  $S_0$  be an arbitrary  $n$ -dimensional subspace of  $\mathbb{R}^{n+k}$ . By the Tilt-excess Decay Theorem 4.1 we then know that there are  $\delta_0 = \delta_0(n, k, p)$ ,  $\eta = \eta(n, k, p)$  so that if  $\delta \leq \delta_0$  then 5.1 implies

$$E_*(\xi, \eta\sigma, S_1) \leq \eta^{2(1-n/p)} E_*(\xi, \sigma, S_0)$$

for suitable  $S_1$ . Notice that this can be repeated; by induction we prove that if  $\xi \in \text{spt } V \cap B_{\delta\rho/2}(0)$ , then, with  $\sigma_0 = \delta\rho/2$ , there is a sequence  $S_1, S_2, \dots$  of  $n$ -dimensional subspaces such that

$$(1) \quad E_*(\xi, \eta^j \sigma_0, S_j) \leq \eta^{2(1-n/p)} E_*(\xi, \eta^{j-1} \sigma_0/2, S_{j-1}) \leq \eta^{2(1-n/p)j} E_*(\xi, \sigma_0, S_0)$$

for each  $j \geq 1$ .

Let  $T_0 = T(0, 2\sigma_0)$ ; then 2.11 tells us that  $E(0, \sigma_0, T_0) \leq C\delta^{\frac{1}{n+1}}$  and hence, with the same  $C$ ,  $E(\xi, \sigma_0/2, T_0) \leq 2^N C\delta^{\frac{1}{n+1}}$  for each  $\xi \in \text{spt } \mu \cap B_{\sigma_0/2}(0)$ , so then the above, always taking  $S_0 = T_0$  (for each  $\xi \in B_{\sigma_0/2}(0) \cap \text{spt } \mu$ ) implies

$$(2) \quad E_*(\xi, \eta^j \sigma_0, S_j) \leq \eta^{2(1-n/p)} E_*(\xi, \eta^{j-1} \sigma_0/2, S_{j-1}) \leq \eta^{2(1-n/p)j} E_0,$$

where, here and subsequently,  $E_0 = E_*(0, \sigma_0, T_0)$ . Notice in particular that this gives (Cf. 4.2)

$$(3) \quad |p_{S_j} - p_{S_{j-1}}|^2 \leq C E_*(\xi, \eta^{j-1} \sigma_0, S_{j-1}) \leq C \eta^{2(1-n/p)j} E_*(\xi, \sigma_0, S_0).$$

for each  $j \geq 1$ .

By summation from  $j + 1$  to  $\ell$ , (3) gives

$$(4) \quad |p_{S_\ell} - p_{S_j}|^2 \leq C \eta^{2(1-n/p)j} E_0$$

for  $\ell \geq j \geq 0$ . (4) evidently implies that there is  $S(\xi) (= \lim_{\ell \rightarrow \infty} S_\ell)$  such that

$$(5) \quad |p_{S(\xi)} - p_{S_j}|^2 \leq C \eta^{2(1-n/p)j} E_0, \quad j = 0, 1, 2, \dots$$

In particular (setting  $j = 0$ )

$$(6) \quad |p_{S(\xi)} - p_{T_0}|^2 \leq C E_0.$$

Now if  $\sigma \in (0, \sigma_0/2]$  is arbitrary we can choose  $j \geq 0$  such that  $\eta^j \sigma_0/2 < \sigma \leq \eta^{j-1} \sigma_0/2$ . (1) and (5) imply

$$(7) \quad E_*(\xi, \sigma, S(\xi)) \leq C (\sigma/\sigma_0)^{2(1-n/p)} E_0, \quad C = C(n, k, p),$$

for each  $\xi \in B_{\sigma_0/2}(0) \cap \text{spt } V$  and each  $0 < \sigma \leq \sigma_0/2$ . Notice also that, by (6) and (7),

$$(8) \quad E_*(\xi, \sigma, T_0) \leq C E_0 \leq C \delta^{\frac{1}{2n+2}}, \quad 0 < \sigma \leq \sigma_0/2.$$

Supposing without loss of generality that  $T_0 = \mathbb{R}^n \times \{0\}$ , we then see, by Corollary 2.14 and (8), if  $L_0 \in (0, \frac{1}{4}]$  is given, and if  $\delta \leq \delta_0 L_0^{2n+2}$  for suitable  $\delta_0 = \delta_0(n, k, p)$ , then

$$(9) \quad \text{spt } V \cap B_{\sigma_0/4}(0) = \text{graph } f \cap B_{\sigma_0/4}(0),$$

where  $f$  is a Lipschitz function  $B_{\sigma_0/2}^n(0) \rightarrow \mathbb{R}^k$  with  $\text{Lip } f \leq L_0$ .

With such an  $f$ , let  $F = \text{graph } f$  and  $\xi = (\xi', f(\xi')) \in F$ , and note that, in view of (9), (7) implies

$$\lim_{\sigma \downarrow 0} \sigma^{-n} \int_{B_{\sigma}(\xi) \cap F} |p_{T_x F} - p_{S(\xi)}|^2 d\mathcal{H}^n = 0$$

for  $\mathcal{H}^n$ -a.e.  $\xi \in F \cap B_{\sigma_0/2}(0)$ , and at all such points  $\xi$  it evidently follows that  $S(\xi)$  is the approximate tangent space of  $F$ ; i.e.  $S(\xi) = p_{T_\xi F}$ , so (7) can be equivalently written

$$(10) \quad \sigma^{-n} \int_{B_{\sigma}(\xi) \cap F} |p_{T_x F} - p_{T_\xi F}|^2 d\mathcal{H}^n \leq C (\sigma/\sigma_0)^{2(1-n/p)} E_0$$

for all  $0 < \sigma \leq \sigma_0/2$ . Now the orthogonal projection  $p_{T_\xi F}$  of  $\mathbb{R}^{n+k}$  onto the subspace  $T_\xi F$  is given by  $p_{T_\xi F}(v) = \sum_{j=1}^n (\tau_j \cdot v) \tau_j$ , where  $\tau_j$  is an orthonormal basis for  $T_\xi F$ , and by the Gram-Schmidt orthogonalization process, starting with the basis  $(e_j, D_j f(\xi'))$ ,  $j = 1, \dots, n$ , for  $T_\xi F$ , where  $(\xi', f(\xi')) = \xi$ , shows that  $p_{T_\xi F}$  has matrix  $P_\xi$  of the form

$$P_\xi = \begin{pmatrix} I_{n \times n} & Df(\xi') \\ (Df(\xi'))^t & O_{k \times k} \end{pmatrix} + \mathcal{F}(Df(\xi')),$$

where  $\mathcal{F}(p)$  is a real analytic function of  $p = (p_{ij})_{i=1, \dots, n, j=1, \dots, k} \in \mathbb{R}^{nk}$  with  $\mathcal{F}(0) = 0$ ,  $D_p \mathcal{F}(0) = 0$  and hence  $|\mathcal{F}(p_1) - \mathcal{F}(p_2)| \leq C(n, k)(|p_1| + |p_2|)|p_1 - p_2|$  for  $|p_1|, |p_2| \leq 1$ . Evidently then (provided we choose  $L_0$  small enough, depending only on  $n, k$ ) we have

$$|Df(x') - Df(\xi')|^2 \leq |p_{T_x F} - p_{T_\xi F}|^2 \leq 3|Df(x') - Df(\xi')|^2$$

and so (10) implies

$$(11) \quad \sigma^{-n} \int_{B_\sigma^n(\xi')} |Df(x) - Df(\xi)|^2 d\mathcal{L}^n(x) \leq C(\sigma/\sigma_0)^{2(1-n/p)} E_0,$$

for all  $0 < \sigma < \sigma_0/4$ . For  $\mu$ -a.e.  $x_1, x_2 \in \text{spt } V \cap B_{\sigma_0/8}(0)$  we can use (11) with  $\sigma = |x_1 - x_2|$  and with  $\xi = x_1, x_2$ . Since  $|Df(x_1) - Df(x_2)|^2 \leq 2|Df(x) - Df(x_1)|^2 + 2|Df(x) - Df(x_2)|^2$  for  $x \in B_\sigma^n(x_1) \cap B_\sigma^n(x_2) \supset B_{\sigma/2}^n((x_1 + x_2)/2)$  we then conclude

$$|Df(x_1) - Df(x_2)| \leq C(|x_1 - x_2|/\sigma_0)^{1-n/p} E_0^{1/2}$$

for  $\mathcal{L}^n$ -a.e.  $x_1, x_2 \in B_{\sigma_0/4}^n(0)$ . Of course it follows that then  $f \in C^{1,1-n/p}$  and this holds for every  $x_1, x_2 \in B_{\sigma_0/4}^n(0)$ . Thus, choosing suitable  $\delta = \delta(n, k, p)$  to satisfy the smallness restrictions imposed in the above argument, the theorem is established with  $u = f$  and  $\gamma = \delta/4$ .  $\square$

As an application of The Conical Approximation Theorem 5.1 of Ch. 4 we establish the following corollary of the regularity theorem (Theorem 5.2), guaranteeing that the conclusion of the regularity theorem holds for any  $\gamma \in (0, 1)$  (rather than for small enough  $\gamma = \gamma(n, k, p)$ ), provided the hypotheses 5.1 hold with  $\delta$  sufficiently small depending on  $\gamma$ :

**5.5 Corollary.** *For each  $\gamma \in (0, 1)$  there is  $\delta_0 = \delta_0(n, k, p, \gamma) \in (0, 1)$  such that the hypotheses 5.1 with  $p > n$  and  $\delta \leq \delta_0$  imply the existence of a linear isometry  $q$  of  $\mathbb{R}^{n+k}$  and a function  $u = (u^1, \dots, u^k) \in C^{1,1-n/p}(B_{\gamma\rho}^n(0); \mathbb{R}^k)$  with  $Du(0) = 0$ ,  $\text{spt } V \cap B_{\gamma\rho}(0) =$*

$q(\text{graph } u) \cap B_{\gamma\rho}(0)$ , and

$$\rho^{-1} \sup |u| + \sup |Du| + \rho^{1-n/p} \sup_{x, y \in B_{\gamma\rho}^n(0), x \neq y} |x - y|^{-(1-n/p)} |Du(x) - Du(y)| \leq C\delta^{\frac{1}{2n+2}},$$

where  $C = C(n, k, p, \gamma)$ .

**Proof:** Let  $\sigma \in (0, \frac{1}{4}]$  and assume 5.1, where without loss of generality we can assume  $\rho = 1$ . Then the monotonicity inequalities 2.8 guarantee that the hypotheses of Theorem 5.1 of Ch. 4 are satisfied for any  $\xi \in \partial B_{1-\sigma}(0)$  with  $C\delta$  in place of  $\delta$  and  $C$  independent of  $\xi$ . Furthermore for  $\tau \leq \delta/2$  sufficiently small we have (again by 2.8) that  $(\omega_n \tau^n)^{-1} \mu(B_\tau(\tau\xi)) \leq 1 + C\delta$ . So Theorem 5.1 of Ch. 4 with  $\lambda = \delta$  and  $\rho = \sigma$  implies that

$$(\omega_n \sigma^n)^{-1} \mu(B_\sigma(\xi)) \leq 1 + C\delta^{1/4},$$

which means that for sufficiently small  $\delta = \delta(n, k, p) > 0$  and  $\xi \in \text{spt } \mu$ , we can apply Theorem 5.2 to give  $\text{spt } \mu \cap B_\delta(\xi) = q(\text{graph } u) \cap B_\delta(\xi)$ . In view of the arbitrariness of  $\xi$  this evidently gives the stated conclusion.  $\square$

## 6 Some Initial Applications of the Allard Theorem

The Allard Theorem of §5 is fundamental in the study of the regularity and compactness properties of rectifiable varifolds (including also smooth submanifolds) with prescribed (generalized) mean curvature, in particular in the study of stationary varifolds. Here we discuss some initial applications. First we have the following corollary of the Allard theorem 5.2.

**6.1 Theorem.** *If  $V = \underline{v}(M, \theta)$ , of dimension  $n$ , has generalized mean curvature  $\underline{H}$  (as in 3.15 of Ch. 4) in an open set  $U \subset \mathbb{R}^{n+k}$  and if  $\underline{H}$  is locally in  $L^p(\mu_V)$  for some  $p > n$ , if  $\theta \geq 1 \mu_V$ -a.e. in  $U$  and if  $\xi \in U$  with  $\Theta^n(\mu_V, \xi) = 1$ , then there is  $\rho > 0$  and an orthogonal  $Q$  and of  $\mathbb{R}^{n+k}$  such that*

$$(\ddagger) \quad Q \circ \tau(\text{spt } \mu_V) \cap \check{B}_\rho(0) = \text{graph } u, \quad \tau : x \mapsto x - \xi,$$

where  $u : W \rightarrow \mathbb{R}^k$ ,  $W$  open in  $\mathbb{R}^n$ , is a  $C^{1,1-n/p}(W, \mathbb{R}^k)$  function with  $u(0) = 0, |Du(0)| = 0$ .

*In case  $\theta$  is positive integer-valued  $\mu_V$ -a.e. in  $U$  and  $\underline{H} = h|_{\text{spt } \mu_V}$ , where  $h$  is a  $C^{q,\alpha}$  function in  $U$  for some  $q \in \{0, 1, 2, \dots\}$  and some  $\alpha \in (0, 1)$ , then, for sufficiently small  $\rho > 0$ , the above  $u$  is automatically  $C^{q+2,\alpha}(W)$  and  $\Theta^n(\mu_V, x) \equiv 1$  on  $\text{spt } \mu_V \cap \check{B}_\rho(\xi)$ .*

Finally, if  $\theta$  is positive integer-valued  $\mu_V$ -a.e. in  $U$ , if  $N \subset U$  is an  $(n + \ell)$ -dimensional  $C^{q+2,\alpha}$  embedded submanifold of  $\mathbb{R}^{n+k}$  (where  $\ell \leq k$ ) with  $\xi \in N$ , if  $V$  is stationary in  $N$  as in 2.8 of Ch. 4 (so that  $V$  has generalized mean curvature  $\underline{H} = \overline{H}_M$  in  $N$  as in 3.15 of Ch. 4), then again  $(\ddagger)$  holds for sufficiently small  $\rho > 0$ , with  $u \in C^{q+2,\alpha}(W)$  and  $\Theta^n(\mu_V, x) \equiv 1$  in  $\text{spt } \mu_V \check{B}_\rho(\xi)$ .

**Proof:** Since  $\lim_{\rho \downarrow 0} (\rho^{p-n} \int_{B_\rho(\xi)} |H|^p d\mu_V)^{1/p} = 0$  and  $\lim_{\rho \downarrow 0} (\omega_n \rho^n)^{-1} \mu_V(B_\rho(\xi)) = 1$ , we can choose  $\rho > 0$  such that the hypotheses of Theorem 5.2 hold, so, after applying the appropriate translation and orthogonal transformation, the required  $u$  exists with

$$(1) \quad \text{graph } u = \text{spt } V \cap B_\sigma(0)$$

with  $\sigma = \gamma\rho$ ,  $\gamma$  as in Theorem 5.2. Since  $\theta$  is integer valued and  $< 2$  a.e., we have  $\theta = 1$   $\mathcal{H}^n$ -a.e. on  $\text{graph } u$ ; but  $\text{graph } u$  is a  $C^1$  embedded submanifold so then  $\Theta^n(\mu_V, x) = 1$  at every point of  $\text{graph } u$ .

Let  $\varepsilon_0 \in (0, 1)$ . Since  $Du(0) = 0$ , by choosing a smaller  $\sigma$  if necessary we can assume that  $|Du| \leq \varepsilon_0$  on  $B_\sigma(0)$  and so the analysis we made in §1 of the present chapter is applicable and tells us that  $u$  satisfies a system of equations of the form 1.3; i.e.

$$(2) \quad \Delta u_i = \sum_{j=1}^n D_j(A_{ij}(Du)) + h_i, \quad i = 1, \dots, k,$$

with  $A_{ij}(P)$  are  $C^\infty$  functions of the variable  $P = (p_{\ell m})_{\ell=1, \dots, n, m=1, \dots, k}$  with  $|A_{ij}(P)| \leq C|P|^2$  and  $|D_P A_{ij}(P)| \leq C|P|$ , where  $C = C(n)$ . Then by the Schauder theory for elliptic equations we see that  $h_i \in C^{q,\alpha}(\check{B}_\sigma(0))$  implies that  $u \in C^{q+2,\alpha}(\check{B}_\sigma(0))$  as claimed.

Finally, assume  $V$  is stationary in  $N$ . Then we can apply Theorem 5.2 for each  $p > n$  so for each  $\alpha \in (0, 1)$  we have  $\sigma$  such that (1) holds with  $u \in C^{1,\alpha}(\check{B}_\sigma(0))$ . Then  $(e_i, D_i u(x)), i = 1, \dots, n$ , is a  $C^{0,\alpha}$  basis for  $T_{(x, u(x))} F$ ,  $F = \text{graph } u$ ,  $x \in \check{B}_\sigma(0)$ . By the Gram-Schmidt orthogonalization theorem we then have functions  $F_j(Du)$ ,  $j = 1, \dots, n$ , such that  $F_j(P)$  is a smooth function of  $P = (p_{ij})_{i=1, \dots, n, j=1, \dots, k}$  and  $F_1(Du(x)), \dots$ , is an orthonormal basis for  $T_{(x, u(x))} F$  for each  $x \in \check{B}_\sigma(0)$ . Then, by 2.8 of Ch. 4,  $F$  has generalized mean curvature at  $(x, u(x))$  equal to  $\sum_{j=1}^n \bar{B}_{u(x)}(F_j(Du(x)), F_j(Du(x)))$ . Thus, in this case (1) can be written

$$(3) \quad \Delta u_i = \sum_{j=1}^n D_j(A_{ij}(Du)) + \sum_{j=1}^n e_{n+i} \cdot \bar{B}_{(x, u(x))}(F_j(Du(x)), F_j(Du(x)))$$

for  $i = 1, \dots, k$ , and again standard elliptic theory implies  $u \in C^{q+2,\alpha}(\check{B}_\sigma(0))$ .  $\square$

**6.2 Definition:** If  $V = \underline{v}(M, \theta)$  is an  $n$ -dimensional rectifiable varifold, we say that a point  $\xi \in \text{spt } V$  is a *regular point* of  $V$  if there is a  $\rho > 0$  such that  $\check{B}_\rho(\xi) \cap \text{spt } V$  is an

$n$ -dimensional  $C^1$  embedded submanifold of  $\mathbb{R}^{n+k}$ . Then we let

$$\begin{aligned}\operatorname{reg} V &= \{\xi \in \operatorname{spt} V : \xi \text{ is a regular point of } V\} \\ \operatorname{sing} V &= \operatorname{spt} V \setminus \operatorname{reg} V.\end{aligned}$$

Notice that then by definition  $\operatorname{reg} V$ ,  $\operatorname{sing} V$  are respectively relatively open in  $\operatorname{spt} V$  and relatively closed in  $U$ .

**6.3 Corollary.** *If  $V = \underline{v}(M, \theta)$ , of dimension  $n$ , has generalized mean curvature  $\underline{H}$  in an open set  $U \subset \mathbb{R}^{n+k}$ , if  $\underline{H}$  is locally in  $L^p(\mu_V)$  for some  $p > n$ , and if  $\theta$  is positive integer-valued  $\mu_V$ -a.e. in  $\operatorname{spt} V$ , then  $\operatorname{reg} V$  is a relatively open dense set in  $\operatorname{spt} V$ ; i.e.  $\operatorname{sing} V$  is nowhere dense in  $\operatorname{spt} V$ , and  $\operatorname{spt} V$  is the closure, taken in  $U$ , of  $\operatorname{reg} V$ .*

**6.4 Remark:** It is an open question whether or not  $\operatorname{sing} V$  has  $\mathcal{H}^n$ -measure zero under the general conditions of the above corollary, even if we assume  $\underline{H} = 0$ ; such results (and much more) are true in the special case when  $V$  is the varifold associated with a minimizing current, as discussed below in Ch. 7.

**Proof of 6.3:** Take any ball  $B_\rho(\xi) \subset U$  and let

$$N = \min\{j \in \{1, 2, \dots\} : \Theta^n(\mu_V, x) = j \text{ for some } x \in \check{B}_\rho(\xi)\}.$$

Then  $\tilde{V} = \underline{v}(M, N^{-1}\theta) \llcorner \check{B}_\rho(\xi)$  has density  $\Theta^n(\mu_{\tilde{V}}, x) \geq 1$  everywhere in  $\operatorname{spt} \tilde{V} \cap \check{B}_\rho(\xi)$  and  $\Theta^n(\mu_{\tilde{V}}, x_0) = 1$  at some point of  $x_0 \in \check{B}_\rho(\xi)$ . Such a point  $x_0$  is in  $\operatorname{reg} \tilde{V} (= \operatorname{reg} V \cap \check{B}_\rho(\xi))$  by Theorem 6.1, so we have shown  $\operatorname{reg} V \cap \check{B}_\rho(\xi) \neq \emptyset$ .  $\square$

The Allard theorem will play a key role later (in Ch. 7) in establishing the regularity theory for solutions of the Plateau problem in arbitrary dimensions.

## CHAPTER 5 PROBLEMS

5.1 Let  $\gamma : [0, 1] \rightarrow \mathbb{R}^2$  be defined by  $\gamma(r) = r(\cos((\log(2/r))^\alpha), \sin((\log(2/r))^\alpha))$  for  $r \in (0, 1]$  and  $\gamma(0) = (0, 0)$ , where  $\alpha \in (0, \frac{1}{2})$ . Prove

(i)  $|\gamma'(r)| \leq 2$  and hence  $\mathcal{H}^1(\gamma([0, 1])) \leq 2$ .

(ii) If  $\Gamma = \gamma([0, 1]) \cup (-\gamma)([0, 1])$ , prove that the approximate tangent space  $T_0\Gamma$  does not exist, but that  $\Gamma$  has the strong affine approximation property at 0, meaning that for each  $\sigma \in (0, 1]$  there is a 1-dimensional subspace  $T_\sigma$  of  $\mathbb{R}^2$  with  $\sigma^{-1} \text{dist}(T_\sigma \cap B_\sigma(0), \Gamma \cap B_\sigma(0)) \rightarrow 0$  as  $\sigma \downarrow 0$ .

Here  $\text{dist}$  means as usual Hausdorff distance  $d(A, B) = \inf$  of all numbers  $\lambda > 0$  such that  $A$  is contained in the  $\lambda$ -nhd. of  $B$  and  $B$  is contained in the  $\lambda$ -nhd. of  $A$ .

5.2 With  $\Gamma$  as in 5.1 above, calculate  $p_{(T_x\Gamma)^\perp}(x)$  for  $x \in \Gamma \setminus \{0\}$ , and check that  $\int_\Gamma r^{-3} |p_{(T_x\Gamma)^\perp}(x)|^2 d\mathcal{H}^1 < \infty$ .

Note: 5.1, 5.2 suggest that finiteness of the term  $\int_\Gamma r^{-3} |p_{(T_x\Gamma)^\perp}(x)|^2 d\mathcal{H}^1$  (which is one of the key terms appearing in the monotonicity identity) does not in itself guarantee any especially strong asymptotic properties of  $\Gamma$  on approach to 0.

5.3 Let  $F(p) = \sqrt{\det(\delta_{ij} + p_i \cdot p_j)}$ , where  $p = (p_i^\alpha)_{i=1, \dots, n, \alpha=1, \dots, \ell} \in \mathbb{R}^{n\ell}$  and  $p_i = (p_i^1, \dots, p_i^\ell) \in \mathbb{R}^\ell$ .

(i) Prove that there is  $\varepsilon = \varepsilon(n, \ell) > 0$  such that  $F(p)$  is a convex function of  $p$  for  $|p| \leq \varepsilon$ .

(ii) Suppose  $u : \check{B}_\rho^n(0) \rightarrow \mathbb{R}^\ell$  is Lipschitz with  $\text{Lip } u \leq \varepsilon$  and let  $\mathcal{A}(u) = \mathcal{H}^n(\text{graph } u)$ . Prove that in fact  $\mathcal{A}(u) = \int_{\check{B}_\rho^n(0)} F(Du) d\mathcal{L}^n$ , and, if  $v : \check{B}_\rho^n(0) \rightarrow \mathbb{R}^\ell$  is also Lipschitz with  $\text{Lip } v \leq \varepsilon$ ,  $\mathcal{A}(u) - \mathcal{A}(v) \leq \int_{\check{B}_\rho(0)} \sum_{i,\alpha} A_i^\alpha(Du) D_i(u^\alpha - v^\alpha)$ , where  $A_i^\alpha(p) = \partial F(p) / \partial p_i^\alpha$ .

Hint: Let  $f(t) = \mathcal{A}(u + t(v - u))$ ,  $t \in [0, 1]$ , and use the 2nd order Taylor expansion  $f(1) = f(0) + f'(0) + \int_0^1 (1-t) f''(t) dt$  together with (i).

5.4  $u$  as in 5.2(ii) is said to be a weak solution of the minimal surface system (MSS) if it is a weak solution of the Euler-Lagrange system for the functional  $\mathcal{A}(u)$ ; that is  $\frac{d}{ds} \big|_{s=0} \mathcal{A}(u + s\zeta) = 0$  for each Lipschitz  $\zeta$  with compact support in  $\check{B}_\rho^n(0)$ .

(i) Prove that this is exactly the requirement that  $\sum_{i,\alpha} \int_{\check{B}_\rho^n(0)} A_i^\alpha(Du) D_i \zeta^\alpha d\mathcal{L}^n = 0$  for for each Lipschitz  $\zeta$  with compact support in  $\check{B}_\rho^n(0)$ .

(ii) Prove using 5.3(ii) that if  $u$  is a Lipschitz weak solution of the MSS as in (i) with  $\text{Lip } u \leq \varepsilon$ , then  $\mathcal{A}(u) \leq \mathcal{A}(v)$  for every Lipschitz  $v : \check{B}_\rho^n(0) \rightarrow \mathbb{R}^\ell$  which is such that  $v - u$  has compact support in  $\check{B}_\rho^n(0)$  and  $\text{Lip } v \leq \varepsilon$ .

(iii) If  $u$  is as in (ii) except that now  $\text{Lip } u \leq \varepsilon/2$ , prove that  $G = \text{graph } u$  (viewed as a multiplicity 1 rectifiable varifold in  $\check{B}_\rho^n(0) \times \mathbb{R}^\ell$ ) is stationary; i.e., prove that if

$\varphi_t(x) = x + tX|_x$  for  $x \in \check{B}_\rho^n(0) \times \mathbb{R}^\ell$  with  $X = (X^1, \dots, X^{n+\ell}) \in C^1$  with compact support in  $\check{B}_\rho^n(0) \times \mathbb{R}^\ell$ , then  $\frac{d}{dt}|_{t=0} \mathcal{H}^n(\varphi_t(G)) = 0$ .

Hint: Show that, for small enough  $t$ ,  $\varphi_t(G)$  is again the graph of a Lipschitz function  $u_t$  with  $\text{Lip } u_t < \varepsilon$ , and then use (ii).

Note: Having proved (iii), we can immediately apply the Allard regularity theorem to deduce that  $u$  is  $C^{1,\alpha}(\check{B}_{(1-\theta)\rho}(0))$  for any  $\alpha, \theta \in (0, 1)$  provided  $\varepsilon = \varepsilon(\alpha, n, \ell, \theta)$  is small enough.

# Chapter 6

## Currents

§ 1	Preliminaries: Vectors, Co-vectors, and Forms.....	155
§ 2	General Currents.....	159
§ 3	Integer Multiplicity Rectifiable Currents.....	172
§ 4	Slicing.....	180
§ 5	The Deformation Theorem.....	185
§ 6	Applications of Deformation Theorem.....	195
§ 7	The Flat Metric Topology.....	196
§ 8	The Rectifiability and Compactness Theorems.....	199

### 1 Preliminaries: Vectors, Co-vectors, and Forms

$e_1, \dots, e_P$  denote the standard orthonormal basis for  $\mathbb{R}^P$ . We let  $\Lambda^1(\mathbb{R}^P)$  denote the dual space of  $\mathbb{R}^P$ ; thus  $\Lambda^1(\mathbb{R}^P)$  is the space of linear functionals  $\omega : \mathbb{R}^P \rightarrow \mathbb{R}$ .  $dx^1, \dots, dx^P \in \Lambda^1(\mathbb{R}^P)$  will denote the basis for  $\Lambda^1(\mathbb{R}^P)$  dual to the standard basis  $e_1, \dots, e_P$  of  $\mathbb{R}^P$ . Thus for  $v = (v^1, \dots, v^P) \in \mathbb{R}^P$  we have

$$dx^j(v) = v^j, \quad j = 1, \dots, P.$$

For  $n \geq 2$ ,  $\Lambda^n(\mathbb{R}^P)$  denotes the space of alternating  $n$ -linear functions on  $\mathbb{R}^P \times \dots \times \mathbb{R}^P$  ( $n$  factors). Thus  $\omega \in \Lambda^n(\mathbb{R}^P)$  means  $\omega(v_1, \dots, v_n)$  is linear in each  $v_j$  and  $\omega(v_1, \dots, v_i, \dots, v_j, \dots, -v_i, \dots, v_j, \dots, v_n)$  for each  $i \neq j$ . If  $\omega_1, \dots, \omega_n \in \Lambda^1(\mathbb{R}^P)$  we define  $\omega_1 \wedge \omega_2 \wedge \dots \wedge \omega_n \in \Lambda^n(\mathbb{R}^P)$  by

$$1.1 \quad \omega_1 \wedge \omega_2 \wedge \dots \wedge \omega_n(v_1, \dots, v_n) = \sum_{\sigma} \operatorname{sgn} \sigma \, \omega_{\sigma(1)}(v_1) \omega_{\sigma(2)}(v_2) \cdots \omega_{\sigma(n)}(v_n) \quad (= \det(\omega_i(v_j))),$$

where the sum is over all permutations  $\sigma$  of  $\{1, \dots, n\}$  and where  $\text{sgn } \sigma$  is the sign of the permutation  $\sigma : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ .

Given  $v_1, \dots, v_n \in \mathbb{R}^P$ , we can write  $v_i = \sum_{j_i=1}^P v_i^{j_i} e_{j_i}$  for each  $i = 1, \dots, n$ , so for any  $\omega \in \Lambda^n(\mathbb{R}^P)$  we have, by the  $n$ -linearity of  $\omega$ ,

$$\omega(v_1, \dots, v_n) = \sum_{j_1, \dots, j_n=1}^P v_1^{j_1} \cdots v_n^{j_n} \omega(e_{j_1}, \dots, e_{j_n}),$$

and because  $\omega$  is alternating this sum can be restricted to *distinct*  $j_1, \dots, j_n$ . Then

$$\omega(v_1, \dots, v_n) = \sum_{\alpha=(j_1, \dots, j_n) \in I_{n,P}} \sum_{\sigma \in P_n} v_1^{j_{\sigma(1)}} \cdots v_n^{j_{\sigma(n)}} \omega(e_{j_{\sigma(1)}}, \dots, e_{j_{\sigma(n)}}),$$

where

$$1.2 \quad I_{n,P} = \{(j_1, \dots, j_n) \in \mathbb{Z}_+^n : 1 \leq j_1 < \cdots < j_n \leq P\},$$

and where  $P_n$  denotes the set of permutations  $\sigma$  of  $\{1, \dots, n\}$ . Since  $\omega$  is alternating we evidently have  $\omega(e_{j_{\sigma(1)}}, \dots, e_{j_{\sigma(n)}}) = \text{sgn}(\sigma) \omega(e_{j_1}, \dots, e_{j_n})$ , so

$$\omega(v_1, \dots, v_n) = \sum_{\alpha=(j_1, \dots, j_n) \in I_{n,P}} \sum_{\sigma \in P_n} \text{sgn}(\sigma) v_1^{j_{\sigma(1)}} \cdots v_n^{j_{\sigma(n)}} \omega(e_{j_1}, \dots, e_{j_n}).$$

But, according to the definition 1.1,  $\sum_{\sigma \in P_n} \text{sgn}(\sigma) v_1^{j_{\sigma(1)}} \cdots v_n^{j_{\sigma(n)}} = dx^\alpha(v_1, \dots, v_n)$ , where we use the notation

$$1.3 \quad dx^\alpha = dx^{j_1} \wedge \cdots \wedge dx^{j_n}, \quad \alpha = (j_1, \dots, j_n).$$

So we have proved that any  $\omega \in \Lambda^n(\mathbb{R}^P)$  can be written

$$1.4 \quad \omega = \sum_{\alpha \in I_{n,P}} \omega_\alpha dx^\alpha,$$

where  $\omega_\alpha = \omega(e_{j_1}, \dots, e_{j_n})$  for each  $\alpha = (j_1, \dots, j_n) \in I_{n,P}$ .

Thus  $\{dx^\alpha : \alpha \in I_{n,P}\}$  are a basis for  $\Lambda^n(\mathbb{R}^P)$  and dimension  $\Lambda^n(\mathbb{R}^P) = \binom{P}{n}$ .

For  $\omega = \sum_{\alpha \in I_{\ell,P}} \omega_\alpha dx^\alpha \in \Lambda^\ell(\mathbb{R}^P)$ ,  $\eta = \sum_{\beta \in I_{m,P}} \eta_\beta dx^\beta \in \Lambda^m(\mathbb{R}^P)$  we can define

$$1.5 \quad \omega \wedge \eta = \sum_{\alpha \in I_{\ell,P}, \beta \in I_{m,P}} \omega_\alpha \eta_\beta dx^\alpha \wedge dx^\beta \in \Lambda^{\ell+m}(\mathbb{R}^P).$$

This is consistent with 1.1, and for  $\omega, \omega_1, \omega_2 \in \Lambda^\ell(\mathbb{R}^P)$ ,  $\eta \in \Lambda^m(\mathbb{R}^P)$ ,  $v \in \Lambda^p(\mathbb{R}^P)$  we have

$$\begin{aligned} (c_1 \omega_1 + c_2 \omega_2) \wedge \eta &= c_1 \omega_1 \wedge \eta + c_2 \omega_2 \wedge \eta \\ (\omega \wedge \eta) \wedge v &= \omega \wedge (\eta \wedge v) \\ \omega \wedge \eta &= (-1)^{\ell m} \eta \wedge \omega. \end{aligned}$$

If  $V$  is a subspace of  $\mathbb{R}^P$  of  $\dim = \ell$  with basis  $v_1, \dots, v_\ell$  then  $\Lambda^n(V)$  denotes the subspace of  $\Lambda^n(\mathbb{R}^P)$  with basis  $\{v_{i_1}^* \wedge \dots \wedge v_{i_n}^* : (i_1, \dots, i_n) \in I_{n,\ell}\}$ , where  $v_j^* \in \Lambda^1(\mathbb{R}^P)$  is the element dual to  $v_j$ , so that, for  $v \in \mathbb{R}^P$ ,  $v^* \in \Lambda^1(\mathbb{R}^P)$  is defined by

$$1.6 \quad v^*(w) = v \cdot w, \quad w \in \mathbb{R}^P.$$

Analogous to the definition of  $\Lambda^n(\mathbb{R}^P)$ , we could similarly define  $\Lambda^n(\Lambda^1(\mathbb{R}^P))$  for  $n \geq 2$  as the space of alternating  $n$ -linear functions on  $\Lambda^1(\mathbb{R}^P)$ . In which case, after making the identification  $(dx^j)^* \simeq e_j$ , we have the space  $\Lambda_n(\mathbb{R}^P) \simeq \Lambda^n(\Lambda^1(\mathbb{R}^P))$  of  $n$ -vectors

$$w = \sum_{\alpha \in I_{n,P}} w^\alpha e_\alpha,$$

where  $w^\alpha \in \mathbb{R}$  and  $e_\alpha = e_{j_1} \wedge \dots \wedge e_{j_n}$  for  $\alpha = (j_1, \dots, j_n) \in I_{n,P}$ , and

$$\begin{aligned} v_1 \wedge \dots \wedge v_n &= \sum_{j_1, \dots, j_n=1}^P v_{1j_1} v_{2j_2} \dots v_{nj_n} e_{j_1} \wedge \dots \wedge e_{j_n} \\ &= \sum_{(\ell_1, \dots, \ell_n) \in I_{n,P}} \det(v_{i\ell_j}) e_{\ell_1} \wedge \dots \wedge e_{\ell_n} \end{aligned}$$

for any  $v_1, \dots, v_n \in \mathbb{R}^P$ .

If  $V$  is a subspace of  $\mathbb{R}^P$  of  $\dim = \ell$  with basis  $v_1, \dots, v_\ell$  then  $\Lambda_n(V)$  is the subspace of  $\Lambda_n(\mathbb{R}^P)$  spanned by  $\{v_{i_1} \wedge \dots \wedge v_{i_n} : (i_1, \dots, i_n) \in I_{n,\ell}\}$ .

$\omega \in \Lambda^n(\mathbb{R}^P)$  (respectively  $w \in \Lambda_n(\mathbb{R}^P)$ ) is called *simple* if it can be expressed  $\omega_1 \wedge \dots \wedge \omega_n$  with  $\omega_j \in \Lambda^1(\mathbb{R}^P)$  (respectively  $w_1 \wedge \dots \wedge w_n$  with  $w_j \in \mathbb{R}^P$ ).

We assume  $\Lambda_n(\mathbb{R}^P)$ ,  $\Lambda^n(\mathbb{R}^P)$  are equipped with the inner products naturally induced from  $\mathbb{R}^P$  (making  $\{e_\alpha\}_{\alpha \in I_{n,P}}$ ,  $\{dx^\alpha\}_{\alpha \in I_{n,P}}$  orthonormal bases). Thus

$$1.7 \quad \left( \sum_{\alpha \in I_{n,P}} \omega_\alpha dx^\alpha \right) \cdot \left( \sum_{\alpha \in I_{n,P}} \eta_\alpha dx^\alpha \right) = \sum_{\alpha \in I_{n,P}} \omega_\alpha \eta_\alpha$$

and

$$1.8 \quad \left( \sum_{\alpha \in I_{n,P}} u^\alpha e_\alpha \right) \cdot \left( \sum_{\alpha \in I_{n,P}} w^\alpha e_\alpha \right) = \sum_{\alpha \in I_{n,P}} u^\alpha w^\alpha$$

The dual pairing between  $\omega \in \Lambda^n(\mathbb{R}^P)$  and  $w \in \Lambda_n(\mathbb{R}^P)$  will be denoted  $\langle \omega, w \rangle$ ; thus

$$1.9 \quad \left\langle \sum_{\alpha \in I_{n,P}} \omega_\alpha dx^\alpha, \sum_{\alpha \in I_{n,P}} w^\alpha e_\alpha \right\rangle = \sum_{\alpha \in I_{n,P}} \omega_\alpha w^\alpha.$$

Given  $\ell : \mathbb{R}^P \rightarrow \mathbb{R}^Q$  linear, the “pull-back”  $\ell^\# : \Lambda^n(\mathbb{R}^Q) \rightarrow \Lambda^n(\mathbb{R}^P)$  is defined by

$$1.10 \quad \ell^\# \omega(v_1, \dots, v_n) = \omega(\ell(v_1), \dots, \ell(v_n)), \quad v_1, \dots, v_n \in \mathbb{R}^Q,$$

and then the “push-forward”  $\ell_\# : \Lambda^n(\mathbb{R}^P) \rightarrow \Lambda^n(\mathbb{R}^Q)$  is defined by duality according to the requirement

$$1.11 \quad \langle \ell^\# \omega, w \rangle = \langle \omega, \ell_\# w \rangle, \quad \omega \in \Lambda^n(\mathbb{R}^Q), \quad w \in \Lambda_n(\mathbb{R}^P),$$

where  $\langle \cdot, \cdot \rangle$  is the dual pairing as in 1.9. More explicitly,  $\ell^\#$ ,  $\ell_\#$  are then characterized as the unique linear maps  $\Lambda^n(\mathbb{R}^Q) \rightarrow \Lambda^n(\mathbb{R}^P)$  and  $\Lambda_n(\mathbb{R}^P) \rightarrow \Lambda_n(\mathbb{R}^Q)$  respectively such that for  $\omega_1, \dots, \omega_n \in \Lambda^1(\mathbb{R}^Q)$  and  $v_1, \dots, v_n \in \mathbb{R}^P$

$$1.12 \quad \begin{cases} \ell^\#(\omega_1 \wedge \dots \wedge \omega_n) = (\ell^\# \omega_1) \wedge \dots \wedge (\ell^\# \omega_n) = (\omega_1 \circ \ell) \wedge \dots \wedge (\omega_n \circ \ell) \\ \ell_\#(v_1 \wedge \dots \wedge v_n) = (\ell_\# v_1) \wedge \dots \wedge (\ell_\# v_n) = \ell(v_1) \wedge \dots \wedge \ell(v_n). \end{cases}$$

For open  $U \subset \mathbb{R}^P$ ,  $\mathcal{E}^n(U) = C^\infty(U, \Lambda^n(\mathbb{R}^P))$  and the elements  $\omega \in \mathcal{E}^n(U)$  are called smooth  $n$ -forms on  $U$ . Thus  $\omega \in \mathcal{E}^n(U)$  means  $\omega = \sum_{\alpha \in I_{n,P}} \omega_\alpha dx^\alpha$  where  $\omega_\alpha \in C^\infty(U)$ .

The value of  $\omega(x) = \sum_{\alpha \in I_{n,P}} \omega_\alpha(x) dx^\alpha$  at a point  $x \in U$  will also at times be denoted  $\omega|_x$ .

The exterior derivative  $\mathcal{E}^n(U) \rightarrow \mathcal{E}^{n+1}(U)$  is defined as usual by

$$1.13 \quad d\omega = \sum_{j=1}^P \sum_{\alpha \in I_{n,P}} \frac{\partial a_\alpha}{\partial x^j} dx^j \wedge dx^\alpha$$

if  $\omega = \sum_{\alpha \in I_{n,P}} a_\alpha dx^\alpha$ . By direct computation (using  $\frac{\partial^2 a_\alpha}{\partial x^i \partial x^j} = \frac{\partial^2 a_\alpha}{\partial x^j \partial x^i}$  and  $dx^i \wedge dx^j = -dx^j \wedge dx^i$ ) one checks that

$$1.14 \quad d^2\omega = 0 \quad \forall \omega \in \mathcal{E}^n(U).$$

Given  $\omega = \sum_{\alpha \in I_{n,Q}} \omega_\alpha(y) dy^\alpha \in \mathcal{E}^n(V)$ ,  $V \subset \mathbb{R}^Q$  open, and a smooth map  $f : U \rightarrow V$ , we define the “pulled back” form  $f^\# \omega \in \mathcal{E}^n(U)$  by

$$1.15 \quad f^\# \omega = \sum_{\alpha=(i_1, \dots, i_n) \in I_{n,Q}} \omega_\alpha \circ f df^{i_1} \wedge \dots \wedge df^{i_n},$$

where  $df^j$  is  $\sum_{i=1}^P \frac{\partial f^j}{\partial x^i} dx^i$ ,  $j = 1, \dots, Q$ . Equivalently this says

$$f^\# \omega|_x = (df_x)^\#(\omega|_{f(x)}),$$

where the right side is defined as in 1.10 with  $\ell = df_x$ .

Notice that the exterior derivative commutes with the pulling back:

$$1.16 \quad df^\# = f^\# d.$$

We let  $\mathcal{D}^n(U)$  denote the set of  $\omega = \sum_{\alpha \in I_{n,P}} \omega_\alpha dx^\alpha \in \mathcal{E}^n(U)$  such that each  $\omega_\alpha$  has compact support. We topologize  $\mathcal{D}^n(U)$  with the usual locally convex topology, characterized by the assertion that  $\omega_k = \sum_{\alpha \in I_{n,P}} \omega_{k\alpha} dx^\alpha \rightarrow \omega = \sum_{\alpha \in I_{n,P}} \omega_\alpha dx^\alpha$  if there is a fixed compact  $K \subset U$  such that  $\text{spt } \omega_{k\alpha} \subset K \quad \forall \alpha \in I_{n,P}, k \geq 1$ , and if

$\lim D^\beta \omega_{k\alpha} = D^\beta \omega_\alpha$  uniformly in  $K \forall \alpha \in I_{n,p}$  and every multi-index  $\beta$ . For any  $\omega \in \mathcal{D}^n(U)$ , we define

$$1.17 \quad |\omega| = \sup_{x \in U} \sqrt{\omega(x) \cdot \omega(x)}$$

If  $f : U \rightarrow V$  is smooth ( $U, V$  open in  $\mathbb{R}^P, \mathbb{R}^Q$  respectively) and if  $f$  is proper (i.e.  $f^{-1}(K)$  is a compact subset of  $U$  whenever  $K$  is a compact subset of  $V$ ) then  $f^\# \omega \in \mathcal{D}^n(U)$  whenever  $\omega \in \mathcal{D}^n(V)$ .

## 2 General Currents

Throughout this section  $U$  is an open subset of  $\mathbb{R}^P$ .

**2.1 Definition:** An  $n$ -dimensional current (briefly called an  $n$ -current) in  $U$  is a continuous linear functional on  $\mathcal{D}^n(U)$ . The set of such  $n$ -currents (i.e. the dual space of  $\mathcal{D}^n(U)$ ) will be denoted  $\mathcal{D}_n(U)$ .

Note that in case  $n = 0$  the  $n$ -currents in  $U$  are just the Schwartz distributions on  $U$ . More importantly though, the  $n$ -currents,  $n \geq 1$ , can be interpreted as a generalization of the  $n$ -dimensional oriented submanifolds  $M$  having locally finite  $\mathcal{H}^n$ -measure in  $U$ . Indeed given such an  $M \subset U$  with orientation  $\xi$  (thus  $\xi(x)$  is continuous on  $M$  with  $\xi(x) = \pm \tau_1 \wedge \cdots \wedge \tau_n \forall x \in M$ , where  $\tau_1, \dots, \tau_n$  is an orthonormal basis for  $T_x M$ )<sup>1</sup>, there is a corresponding  $n$ -current  $[[M]] \in \mathcal{D}_n(U)$  defined by

$$2.2 \quad [[M]](\omega) = \int_M \langle \omega(x), \xi(x) \rangle d\mathcal{H}^n(x), \quad \omega \in \mathcal{D}^n(U),$$

where  $\langle \cdot, \cdot \rangle$  denotes the dual pairing for  $\Lambda^n(\mathbb{R}^P)$ ,  $\Lambda_n(\mathbb{R}^P)$  as in 1.9. (That is, the  $n$ -current  $[[M]]$  is obtained by integration of  $n$ -forms over  $M$  in the usual sense of differential geometry:  $[[M]](\omega) = \int_M \omega$  in the usual notation of differential geometry.) In the special case then  $M = U$  (i.e.  $k = 0$  and  $M$  is just the open set  $U$  equipped with the standard orientation  $e_1 \wedge \cdots \wedge e_n$ ) we have, for  $\omega = a dx^1 \wedge \cdots \wedge dx^n \in \mathcal{D}^n(U)$ ,

$$2.3 \quad [[U]](\omega) = \int_U a(x) d\mathcal{L}^n(x).$$

Motivated by the classical Stokes' theorem ( $\int_M d\omega = \int_{\partial M} \omega$  if  $M$  is a compact smooth manifold with smooth boundary) we are led (by 2.2) to quite generally define the boundary  $\partial T$  of an  $n$ -current  $T \in \mathcal{D}_n(U)$  by

$$2.4 \quad \partial T(\omega) = T(d\omega), \quad \omega \in \mathcal{D}^n(U)$$

---

<sup>1</sup>Thus  $\xi(x) \in \Lambda_n(T_x M)$ ; notice this differs from the usual convention of differential geometry where we would take  $\xi(x) \in \Lambda^n(T_x M)$ .

(and  $\partial T = 0$  if  $n = 0$ ); thus  $\partial T \in \mathcal{D}_{n-1}(U)$  if  $T \in \mathcal{D}_n(U)$ , where to handle the case  $n = 0$  we make the notational agreement that  $\mathcal{D}_{-1}(U) = \{0\}$ .

Notice that  $\partial^2 T = 0$  by 1.14.

Again motivated by the special example  $T = \llbracket M \rrbracket$  as in 2.2 we define the *mass* of  $T$ ,  $\mathbb{M}(T)$ , for  $T \in \mathcal{D}_n(U)$  by

$$2.5 \quad \mathbb{M}(T) = \sup_{|\omega| \leq 1, \omega \in \mathcal{D}^n(U)} T(\omega)$$

(so that  $\mathbb{M}(T) = \mathcal{H}^n(M)$  in case  $T = \llbracket M \rrbracket$  as in 2.2). More generally for any open  $W \subset U$  we define

$$2.6 \quad \mathbb{M}_W(T) = \sup_{|\omega| \leq 1, \omega \in \mathcal{D}^n(U), \text{spt } \omega \subset W} T(\omega)$$

**2.7 Remark:** We here adopt the definition of  $\mathbb{M}(T)$  using the inner product norm  $|\omega|$ , but notice that there is some flexibility in this; we would still get the “correct” value  $\mathcal{H}^n(M)$  for the case  $T = \llbracket M \rrbracket$  if we were to make the definition  $\mathbb{M}(T) = \sup_{\|\omega(x)\| \leq 1, \omega \in \mathcal{D}^n(U)} T(\omega)$  where  $\|\omega(x)\|$  denotes the comass norm of  $\omega$  at  $x$ ; thus

$$\|\omega\| = \sup_{\xi \in \Lambda_n(\mathbb{R}^P), |\xi|=1, \xi \text{ simple}} \langle \omega, \xi \rangle.$$

Indeed in general this works (for  $T = \llbracket M \rrbracket$ ) provided only that  $\|\cdot\|$  is a norm for  $\Lambda^n(\mathbb{R}^P)$  with the properties:

- (a)  $\langle \omega, \xi \rangle \leq \|\omega\| |\xi|$  whenever  $\xi \in \Lambda_n(\mathbb{R}^P)$  is *simple*
- (b) For each fixed simple  $\xi \in \Lambda_n(\mathbb{R}^P)$ , equality holds in (a) for some  $\omega \neq 0$ .

Evidently the inner product norm and the comass norm are two such norms, but the comass norm is the *smallest* possible norm for  $\Lambda^n(\mathbb{R}^P)$  having these properties, which gives *maximality* of the corresponding definition of  $\mathbb{M}(T)$ . The reader is warned that  $\mathbb{M}(T)$  is usually defined in terms of the comass norm—this makes no significant difference to later discussion here but of course there will be contexts in which the difference becomes significant.

Suppose now  $T \in \mathcal{D}_n(U)$  satisfies  $\mathbb{M}_W(T) < \infty$  for every open  $W \subset\subset U$  and let  $\mathcal{C}^n(U)$  denote the set of *continuous*  $n$ -forms with compact support in  $U$ ; thus  $\omega \in \mathcal{C}^n(U)$  means  $\omega = \sum_{\alpha \in I_{n,P}} a_\alpha dx^\alpha$ , where  $a_\alpha$  are continuous functions with compact support in  $U$ . Given such a continuous  $\omega$  we can find a sequence  $\omega_j = \sum_{\alpha \in I_{n,P}} a_{j\alpha} dx^\alpha \in \mathcal{D}^n(U)$  with  $a_{j\alpha}$  converging to  $a_\alpha$  uniformly on  $U$  and with all  $a_{j\alpha}$  having compact support in a fixed  $W \subset\subset U$ . Then of course  $|T(\omega_j) - T(\omega_k)| = |T(\omega_j - \omega_k)| \leq \mathbb{M}_W(T) |\omega_j - \omega_k| \rightarrow 0$  as  $j, k \rightarrow \infty$ , so  $T(\omega_j)$  is a Cauchy sequence on  $\mathbb{R}$  and hence converges to some real number which we denote  $\bar{T}(\omega)$ . Evidently  $\bar{T}(\omega)$  is independent of the particular approximating

sequence  $\omega_j$  and also, so defined,  $\bar{T}$  is a bounded linear map  $\{\omega \in \mathcal{C}^n(U) : \text{spt } \omega \subset K\} \rightarrow \mathbb{R}$  for each compact  $K \subset U$ . But then the Riesz Representation Theorem 5.14 of Ch.1 is applicable and we deduce that there is a Radon measure  $\mu_T$  on  $U$  and an  $\mathcal{H}^n$ -measurable function  $\vec{T} : U \rightarrow \Lambda_n(\mathbb{R}^P)$  such that  $|\vec{T}| = 1$   $\mu_T$ -a.e. and

$$2.8 \quad \bar{T}(\omega) = \int_U \langle \omega, \vec{T} \rangle d\mu_T, \quad \omega \in \mathcal{C}^n(U),$$

and hence

$$2.9 \quad T(\omega) = \int_U \langle \omega, \vec{T} \rangle d\mu_T, \quad \omega \in \mathcal{D}^n(U).$$

$$2.10 \quad \mu_T(W) = \mathbb{M}_W(T) (= \sup_{\omega \in \mathcal{D}^n(U), \|\omega\| \leq 1, \text{spt } \omega \subset W} T(\omega))$$

for any open  $W$  with  $\bar{W}$  a compact subset of  $U$ . In particular

$$\mu_T(U) = \mathbb{M}(T).$$

$(\omega)$ ,

Notice that for such a  $T$  we can define, for any  $\mu_T$ -measurable subset  $A$  of  $U$  (and in particular for any Borel set  $A \subset U$ ), a new current  $T \llcorner A \in \mathcal{D}_n(U)$  by

$$2.11 \quad (T \llcorner A)(\omega) = \int_A \langle \omega, \vec{T} \rangle d\mu_T.$$

More generally, if  $\varphi$  is any locally  $\mu_T$ -integrable function on  $U$  then we can define  $T \llcorner \varphi \in \mathcal{D}_n(U)$  by

$$2.12 \quad (T \llcorner \varphi)(\omega) = \int \langle \omega, \vec{T} \rangle \varphi d\mu_T.$$

Given  $T \in \mathcal{D}_n(U)$  we define the *support*,  $\text{spt } T$ , of  $T$  to be the relatively closed subset of  $U$  defined by

$$2.13 \quad \text{spt } T = U \setminus \cup W$$

where the union is over all open sets  $W \subset\subset U$  such that  $T(\omega) = 0$  whenever  $\omega \in \mathcal{D}^n(U)$  with  $\text{spt } \omega \subset W$ . Notice that if  $\mathbb{M}_W(T) < \infty$  for each  $W \subset\subset U$  and if  $\mu_T$  is the corresponding total variation measure (as in 2.9, 2.10) then

$$2.14 \quad \text{spt } T = \text{spt } \mu_T$$

where  $\text{spt } \mu_T$  is the support of  $\mu_T$  in the usual sense of Radon measures in  $U$ .

Given a sequence  $\{T_q\} \subset \mathcal{D}_n(U)$ , we write  $T_q \rightharpoonup T$  in  $U$  ( $T \in \mathcal{D}_n(U)$ ) if  $\{T_q\}$  converges weakly to  $T$  in the usual sense of distributions:

$$2.15 \quad T_q \rightharpoonup T \iff \lim T_q(\omega) = T(\omega) \quad \forall \omega \in \mathcal{D}^n(U).$$

Notice that mass is trivially lower semi-continuous with respect to weak convergence: if  $T_q \rightharpoonup T$  in  $U$  then

$$2.16 \quad \mathbb{M}_W(T) \leq \liminf_{q \rightarrow \infty} M_W(T_q) \quad \forall \text{ open } W \subset U.$$

We also observe that if  $\sup_q \mathbb{M}_W(T_q) < \infty$  for each open  $W \subset\subset U$  then, by the discussion preceding 2.8,  $T_q$  extends uniquely to a linear functional  $\bar{T}_q$  on  $\mathcal{C}^n(U)$  such that  $|\bar{T}_q(\omega)| \leq \mathbb{M}_W(T_q)|\omega|$  for each  $\omega \in \mathcal{C}^n(U)$  with  $\text{spt } \omega \subset W$ . The weak convergence  $T_q \rightharpoonup T$  is thus equivalent to weak\* convergence with respect to continuous forms with compact support (i.e.  $\bar{T}_q(\omega) \rightarrow \bar{T}(\omega)$  for all continuous  $n$ -forms  $\omega$  on  $U$  with compact support), and hence by applying the standard Banach-Alaoglu theorem [Roy88] (in the Banach spaces  $\mathcal{M}_n(W) = \{T \in \mathcal{D}_n(W) : \mathbb{M}_W(T) < \infty\}$ ,  $W \subset\subset U$ ) we deduce

**2.17 Lemma.** *If  $\{T_q\} \subset \mathcal{D}_n(U)$  and  $\sup_{q \geq 1} \mathbb{M}_W(T_q) < \infty$  for each open  $W \subset\subset U$ , then there is a subsequence  $\{T_{q'}\}$  and a  $T \in \mathcal{D}_n(U)$  such that*

$$\int_U \langle \omega, \bar{T}_{q'} \rangle d\mu_{T_{q'}} \rightarrow \int_U \langle \omega, \bar{T} \rangle d\mu_T$$

for each continuous  $n$ -form  $\omega$  with compact support in  $U$ .

The following terminology will be used frequently:

**2.18 Terminology:** Given  $T_1 \in \mathcal{D}_n(U_1)$ ,  $T_2 \in \mathcal{D}_n(U_2)$  and an open  $W \subset U_1 \cap U_2$ , we say  $T_1 = T_2$  in  $W$  if  $T_1(\omega) = T_2(\omega)$  whenever  $\omega$  is a smooth  $n$ -form in  $\mathbb{R}^{n+k}$  with  $\text{spt } \omega \subset W$ .

Next we want to describe the cartesian product of currents  $T_1 \in \mathcal{D}_s(U_1)$ ,  $T_2 \in \mathcal{D}_t(U_2)$ ,  $U_1 \subset \mathbb{R}^{P_1}$ ,  $U_2 \subset \mathbb{R}^{P_2}$  open. We are motivated by the case when  $T_1 = \llbracket M_1 \rrbracket$  and  $T_2 = \llbracket M_2 \rrbracket$  (Cf. 2.2) where  $M_1, M_2$  are oriented submanifolds of dimension  $s, t$  respectively. We want to define  $T_1 \times T_2 \in \mathcal{D}_{s+t}(U_1 \times U_2)$  in such a way that for this special case (when  $T_j = \llbracket M_j \rrbracket$ ) we get  $\llbracket M_1 \rrbracket \times \llbracket M_2 \rrbracket = \llbracket M_1 \times M_2 \rrbracket$ . Since  $M_1 \times M_2$  has the natural orienting  $(s+t)$ -vector  $p_\#(\xi) \wedge q_\#(\eta)$ , where  $\xi$  and  $\eta$  are the orienting  $s$ -vector and  $t$ -vector for  $M_1, M_2$  respectively, and where  $p(x) = (x, 0)$ ,  $x \in \mathbb{R}^{P_1}$ , and  $q(y) = (0, y)$ ,  $y \in \mathbb{R}^{P_2}$ , we are thus led to the following definition:

**2.19 Definition:** If  $\omega \in \mathcal{D}^{s+t}(U_1 \times U_2)$  is written in the form

$$\omega = \sum_{(\alpha, \beta) \in I_{s', P_1} \times I_{t', P_2}, s' + t' = s + t} a_{\alpha\beta}(x, y) dx^\alpha \wedge dy^\beta$$

then we define

$$S \times T(\omega) = T(\sum_{\beta \in I_{t, P_2}} S(\sum_{\alpha \in I_{s, P_1}} a_{\alpha\beta}(x, y) dx^\alpha) dy^\beta),$$

which makes sense because if  $\text{spt } \omega = K$  then  $K \subset P(K) \times Q(K)$ , where  $P$  denotes the projection  $(x, y) \mapsto x$  of  $U_1 \times U_2 \rightarrow U_1$  and  $Q$  denotes the projection  $(x, y) \mapsto y$  of  $U_1 \times U_2 \rightarrow U_2$ , and one can check that  $S(\sum_{\alpha \in I_{s, P_1}} \omega_{\alpha\beta}(x, y) dx^\alpha)$  is a  $C_c^\infty(U_2)$  function of  $y$  with support in  $Q(K)$ .

Notice in particular this gives, for  $\omega_1 \in \mathcal{D}^{s'}(U_1)$ ,  $\omega_2 \in \mathcal{D}^{t'}(U_2)$  with  $s' + t' = s + t$  and with  $P, Q$  as above,

$$2.20 \quad S \times T((P^\# \omega_1) \wedge (Q^\# \omega_2)) = \begin{cases} S(\omega_1)T(\omega_2) & \text{if } (s', t') = (s, t) \\ 0 & \text{if } (s', t') \neq (s, t). \end{cases}$$

One readily checks, using Definition 2.19 and the definition of  $\partial$  (in 2.4), that

$$2.21 \quad \partial(S \times T) = (\partial S) \times T + (-1)^s S \times \partial T.$$

Notice this is valid also in case  $r$  or  $s = 0$  if we interpret the appropriate terms as zero; e.g.

$$2.22 \quad \partial(S \times T) = S \times \partial T \text{ if } s = 0.$$

Also (Cf. 4.5 of Ch.2), by 2.19 and 2.20,

$$2.23 \quad \mathbb{M}_{W_1 \times W_2}(S \times T) = \mathbb{M}_{W_1}(S) \mathbb{M}_{W_2}(T)$$

for any open  $W_1 \subset U_1$ ,  $W_2 \subset U_2$ , so if  $\mathbb{M}_W(S), \mathbb{M}_W(T) < \infty$  for each open  $W \subset U$  then also  $\mathbb{M}_W(S \times T) < \infty$  for each open  $W \subset U \times V$ . Also in this case one checks directly from the definition 2.19 that

$$2.24 \quad \overrightarrow{S \times T} = p_\# \vec{S} \wedge q_\# \vec{T}, \quad \mu_{S \times T} = \mu_S \times \mu_T,$$

where  $p(x) = (x, 0)$  and  $q(y) = (0, y)$ , and where  $\mu_S \times \mu_T$  is the product Borel regular measure characterized by the property

$$\int_{U \times V} f(x, y) d\mu_S \times \mu_T = \int_V \left( \int_U f(x, y) d\mu_S(x) \right) d\mu_T(y), \quad f \in C_c^0(U \times V).$$

An important special case of 2.21 occurs when we take  $T \in \mathcal{D}_n(U)$ ,  $U \subset \mathbb{R}^P$ , and we let  $\llbracket(0, 1)\rrbracket$  be the 1-current  $\in \mathcal{D}_1(\mathbb{R})$  defined as in 2.4 with  $M = (0, 1) \subset \mathbb{R}$  ( $(0, 1)$  having its usual orientation), so

$$\overrightarrow{\llbracket(0, 1)\rrbracket} = 1, \quad \mu_{\llbracket(0, 1)\rrbracket} = \mathcal{L}^1 \llcorner (0, 1)$$

and 2.24 says

$$2.25 \quad \overrightarrow{\llbracket(0, 1)\rrbracket \times T} = e_1 \wedge q_\# \vec{T}, \quad \mu_{S \times T} = \mathcal{L}^1 \times \mu_T.$$

Also, 2.21 gives

$$\begin{aligned} 2.26 \quad \partial(\llbracket(0, 1)\rrbracket \times T) &= (\{1\} - \{0\}) \times T - \llbracket(0, 1)\rrbracket \times \partial T \\ &= \{1\} \times T - \{0\} \times T - \llbracket(0, 1)\rrbracket \times \partial T. \end{aligned}$$

Here and subsequently  $\{p\}$ , for a point  $p \in U$ , means the 0-current  $\in \mathcal{D}_0(U)$  defined by

$$\{p\}(\omega) = \omega(p), \quad \omega \in \mathcal{D}^0(U) (= C_c^\infty(\tilde{U})).$$

Observe that then

$$\overrightarrow{\{p\}} = 1 \text{ and } \mu_{\{p\}} = \delta_{\{p\}},$$

where  $\delta_{\{p\}}$  is the point mass at  $p$  (i.e.  $\int_U f d\delta_{\{p\}} = f(p)$  for  $f \in C^0(U)$ ), and then 2.24 says

$$2.27 \quad \overrightarrow{\{p\} \times T} = q_{\#} \tilde{T}, \quad \mu_{\{p\} \times T} = \delta_{\{p\}} \times \mu_T.$$

Thus if  $\omega = \sum_{\alpha \in I_{s,P}, \beta \in I_{t,Q}, s+t=n} \omega_{\alpha\beta}(x, y) dx^\alpha dy^\beta \in \mathcal{D}^n(U \times V)$  with  $U \subset \mathbb{R}^P$  and  $V \subset \mathbb{R}^Q$  open, and if  $T \in \mathcal{D}_n(V)$ , then

$$2.28 \quad (\{p\} \times T)(\omega) = T(\sum_{\beta \in I_{n,Q}} \omega_{0,\beta}(p, y) dy^\beta).$$

Next we want to discuss the notion of “pushing forward” a current  $T$  via a smooth map  $f : U \rightarrow V$ ,  $U \subset \mathbb{R}^P$ ,  $V \subset \mathbb{R}^Q$  open. The main restriction needed is that  $f|_{\text{spt } T}$  is *proper*; that is  $f^{-1}(K) \cap \text{spt } T$  is a compact subset of  $U$  whenever  $K$  is a compact subset of  $V$ . Assuming this, we can define

$$2.29 \quad f_{\#}T(\omega) = T(\zeta f^{\#}\omega) \quad \forall \omega \in \mathcal{D}^n(V),$$

where  $\zeta$  is any function  $\in C_c^\infty(U)$  such that  $\zeta$  is identically equal to 1 in a neighborhood of the compact set  $\text{spt } T \cap \text{spt } f^{\#}\omega$ . The right side here certainly is defined because  $\zeta f^{\#}\omega$  has compact support in  $U$  (independent of any properness requirements on  $f$ ) and also the definition is independent of the particular choice of  $\zeta$ —if  $\tilde{\zeta}$  is another such choice then  $T(\zeta f^{\#}\omega) - T(\tilde{\zeta} f^{\#}\omega) = T((\zeta - \tilde{\zeta}) f^{\#}\omega) = 0$  because  $(\zeta - \tilde{\zeta}) f^{\#}\omega$  has compact support and is identically zero in a neighborhood of  $\text{spt } T$ .

**2.30 Remarks:** (1) Notice that if  $\mathbb{M}_W(T) < \infty$  for each  $W \subset\subset U$ , so that  $T$  has a representation as in 2.9, then, with  $f$  as in 2.29,  $f_{\#}T$  is given explicitly by

$$\begin{aligned} f_{\#}T(\omega) &= \int_U \langle f^{\#}\omega, \tilde{T} \rangle d\mu_T \\ &= \int_U \langle (df_x)^{\#}(\omega|_{f(x)}), \tilde{T}(x) \rangle d\mu_T(x) \\ &= \int_U \langle \omega|_{f(x)}, (df_x)_{\#}(\tilde{T}(x)) \rangle d\mu_T(x). \end{aligned}$$

Thus if  $\mathbb{M}_W(T) < \infty \forall W \subset\subset U$  we can make sense of  $f_\#T$  in case  $f$  is merely  $C^1$  with  $f|_{\text{spt } T}$  proper; note also that in this case

$$\mathbb{M}_W(f_\#T) \leq C \sup_{x \in f^{-1}W \cap \text{spt } T} \|df_x\|^n \mathbb{M}_{f^{-1}W}(T), \quad \forall W \subset\subset V,$$

with  $C = \binom{P}{n}^{1/2}$  (with  $C = 1$  if  $\vec{T}(x)$  is simple for  $\mu_T$ -a.e.  $x$ ).

(2) If  $T \in \mathcal{D}_n(U)$  with locally finite mass in  $U$ , if  $f : U \rightarrow V$  is  $C^1$  with  $f|_{\text{spt } T}$  proper, and if  $F$  is a closed subset of  $V$ , then, using the notation of 2.11 and 2.12,

$$(f_\#T) \llcorner F = f_\#(T \llcorner f^{-1}F).$$

One checks this by observing that there is a decreasing sequence  $g_j$  of  $C^\infty(U)$  functions with  $g_j(x) \rightarrow \chi_F(x)$  for every  $x \in U$ , so, using 2.9,  $((f_\#T) \llcorner F)(\omega) = \lim((f_\#T) \llcorner g_j)(\omega) = \lim(f_\#T)(g_j\omega) = \lim T(g_j \circ f \circ f^\#\omega) = \lim(f_\#(T \llcorner g_j \circ f))(\omega) = (f_\#(T \llcorner f^{-1}F))(\omega)$ ,  $\omega \in \mathcal{D}_n(V)$ .

(3) If  $T = \llbracket M \rrbracket$  as in 2.2, the above remark tells us that if  $f|_{\overline{M} \cap U}$  is proper, then

$$f_\#T(\omega) = \int_M \langle \omega|_{f(x)}, df_{x\#}\xi(x) \rangle d\mathcal{H}^n(x),$$

where  $\xi$  is the orientation for  $M$ . Notice that this makes sense if  $f$  is only Lipschitz (by virtue of Rademacher's Theorem 1.4 of Ch.2). If  $f$  is 1:1 and if  $J_f$  is the Jacobian of  $f$  as in ?? of Ch.2, then the area formula evidently tells us that (since  $df_{x\#}\xi(x) = J_f(x)\eta(f(x))$ , where  $\eta$  is the orientation for  $f(M_+)$ ,  $M_+ = \{x \in M : J_f(x) > 0\}$ , induced by  $f$ )

$$f_\#T(\omega) = \int_{f(M_+)} \langle \omega(y), \eta(y) \rangle d\mathcal{H}^n(y).$$

(Which confirms that our definition of  $f_\#T$  is "correct.")

Notice that the operations of pushing forward and taking boundaries *commute*:

$$2.31 \quad \partial f_\#T = f_\#\partial T, \quad T \in \mathcal{D}_n(U),$$

because, with  $\zeta$  as in 2.29,  $\partial f_\#T(\omega) = f_\#T(d\omega) = T(\zeta f^\#d\omega) = T(\zeta df^\#\omega) = T(d(\zeta f^\#\omega)) = \partial T(\zeta f^\#\omega) = f_\#\partial T$ , where we used 1.16.

We can now derive the important homotopy formula for currents as follows:

If  $f, g : U \rightarrow V$  are  $C^\infty$  ( $V \subset \mathbb{R}^Q$  open) and  $h : [0, 1] \times U \rightarrow V$  is  $C^\infty$  with  $h(0, x) \equiv f(x)$ ,  $h(1, x) \equiv g(x)$ , if  $T \in \mathcal{D}_n(U)$ , and if  $h|_{[0, 1] \times \text{spt } T}$  is proper, then, by 2.25, 2.26 and 2.27,

$$\begin{aligned} \partial h_\#(\llbracket(0, 1)\rrbracket \times T) &= h_\#\partial(\llbracket(0, 1)\rrbracket \times T) \\ &= h_\#(\{1\} \times T - \{0\} \times T - \llbracket(0, 1)\rrbracket \times \partial T) \\ &= g_\#T - f_\#T - h_\#(\llbracket(0, 1)\rrbracket \times \partial T). \end{aligned}$$

Thus, subject to the conditions stated above, we obtain the *homotopy formula*

$$2.32 \quad g_{\#}T - f_{\#}T = \partial h_{\#}(\llbracket(0, 1)\rrbracket \times T) + h_{\#}(\llbracket(0, 1)\rrbracket \times \partial T).$$

Notice that an important case of the above is given by

$$2.33 \quad h(t, x) = tg(x) + (1 - t)f(x) = f(x) + t(g(x) - f(x))$$

(i.e.  $h$  is an “affine homotopy” from  $f$  to  $g$ ). In this case we note that if  $h|_{\text{spt } T}$  is a proper map into  $V$  then  $W \subset\subset V \Rightarrow \text{spt}(\llbracket(0, 1)\rrbracket \times T) \cap h^{-1}(W) \subset\subset [0, 1] \times U$  and hence  $\text{spt } T \cap P(h^{-1}(W)) \subset\subset U$ , where  $P$  is the projection  $(t, x) \mapsto x$ . Then by the integral representation 2.9 and Remark 2.30(1) above we have, for any open  $W \subset\subset V$ ,

$$2.34 \quad \mathbb{M}_W(h_{\#}\llbracket(0, 1)\rrbracket \times T) \leq C \sup_{x \in \text{spt } T \cap W_h} |f - g| \cdot \sup_{x \in \text{spt } T \cap W_h} (|df_x| + |dg_x|)^n \mathbb{M}_{W_h}(T),$$

where  $W_h = Q(h^{-1}(W))$ , with  $Q : (t, x) \mapsto x$ . Indeed by 2.25 and 2.30(1) we have, for any  $\omega \in \mathcal{D}^n(V)$ ,

$$\begin{aligned} h_{\#}(\llbracket(0, 1)\rrbracket \times T)(\omega) &= \int_0^1 \int_U \langle \omega|_{h(t, x)}, dh_{(t, x)\#}(e_1 \wedge q_{\#}\tilde{T}(x)) \rangle d\mu_T(x) dt \\ &= \int_0^1 \int_U \langle \omega|_{h(t, x)}, (g(x) - f(x)) \\ &\quad \wedge (tdg_x + (1 - t)df_x)_{\#}\tilde{T}(x) \rangle d\mu_T(x) dt, \end{aligned}$$

and 2.34 follows immediately.

We now give a couple of important applications of the above homotopy formula.

**2.35 Lemma:** *If  $T \in \mathcal{D}_n(U)$ ,  $\mathbb{M}_W(T), \mathbb{M}_W(\partial T) < \infty \forall W \subset\subset U$  and if  $f, g : U \rightarrow V$  are  $C^1$  with  $f = g$  on  $\text{spt } T$ , and if  $h$  is as in 2.33 with  $h(\text{spt } T) \subset V$  and  $h|_{\text{spt } T}$  proper, then  $f_{\#}T = g_{\#}T$ . (Note that  $f_{\#}T, g_{\#}T$  are well-defined by 2.30(1).)*

**Proof:** By the homotopy formula 2.32 we have, with  $h(t, x) = tg(x) + (1 - t)f(x)$ ,

$$\begin{aligned} g_{\#}T(\omega) - f_{\#}T(\omega) &= \partial h_{\#}(\llbracket(0, 1)\rrbracket \times T) + h_{\#}(\llbracket(0, 1)\rrbracket \times \partial T)(\omega) \\ &= h_{\#}(\llbracket(0, 1)\rrbracket \times T)(d\omega) + h_{\#}(\llbracket(0, 1)\rrbracket \times \partial T)(\omega), \end{aligned}$$

so that, by 2.34, for a suitable  $C$  depending on  $T$  and  $\omega$ , we have

$$|f_{\#}T(\omega) - g_{\#}T(\omega)| \leq C \sup_{x \in \text{spt } T} |f - g| = 0$$

since  $f = g$  on  $\text{spt } T$ .  $\square$

The homotopy formula also enables us to define  $f_{\#}T$  in case  $f$  is merely Lipschitz, provided  $f|_{\text{spt } T}$  is proper and  $\mathbb{M}_W(T), \mathbb{M}_W(\partial T) < \infty \forall W \subset\subset U$ . In the following

lemma we let  $f_\sigma = f * \varphi^{(\sigma)}$ ,  $\varphi^{(\sigma)}(x) = \sigma^{-P} \varphi(\sigma^{-1}x)$ , with  $\varphi$  a mollifier as in §2 of Ch. 2.

**2.36 Lemma.** *If  $T \in \mathcal{D}_n(U)$ ,  $\mathbb{M}_W(T)$ ,  $\mathbb{M}_W(\partial T) < \infty \forall W \subset\subset U$ , and if  $f : U \rightarrow V$  is Lipschitz with  $f|_{\text{spt } T}$  proper, then  $\lim_{\sigma \downarrow 0} f_{\sigma\#}T(\omega)$  exists for each  $\omega \in \mathcal{D}^n(V)$ ;  $f_{\#}T(\omega)$  is defined to be this limit; then  $\text{spt } f_{\#}T \subset f(\text{spt } T)$  and  $\mathbb{M}_W(f_{\#}T) \leq (\text{ess sup}_{f^{-1}(W)} |Df|)^n \mathbb{M}_{f^{-1}(W)} \forall W \subset\subset V$ .*

**Proof:** If  $\sigma, \tau$  are sufficiently small (depending on  $\omega$ ) then the homotopy formula gives

$$f_{\sigma\#}T(\omega) - f_{\tau\#}T(\omega) = h_{\#}(\llbracket(0, 1)\rrbracket \times T)(d\omega) + h_{\#}(\llbracket(0, 1)\rrbracket \times \partial T)(\omega)$$

where  $h : [0, 1] \times U \rightarrow V$  is defined by  $h(t, x) = tf_\sigma(x) + (1 - t)f_\tau(x)$ . Then by 2.34, for sufficiently small  $\sigma, \tau$ , we have

$$|f_{\sigma\#}T(\omega) - f_{\tau\#}T(\omega)| \leq C \sup_{f^{-1}(K) \cap \text{spt } T} |f_\sigma - f_\tau| \cdot (\text{Lip } f)^n,$$

where  $K$  is a compact subset of  $V$  with  $\text{spt } \omega \subset \text{interior}(K)$ . Since  $f_\sigma \rightarrow f$  uniformly on compact subsets of  $U$ , the result now clearly follows.  $\square$

Next we want to define the notion of the *cone* over a given current  $T \in \mathcal{D}_n(U)$ . We want to define this in such a way that if  $T = \llbracket M \rrbracket$  where  $M$  is a submanifold of  $\mathbb{S}^{P-1} \subset \mathbb{R}^P$  then the cone over  $T$  is just  $\llbracket C_M \rrbracket$ ,  $C_M = \{\lambda x : x \in M, 0 < \lambda \leq 1\}$ . We are thus led generally to make the definition that the cone over  $T$ , denoted  $0 \mathbb{M} T$ , is defined by

$$2.37 \quad 0 \mathbb{M} T = h_{\#}(\llbracket(0, 1)\rrbracket \times T)$$

whenever  $T \in \mathcal{D}_n(U)$ ,  $n \geq 1$ , with  $U$  star-shaped relative to 0 and  $\text{spt } T$  compact, where  $h : [0, 1] \times \mathbb{R}^P \rightarrow \mathbb{R}^P$  is defined by  $h(t, x) = tx$ . Notice that  $h$  is an affine homotopy  $tg(x) + (1 - t)f(x)$ , where  $g(x) = x$  and  $f(x) = 0$ . Thus  $0 \mathbb{M} T \in \mathcal{D}_{n+1}(U)$  and (by the homotopy formula)

$$2.38 \quad \partial(0 \mathbb{M} T) = T - 0 \mathbb{M} \partial T.$$

Notice in particular that, with  $R = 0 \mathbb{M} T$ , we have thus established that

$$2.39 \quad \begin{aligned} &U \text{ star-shaped relative to } 0 \text{ and } T \in \mathcal{D}_n(U), n \geq 1, \text{ with } \text{spt } T \\ &\text{compact and } \partial T = 0 \Rightarrow \exists R \in \mathcal{D}_{n+1}(U) \text{ with } \text{spt } R \text{ compact and } \partial R = T. \end{aligned}$$

As a final application of the homotopy formula we have the following lemma which gives a sufficient condition to ensure that a given current of locally finite mass is conical—i.e. invariant under homotheties  $\eta_{0,\lambda}$ :

**2.40 Lemma.** Suppose  $C \in \mathcal{D}_n(\mathbb{R}^{n+k})$  with  $\mathbb{M}_{\check{B}_R(0)}(C) < \infty$  for each  $R > 0$ ,  $\partial C = 0$ , and  $x \wedge \vec{C}|_x = 0$   $\mu_C$ -a.e. Then  $\eta_{0,\lambda\#}C = C$  for each  $\lambda > 0$ .

**Proof:** We apply the homotopy formula 2.32 with  $f(x) = x$  and  $g(x) = \lambda^{-1}x$ , and  $h(t, x) = tg(x) + (1-t)f(x)$ . Then

$$\eta_{0,\lambda\#}C - C = \partial h_{\#}(\overline{[(0,1)]} \times C).$$

The right side here is zero because  $\overline{[(0,1)]} \times \vec{C} = e_1 \wedge q_{\#}\vec{C}$ , where  $q(x) = (0, x)$ , and hence

$$h_{\#}|_{(t,x)} \overline{[(0,1)]} \times \vec{C}|_{(t,x)} = (1+t(\lambda^{-1}-1))^n (\lambda^{-1}-1)x \wedge \vec{C}|_x = 0. \quad \square$$

The following Constancy Theorem is very useful:

**2.41 Theorem.** If  $U$  is open in  $\mathbb{R}^n$  (i.e.  $P = n$ ), if  $U$  is connected, if  $T \in \mathcal{D}_n(U)$  and  $\partial T = 0$ , then there is a constant  $c$  such that  $T = c\llbracket U \rrbracket$  (using the notation of 2.3).

**Proof:** Let  $\varphi^{(\sigma)}(x) = \sigma^{-n}\varphi(\sigma^{-1}x)$ , with  $\varphi$  a mollifier as in §2 of Ch.2. For any closed ball  $B_{\rho}(x_0) \subset U$  pick  $\sigma_0 > 0$  such that  $B_{\rho+\sigma_0}(x_0) \subset U$  and take  $a \in L^1(\mathbb{R}^n)$  with  $a = 0$  on  $\mathbb{R}^n \setminus B_{\rho}(x_0)$ . Then we have  $a_{\sigma} \in C_c^{\infty}(U)$  for  $\sigma < \sigma_0$ , ( $a_{\sigma} = \varphi^{(\sigma)} * a$ ), and  $D^{\beta}a_{\sigma} = (D^{\beta}\varphi^{(\sigma)}) * a$  for each multi-index  $\beta$ , so if  $a_j \rightarrow a$  in  $L^1(\check{B}_{\rho}(x_0))$  with  $a_j = 0$  on  $\mathbb{R}^n \setminus \check{B}_{\rho}(x_0)$  then  $a_{j\sigma} dx^1 \wedge \cdots \wedge dx^n \rightarrow a_{\sigma} dx^1 \wedge \cdots \wedge dx^n$  in  $\mathcal{D}^n(U)$  for  $\sigma < \sigma_0$ , and hence

$$T(a_{j\sigma} dx^1 \wedge \cdots \wedge dx^n) \rightarrow T(a_{\sigma} dx^1 \wedge \cdots \wedge dx^n).$$

Thus the functional  $F_{\sigma} : L^1(\check{B}_{\rho}(x_0)) \rightarrow \mathbb{R}$  defined by

$$F_{\sigma}(a) = T(a_{\sigma} dx^1 \wedge \cdots \wedge dx^n)$$

is a bounded linear functional on  $L^1(\check{B}_{\rho}(x_0))$ , and by the Riesz Representation Theorem for  $L^1(\check{B}_{\rho}(x_0))$  there is a bounded measurable function  $\theta^{(\sigma)}$  in  $\check{B}_{\rho}(x_0)$  with  $F_{\sigma}(a) = \int_{\check{B}_{\rho}(x_0)} a \theta^{(\sigma)} d\mathcal{L}^n$  for  $a \in L^1(\check{B}_{\rho}(x_0))$ , and hence in particular

$$(1) \quad T(a_{\sigma} dx^1 \wedge \cdots \wedge dx^n) = \int a \theta^{(\sigma)} d\mathcal{L}^n, \quad a \in C_c^{\infty}(\check{B}_{\rho}(x_0)).$$

Now for  $j = 1, \dots, n$  let  $\omega_{j\sigma} = (-1)^{j-1} a_{\sigma} dx^1 \wedge \cdots \wedge dx^{j-1} \wedge dx^{j+1} \wedge \cdots \wedge dx^n$ , and observe that  $d\omega_{j\sigma} = D_j(a_{\sigma}) dx^1 \wedge \cdots \wedge dx^n = (D_j a)_{\sigma} dx^1 \wedge \cdots \wedge dx^n$ , so (1) with  $D_j a$  in place of  $a$  implies

$$(2) \quad \int D_j a \theta^{(\sigma)} d\mathcal{L}^n = T(d\omega_{j\sigma}) = \partial T(\omega_{j\sigma}) = 0, \quad j = 1, \dots, n.$$

Observe that with  $\xi \in B_{\rho/2}(x_0)$  and the choice  $a(x) = \varphi^{(\tau)}(x - \xi)$  with  $\tau < \rho/2$  this in particular says that  $D_j(\theta^{(\sigma)})_{\tau}(\xi)$  is zero for  $\xi \in B_{\rho/2}(x_0)$  and for  $j = 1, \dots, n$ , and

hence  $(\theta^{(\sigma)})_\tau$  is constant on  $B_{\rho/2}(x_0)$ . Thus letting  $\tau \downarrow 0$  we have that  $\theta^{(\sigma)}$  is constant on  $B_{\rho/2}(x_0)$ . Finally letting  $\sigma \downarrow 0$  in (1) we conclude there is a constant  $c$  such that  $T(\omega) = c \llbracket U \rrbracket(\omega)$  if  $\omega \in \mathcal{D}^n(U)$  with  $\text{spt } \omega \subset B_{\rho/2}(x_0)$ . In view of the arbitrariness of  $B_\rho(x_0)$  this completes the proof.  $\square$

**2.42 Remark:** Notice that if we merely have  $\mathbb{M}_W(\partial T) < \infty$  for each  $W \subset\subset U$  rather than  $\partial T = 0$  then the above proof still gives a bounded measurable function  $\theta^{(\sigma)}$  as in (1), but now instead of (2) we get only that

$$\left| \int D_j a \theta^{(\sigma)} d\mathcal{L}^n \right| \leq C \sup |a| \mathbb{M}_{\check{B}_\rho(x_0)}(\partial T), \quad a \in C_c^\infty(\check{B}_\rho(x_0)),$$

with  $C$  independent of  $\sigma$ . We claim that  $\theta^{(\sigma_k)}$  is convergent in  $L^1(B)$  for some sequence  $\sigma_k \downarrow 0$ ,  $B = B_\rho(x_0)$ . Indeed by 2.7 of Ch.2, since  $\partial T$  has locally finite mass in  $U$ , there are constants  $\lambda_k$  such that  $\theta^{(\sigma_k)} - \lambda_k$  is bounded in  $L^1(B_\rho(x_0))$ , and hence  $T_{\sigma_k} - \lambda_k \llbracket B \rrbracket$  has bounded mass in  $\check{B}_\rho(x_0)$ , where  $T_\sigma(adx^1 \wedge \cdots \wedge dx^n) = T(a_\sigma dx^1 \wedge \cdots \wedge dx^n)$  for  $a \in C_c^\infty(\check{B}_\rho(x_0))$ . But  $T_{\sigma_k} \rightarrow T$  in  $\check{B}_\rho(x_0)$  and hence  $\{\lambda_k\}$  is bounded. Thus (see §2 of Ch.2 and in particular 2.6 of Ch.2) we deduce that  $\theta^{(\sigma_k)} \rightarrow \theta$  in  $L^1_{\text{loc}}(U)$  for some sequence  $\sigma_k \downarrow 0$ , with  $\theta \in BV_{\text{loc}}(U)$ , and

$$(\ddagger) \quad T(\omega) = \int a \theta d\mathcal{L}^n, \quad \omega = a dx^1 \wedge \cdots \wedge dx^n \in \mathcal{D}^n(\check{B}_\rho(x_0)).$$

Using the definition of  $\mathbb{M}(\partial T)$ , we easily then check that  $\mathbb{M}_W(\partial T) = |D\theta|(W)$  for each open  $W \subset U$  (and  $\mathbb{M}_W(T) = \int_W |\theta| d\mathcal{L}^n$ ). Indeed in the present case  $n = P$ , any  $\omega \in \mathcal{D}^{n-1}(U)$  can be written  $\omega = \sum_{j=1}^n (-1)^j a_j dx^1 \wedge \cdots \wedge dx^{j-1} \wedge dx^{j+1} \wedge \cdots \wedge dx^n$  for suitable  $a_j \in C_c^\infty(U)$ , and  $d\omega = \text{div } \underline{a} dx^1 \wedge \cdots \wedge dx^n$  for such  $\omega$  ( $\underline{a} = (a_1, \dots, a_n)$ ). Therefore by  $(\ddagger)$  above we have

$$\partial T(\omega) = T(d\omega) = \int \text{div } \underline{a} \theta d\mathcal{L}^n$$

and the assertion  $\mathbb{M}_W(\partial T) = |D\theta|(W)$  then follows directly from the definition of  $\mathbb{M}_W(\partial T)$  and  $|D\theta|$  (in §2 of Ch.2).

**2.43 Theorem.** Suppose  $U$  is open in  $\mathbb{R}^P$ , and let  $T \in \mathcal{D}_n(U)$  with  $\mathbb{M}_W(T), \mathbb{M}_W(\partial T) < \infty$  for every  $W \subset\subset U$ . Then  $\mu_T$  is absolutely continuous with respect to  $\mathcal{H}^n$  on  $U$ . That is  $\mathcal{H}^n(E) = 0 \Rightarrow \mu_T(E) = 0$ , hence (by the abstract Radon-Nikodym theorem 4.17) we have  $\mu_T = \mathcal{H}^n \llcorner \theta$  for some non-negative Borel measurable function  $\theta$  on  $U$ .

**Proof of 2.43:** Take any  $E \subset U$  with  $\mathcal{H}^n(E) = 0$ . Since there is a Borel set  $B \supset E$  with  $\mathcal{H}^n(B) = 0$ , we can assume  $E$  is Borel. We have to show  $\mu_T(E) = 0$ . Since

$\mu_T(E) = \sup_{K \subset E, K \text{ compact}} \mu_T(K)$  by 1.22 of Ch. 1, we can (and we shall) without loss of generality assume  $E$  is compact.

In the proof we use the notation that, for  $\alpha = (i_1, \dots, i_n) \in I_{n,P}$ ,  $p_\alpha$  denotes the orthogonal projection of  $\mathbb{R}^P$  onto  $\mathbb{R}^n$  given by

$$(x^1, \dots, x^P) \mapsto (x^{i_1}, \dots, x^{i_n}).$$

If  $\omega \in \mathcal{D}^n(U)$  then  $\omega = \sum_{\alpha \in I_{n,P}} \omega_\alpha dx^\alpha$ ,  $\omega_\alpha \in C_c^\infty(U)$ , and hence

$$\begin{aligned} T(\omega) &= \sum_\alpha T(\omega_\alpha dx^\alpha) = \sum_\alpha (T \lrcorner \omega_\alpha)(dx^\alpha) \\ &= \sum_\alpha (T \lrcorner \omega_\alpha) p_\alpha^\# dy. \end{aligned}$$

( $dy = dy^1 \wedge \dots \wedge dy^n$ ,  $y^1, \dots, y^n$  the standard coordinate functions in  $\mathbb{R}^n$ .) Thus

$$(1) \quad T(\omega) = \sum_\alpha p_{\alpha\#}(T \lrcorner \omega_\alpha)(dy)$$

(which makes sense because  $\text{spt}(T \lrcorner \omega_\alpha) \subset \text{spt} \omega_\alpha = \text{a compact subset of } U$ ). Now observe that  $\mathbb{M}(\partial(T \lrcorner \omega_\alpha)) < \infty$ , because, for any  $\eta \in \mathcal{D}^{n-1}(U)$ ,

$$\begin{aligned} \partial(T \lrcorner \omega_\alpha)(\eta) &= (T \lrcorner \omega_\alpha)(d\eta) \\ &= T(\omega_\alpha d\eta) \\ &= T(d(\omega_\alpha \eta)) - T(d\omega_\alpha \wedge \eta) \\ &= \partial T(\omega_\alpha \eta) - T(d\omega_\alpha \wedge \eta), \end{aligned}$$

so

$$\mathbb{M}_W(\partial(T \lrcorner \omega_\alpha)) \leq \mathbb{M}_W(\partial T)|\omega_\alpha| + \mathbb{M}_W(T)|d\omega_\alpha| < \infty$$

as claimed.

Of course then  $\mathbb{M}(\partial p_{\alpha\#} \partial(T \lrcorner \omega_\alpha)) = \mathbb{M}(p_{\alpha\#} \partial(T \lrcorner \omega_\alpha)) < \infty$ , and hence by 2.42(‡) we have a  $\theta_\alpha \in L^1(p_\alpha(U))$  (depending on both  $\alpha$  and  $\omega_\alpha$ ) such that

$$p_{\alpha\#}(T \lrcorner \omega_\alpha)(\eta) = \int_{p_\alpha(U)} \langle \eta, e_1 \wedge \dots \wedge e_n \rangle \theta_\alpha d\mathcal{L}^n,$$

and hence  $p_{\alpha\#}(T \lrcorner \omega_\alpha) \lrcorner p_\alpha(E) = 0$  because  $\mathcal{L}^n(p_\alpha(E)) \leq \mathcal{H}^n(E) = 0$ . Then

$$\begin{aligned} (2) \quad \mathbb{M}(p_{\alpha\#}(T \lrcorner \omega_\alpha)) &\leq \mathbb{M}(p_{\alpha\#}(T \lrcorner \omega_\alpha) \lrcorner (\mathbb{R}^n \setminus p_\alpha(E))) \\ &= \mathbb{M}(p_{\alpha\#}((T \lrcorner \omega_\alpha) \lrcorner (\mathbb{R}^P \setminus p_\alpha^{-1} p_\alpha(E)))) \quad (\text{by 2.30(2)}) \\ &\leq C \mathbb{M}((T \lrcorner \omega_\alpha) \lrcorner (\mathbb{R}^P \setminus p_\alpha^{-1} p_\alpha(E))) \quad (\text{by 2.30(1)}) \\ &\leq C \mathbb{M}_W(T \lrcorner (\mathbb{R}^P \setminus p_\alpha^{-1} p_\alpha(E))) \sup_W |\omega_\alpha| \end{aligned}$$

for any  $W$  such that  $\text{spt} \omega \subset W \subset\subset U$ , where  $C = \binom{P}{n}^{1/2}$ . Since  $E \subset p_\alpha^{-1} p_\alpha E$ , we have thus proved

$$\mathbb{M}(p_{\alpha\#}(T \lrcorner \omega_\alpha)) \leq C \mathbb{M}_W(T \lrcorner (\mathbb{R}^P \setminus E)) \sup_W |\omega_\alpha| \leq C \mathbb{M}_W(T \lrcorner (\mathbb{R}^P \setminus E)) |\omega|.$$

Using this in (1) we then have

$$|T(\omega)| \leq C \mathbb{M}_W(T \llcorner (\mathbb{R}^P \setminus E)) \sup_W |\omega|$$

for suitable  $C = C(P, n)$  and  $\forall \omega \in \mathcal{D}^n(U)$  with  $\text{spt } \omega \subset W$ , so

$$\mathbb{M}_W(T) \leq C \mathbb{M}_W(T \llcorner (\mathbb{R}^P \setminus E)),$$

which says

$$(3) \quad \mu_T(W) \leq C \mu_T(W \setminus E).$$

Since  $E$  is compact, we can choose  $\{W_q\}$  so that  $W_q \subset\subset U$ ,  $W_{q+1} \subset W_q$ ,  $\cap_{q=1}^{\infty} W_q = E$ ; using (3) with  $W = W_q$  then gives  $\mu_T(E) = 0$ .  $\square$

In the following corollary, we continue to use the notation that, for  $\alpha = (i_1, \dots, i_n) \in I_{n,P}$ ,  $p_\alpha$  denotes the orthogonal projection of  $\mathbb{R}^P$  onto  $\mathbb{R}^n$  given by

$$(x^1, \dots, x^P) \mapsto (x^{i_1}, \dots, x^{i_n}).$$

Observe that the proof of Theorem 2.43 used only the fact that  $\mathcal{L}^n(p_\alpha E) = 0$  for each  $\alpha \in I_{n,P}$ , hence we have the following corollary:

**2.44 Corollary.** *Suppose  $T \in \mathcal{D}_n(U)$  with  $\mathbb{M}_W(T), \mathbb{M}_W(\partial T) < \infty$  for every  $W \subset\subset U$ , and suppose  $E$  is a closed subset of  $U$  with  $\mathcal{L}^n(p_\alpha(E)) = 0$  for each multi-index  $\alpha \in I_{n,P}$ ,  $1 \leq i_1 < i_2 < \dots < i_n \leq P$ . Then  $T \llcorner E = 0$ .*

**2.45 Remarks:** (1) The hypothesis  $\mathcal{L}^n(p_\alpha(E)) = 0 \forall \alpha$  can be satisfied even if  $E$  has positive  $\mathcal{H}^n$ -measure (as example 3.4 of Ch.3 shows), but, in the case when  $E$  is  $\mathcal{H}^n$   $\sigma$ -finite, only if  $E$  is purely  $n$ -unrectifiable, as shown by 3.3 of Ch.3.

(2) Let  $Q$  be any orthogonal transformation of  $\mathbb{R}^P$ . Since  $T \in \mathcal{D}_n(U) \Rightarrow Q_\# T \in \mathcal{D}_n(QU)$  and  $\mathbb{M}_W(T) = \mathbb{M}_{QW}(Q_\# T)$  for each  $W \subset U$ . So if  $\mathbb{M}_W(T) < \infty$  for each  $W \subset\subset U$  we have  $\mu_{Q_\# T}(Q(A)) = \mu_T(A)$  for each  $A \subset U$ , hence the above lemma guarantees  $\mathcal{L}^n(p_\alpha(Q(E))) = 0$  for each  $\alpha \Rightarrow \mu_T(E) = 0$ . On the other hand the Rectifiability Theorem 3.7 of Ch.3 implies that if  $E$  is a  $\mathcal{H}^n$   $\sigma$ -finite set which is purely  $n$ -unrectifiable then almost all (with respect to Haar measure) orthogonal projections  $p$  of  $\mathbb{R}^{n+k}$  onto an  $n$ -dimensional subspace of  $\mathbb{R}^{n+k}$  have  $\mathcal{H}^n(p(E)) = 0$ . But for each  $\alpha \in I_{n,P}$  any such orthogonal projection  $p$  can be expressed  $p = Q^* \circ \tilde{p}_\alpha \circ Q$  for some orthogonal transformation  $Q$  of  $\mathbb{R}^{n+k}$ , where  $\tilde{p}_\alpha(x) = (p_\alpha(x), 0) \in \mathbb{R}^{n+k}$ . Hence there must be many orthogonal  $Q$  such that  $\mathcal{L}^n(p_\alpha(Q(E))) = 0$  for each  $\alpha \in I_{n,P}$ , so, by applying Corollary 2.44 with  $Q_\# T$  in place of  $T$ , we conclude  $T \llcorner E = 0$  for any purely  $n$ -unrectifiable  $\mathcal{H}^n$   $\sigma$ -finite set  $E \subset U$ .

### 3 Integer Multiplicity Rectifiable Currents

In this section we want to develop the theory of integer multiplicity currents  $T \in \mathcal{D}_n(U)$ , which, roughly speaking are those currents obtained by assigning (in a  $\mathcal{H}^n$ -measurable fashion) an orientation to the tangent spaces  $T_x V$  of an integer multiplicity varifold  $V$ . (See Ch. 4 for terminology.)

These currents are precisely those called locally *locally rectifiable currents* by Federer and Fleming [FF60], [Fed69].

Throughout this section  $n \geq 1, k \geq 1$  are integers and  $U$  is an open subset of  $\mathbb{R}^{n+k}$ .

**3.1 Definition:** If  $T \in \mathcal{D}_n(U)$  we say that  $T$  is an integer multiplicity rectifiable  $n$ -current (briefly an integer multiplicity current) if it can be expressed

$$(\ddagger) \quad T(\omega) = \int_M \langle \omega(x), \xi(x) \rangle \theta(x) d\mathcal{H}^n(x), \quad \omega \in \mathcal{D}^n(U),$$

where  $M$  is an  $\mathcal{H}^n$ -measurable countably  $n$ -rectifiable subset of  $U$ ,  $\theta$  is a positive locally  $\mathcal{H}^n$ -integrable function which is integer-valued  $\mathcal{H}^n$ -a.e., and  $\xi : M \rightarrow \Lambda_n(\mathbb{R}^{n+k})$  is a  $\mathcal{H}^n$ -measurable function such that for  $\mathcal{H}^n$ -a.e. point  $x \in M$ ,  $\xi(x)$  can be expressed in the form  $\tau_1 \wedge \cdots \wedge \tau_n$ , where  $\tau_1, \dots, \tau_n$  form an orthonormal basis for the approximate tangent space  $T_x M$ . (See Ch. 3 and Ch. 4.) Thus  $\xi (= \vec{T})$  orients the approximate tangent spaces of  $M$  in an  $\mathcal{H}^n$ -measurable way. The function  $\theta$  in 3.1 ( $\ddagger$ ) is called the *multiplicity* and  $\xi$  is called the *orientation* for  $T$ . If  $T$  is as in 3.1 ( $\ddagger$ ) we shall often write

$$T = \underline{\tau}(M, \theta, \xi).$$

In this case

$$V = \underline{v}(M, \theta)$$

will be referred to as the *integer multiplicity varifold associated with  $T$* . In case  $\theta = 1$   $\mathcal{H}^n$ -a.e. on  $M$  we use the abbreviated notation

$$T = \underline{\tau}(M, \xi), \quad V = \underline{v}(M).$$

**3.2 Remarks:** (1) If  $T_1, T_2 \in \mathcal{D}_n(U)$  are integer multiplicity, then so is  $p_1 T_1 + p_2 T_2$ ,  $p_1, p_2 \in \mathbb{Z}$ .

(2) If  $T_1 = \underline{\tau}(M_1, \theta_1, \xi_1) \in \mathcal{D}_r(U)$ ,  $T_2 = \underline{\tau}(M_2, \theta_2, \xi_2) \in \mathcal{D}_s(W)$  ( $W \subset \mathbb{R}^Q$  open), then  $T_1 \times T_2 \in \mathcal{D}_{r+s}(U \times W)$  is also integer multiplicity, and in fact

$$T_1 \times T_2 = \underline{\tau}(M_1 \times M_2, \theta_1 \theta_2, p_{\#}(\xi_1) \wedge q_{\#}(\xi_2)),$$

where  $p(x) = (x, 0)$  and  $q(y) = (0, y)$  and  $(\theta_1 \theta_2)(x, y) = \theta_1(x) \theta_2(y)$ .

(3) If  $T = \tau(M, \theta, \xi) \in \mathcal{D}_n(U)$  is an integer multiplicity current then

$$\mathbb{M}_W(T) = \int_M \theta \, d\mathcal{H}^n = \mathbb{M}_W(V) \quad \forall \text{ open } W \subset U,$$

where  $V = \underline{v}(M, \theta)$  is the rectifiable varifold associated with  $T$ .

Next we want to discuss pushing forward an integer multiplicity  $T = \tau(M, \theta, \xi) \in \mathcal{D}_n(U)$  ( $M \subset U$ ) by a Lipschitz map  $f : U \rightarrow W$  such that  $f|_{\text{spt } T}$  is proper. First, if  $f$  is  $C^1$ , 1:1,  $f|_{\text{spt } T}$  is proper,  $M$  is an embedded  $C^1$  submanifold,  $\xi$  is any  $\mathcal{H}^n$ -measurable orientation for  $M$ , and  $\theta$  is any  $\mathcal{H}^n$ -measurable positive integer valued function on  $M$ , then we have, by Remark 2.30(3),

$$\begin{aligned} 3.3 \quad f_{\#}T(\omega) &= \int_M \langle f^{\#}\omega, \xi \rangle \theta \, d\mathcal{H}^n \\ &= \int_M \langle (df_x)^{\#}(\omega|_{f(x)}), \xi|_x \rangle \theta(x) \, d\mathcal{H}^n(x) \\ &= \int_M \langle \omega|_{f(x)}, (df_x)_{\#}(\xi|_x) \rangle \theta(x) \, d\mathcal{H}^n(x). \end{aligned}$$

Now  $\xi|_x = \pm \tau_1 \wedge \cdots \wedge \tau_n$ , where  $\tau_1, \dots, \tau_n$  is an orthonormal basis for the tangent space  $T_x M$ , so

$$\begin{aligned} 3.4 \quad df_{x\#}\xi|_x &= \pm df_{x\#}\tau_1 \wedge \cdots \wedge df_{x\#}\tau_n \\ &= \pm D_{\tau_1} f(x) \wedge \cdots \wedge D_{\tau_n} f(x) \end{aligned}$$

which = 0 at points  $x \in M$  where  $J_f^M(x) = 0$ , because  $J_f^M(x) = 0 \Leftrightarrow \text{rank}(d^M f_x) < n$ . On the other hand at points where  $J_f^M(x) \neq 0$  the rank is  $n$  and hence there is  $\rho > 0$  such that  $f|M \cap \check{B}_{\rho}(x)$  is a diffeomorphism onto an  $n$ -dimensional embedded  $C^1$  manifold  $N$ , and at the image point  $y = f(x)$  we let  $\eta_1, \dots, \eta_n$  be an orthonormal basis for  $T_y N$ . Then, since  $D_{\tau_i} f(x) \in T_y N$ , we have  $D_{\tau_i} f(x) = \sum_{j=1}^n D_{\tau_i} f \cdot \eta_j \eta_j$ , and so

$$D_{\tau_1} f(x) \wedge \cdots \wedge D_{\tau_n} f(x) = \det(D_{\tau_i} f \cdot \eta_j) \eta_1 \wedge \cdots \wedge \eta_n.$$

On the other hand

$$\begin{aligned} J_f^M(x) &= \sqrt{\det(D_{\tau_i} f(x) \cdot D_{\tau_j} f(x))} = \sqrt{(\det(D_{\tau_i} f(x) \cdot \eta_j))^2} \\ &= |\det(D_{\tau_i} f(x) \cdot \eta_j)|. \end{aligned}$$

Thus we see that 3.4 implies, at points  $x \in M$  where  $J_f^M(x) \neq 0$ ,

$$3.5 \quad df_{x\#}\xi|_x = J_f^M(x) \eta,$$

where  $\eta$  is an orienting  $n$ -vector for  $N$  (so  $\eta = \pm \eta_1 \wedge \cdots \wedge \eta_n$ ).  $\eta$  is called *the orientation for  $N$  induced by  $f$*  at each point  $x$  where  $J_f^M(x) \neq 0$ .

Now suppose  $f : U \rightarrow W$  is Lipschitz,  $T = \tau(M, \theta, \xi) \in \mathcal{D}_n(U)$  ( $M \subset U$ ) is an integer multiplicity current, and  $f|_{\text{spt } T}$  is proper, then we can define  $f_\#T \in \mathcal{D}_n(W)$  by

$$f_\#T(\omega) = \int_M \langle \omega|_{f(x)}, d^M f_{x\#} \xi(x) \rangle \theta(x) d\mathcal{H}^n(x).$$

Let  $\eta_0$  be  $\mathcal{H}^n$ -measurable such that, for  $\mathcal{H}^n$ -a.e.  $y \in f(M)$ ,  $\eta_0(y) = \pm \eta_1 \wedge \cdots \wedge \eta_n$ , where  $\eta_1, \dots, \eta_n$  are an orthonormal basis for the approximate tangent space  $T_y(f(M))$ . At  $\mathcal{H}^n$ -a.e. point  $x$  where  $J_f^M(x) \neq 0$  the above discussion shows

$$3.6 \quad d^M f_{x\#} \xi(x) = \sigma(x) \theta(x) J_f^M(x) \eta_0 \text{ where } \sigma(x) = \pm 1,$$

so by the area formula

$$3.7 \quad f_\#T(\omega) = \int_{f(M)} \langle \omega|_y, \eta_0(y) \rangle \sum_{x \in f^{-1}(y) \cap M_+} \sigma(x) \theta(x) d\mathcal{H}^n(y),$$

where  $M_+ = \{x \in M : J_M f(x) > 0\}$ . Of course since  $f(M)$  has locally finite  $\mathcal{H}^n$ -measure in  $V$  we know by the area formula  $\int_A J_f^M \theta d\mathcal{H}^n = \int_{f(A)} \sum_{x \in f^{-1}(y)} \theta(x) d\mathcal{H}^n(y)$ , so  $f^{-1}(y)$  is a finite set for  $\mathcal{H}^n$ -a.e.  $y \in f(M)$  and  $\sum_{x \in f^{-1}(y) \cap M_+} \sigma(x) \theta(x) \in \mathbb{Z}$  for  $\mathcal{H}^n$ -a.e.  $y \in f(M)$ . By replacing  $\eta_0$  be  $-\eta_0$  at all points  $y \in f(M)$  where  $\sum_{x \in f^{-1}(y) \cap M_+} \sigma(x) \theta(x) < 0$ , we get a new orientation  $\eta$  for  $f(M)$  (called *the orientation of  $f(M)$  induced by  $f$* ) such that

$$3.8 \quad f_\#T(\omega) = \int_{f(M)} \langle \omega|_y, \eta(y) \rangle N(y) d\mathcal{H}^n(y),$$

where  $\eta(y)$  is a suitable orientation for the approximate tangent space  $T_y(f(M))$  and  $N(y)$  is a non-negative integer given by

$$3.9 \quad N(y) = \left| \sum_{x \in f^{-1}(y) \cap M_+} \sigma(x) \theta(x) \right|, \quad \mathcal{H}^n\text{-a.e. } y \in f(M).$$

with  $\sigma(x) = \pm 1$  according as  $d^M f_{x\#} \xi(x) = \pm J_f^M(x) \eta(y)$ . Thus for  $\mathcal{H}^n$ -a.e.  $y \in f(M)$  we have

$$3.10 \quad N(y) \leq \sum_{x \in f^{-1}(y) \cap M_+} \theta(x), \quad N(y) \equiv \sum_{x \in f^{-1}(y) \cap M_+} \theta(x) \pmod{2}.$$

Also of course

$$3.11 \quad N(y) = \theta \circ f^{-1}(y) \text{ in case } f \text{ is 1:1.}$$

Thus we have proved

**3.12 Lemma.** *If  $f : U \rightarrow W$  is locally Lipschitz and  $f|_{\text{spt } T}$  is proper, with  $T = \tau(M, \xi, \theta) \in \mathcal{D}_n(U)$  an integer multiplicity current, then  $f_\#T$  is an integer multiplicity current in  $W$ ; in fact  $f_\#T = \tau(f(M), \eta, N)$ , as in 3.8, 3.9 above.*

**3.13 Remark:** If  $f : U \rightarrow W$  is Lipschitz and if  $V = \underline{v}(M, \theta)$  is the varifold associated with  $T = \underline{\tau}(M, \xi, \theta)$ , then

$$\mu_{f\#T} \leq \mu_{f\#V}$$

(in the sense of measures) with equality if and only if, for  $\mathcal{H}^n$ -a.e.  $y \in f(M)$ , the sign  $\sigma(x)$  in 3.6 above remains constant as  $x$  varies over  $f^{-1}(y) \cap M_+$ , which is the same as saying  $N(y)$  (in 3.9) satisfies  $N(y) = \sum_{x \in f^{-1}(y) \cap M} \theta(x)$  for  $\mathcal{H}^n$ -a.e.  $y \in f(M)$ . In particular we have  $\mu_{f\#T} = \mu_{f\#V}$  in case  $f$  is 1:1.

Notice also that, by applying 3.12 to the current  $R = 0 \times T$  in 2.39, we have

3.14  $U$  star-shaped from 0,  $T$  integer multiplicity in  $U$ ,  $\text{spt } T$  compact,  $\partial T = 0$   
 $\Rightarrow \exists$  an integer multiplicity  $R$  with  $\partial R = T$ ,  $\text{spt } R$  compact.

A fact of central importance concerning integer multiplicity currents is the following compactness theorem, first proved by Federer and Fleming [FF60]:

**3.15 Theorem. (Federer-Fleming Compactness Theorem.)** *If  $\{T_j\} \subset \mathcal{D}_n(U)$  is a sequence of integer multiplicity currents with*

$$(x) < \sup_{j \geq 1} (\mathbb{M}_W(T_j) + \mathbb{M}_W(\partial T_j)) < \infty \quad \forall W \subset\subset U,$$

*then there is an integer multiplicity  $T \in \mathcal{D}_n(U)$  and a subsequence  $\{T_{j'}\}$  such that  $T_{j'} \rightarrow T$  in  $U$ .*

We shall give the proof of this in §8. Notice that the existence of a  $T \in \mathcal{D}_n(U)$  and a subsequence  $\{T_{j'}\}$  with  $T_{j'} \rightarrow T$  is a consequence of the elementary 2.17; only the fact that  $T$  is an integer multiplicity rectifiable current is non-trivial.

**3.16 Remark:** Notice that the proof of 3.15 in the codimension 1 case (when  $P = n$ ) is a direct consequence of Remark 2.42 and the Compactness Theorem for BV functions (§ 2.6 of Ch. 2).

In contrast to the difficulty in proving 3.15, it is quite straightforward to prove that if  $T_j$  converges to  $T$  in the strong sense that  $\lim \mathbb{M}_W(T_j - T) = 0 \quad \forall W \subset\subset U$ , and if  $T_j$  are integer multiplicity  $\forall j$ , then  $T$  is integer multiplicity. Indeed we have the following lemma.

**3.17 Lemma.** *The set of integer multiplicity currents in  $\mathcal{D}_n(U)$  is complete with respect to the topology given by the family  $\{\mathbb{M}_W\}_{W \subset\subset U}$  of semi-norms.*

**Proof:** Let  $\{T_Q\}$  be a sequence of integer multiplicity currents in  $\mathcal{D}_n(U)$  and  $\{T_Q\}$  is Cauchy with respect to the semi-norms  $\mathbb{M}_W$ ,  $W \subset\subset U$ . Suppose

$$T_Q = \underline{\tau}(M_Q, \theta_Q, \xi_Q)$$

( $\theta_Q$  positive integer-valued on  $M_Q$ ,  $M_Q$  countably  $n$ -rectifiable,  $\mathcal{H}^n(M_Q \cap W) < \infty$  for each  $W \subset\subset U$ ). Then

$$(1) \quad \mathbb{M}_W(T_Q - T_P) \equiv \int_W |\theta_P \xi_P - \theta_Q \xi_Q| d\mathcal{H}^n < \varepsilon_W(Q)$$

$\forall P \geq Q$ , where  $\varepsilon_W(Q) \downarrow 0$  as  $Q \rightarrow \infty$  and where we adopt the convention  $\xi_P = 0$ ,  $\theta_P = 0$  on  $U \setminus M_P$ . In particular, since  $|\xi_P| = 1$  on  $M_P$ , we get

$$(2) \quad \int_W |\theta_P - \theta_Q| d\mathcal{H}^n < \varepsilon_W(Q) \quad \forall P \geq Q,$$

and hence  $\theta_P$  converges in  $L^1(\mathcal{H}^n)$  locally in  $U$  to an integer-valued function  $\theta$ . Of course (2) implies

$$(3) \quad \mathcal{H}^n(((M_+ \setminus M_Q) \cup (M_Q \setminus M_+)) \cap W) \leq \varepsilon_W(Q),$$

where  $M_+ = \{x \in U : \theta(x) > 0\}$ . (1), (2) also imply

$$\int_W \theta_P |\xi_P - \xi_Q| d\mathcal{H}^n \leq 2\varepsilon_W(Q) \quad \forall P \geq Q,$$

and hence by (3)  $\xi_P$  converges in  $L^1(\mathcal{H}^n)$  locally in  $U$  to a function  $\xi$  with values in  $\Lambda_n(\mathbb{R}^{n+k})$  with  $|\xi| = 1$  and  $\xi$  simple on  $M_+$ .

Now  $\xi_q(x) \in \Lambda_n(T_x M_Q)$ ,  $\mathcal{H}^n$ -a.e.  $x \in M_Q$ , and (by (3))  $T_x M_+ = T_x M_Q$  except for a set of measure  $\leq \varepsilon_W(Q)$  in  $M_+ \cap W$ . It follows that  $\xi(x) \in \Lambda_n(T_x M_+)$  for  $\mathcal{H}^n$ -a.e.  $x \in M_+$  and we have shown that  $\mathbb{M}_W(T_P - T) \rightarrow 0$ , where  $T = \underline{T}(M_+, \theta, \xi)$  is an integer multiplicity  $n$ -current in  $U$ .  $\square$

Finally, we shall need the following useful *decomposition theorem* for codimension 1 integer multiplicity currents.

**3.18 Theorem.** *Suppose  $P = n + 1$  (i.e.  $U$  is open in  $\mathbb{R}^{n+1}$ ) and  $R$  is an integer multiplicity current in  $\mathcal{D}_{n+1}(U)$  with  $\mathbb{M}_W(\partial R) < \infty \quad \forall W \subset\subset U$ . Then  $T = \partial R$  is integer multiplicity, and in fact we can find a decreasing sequence of  $\mathcal{L}^{n+1}$ -measurable sets  $\{U_j\}_{j=-\infty}^\infty$  of locally finite perimeter in  $U$  such that (in  $U$ )*

$$\begin{aligned} R &= \sum_{j=1}^\infty \llbracket U_j \rrbracket - \sum_{j=-\infty}^0 \llbracket V_j \rrbracket, \quad V_j = U \setminus U_j, \quad j \leq 0, \\ T &= \sum_{j=-\infty}^\infty \partial \llbracket U_j \rrbracket, \\ \mu_T &= \sum_{j=-\infty}^\infty \mu_{\partial \llbracket U_j \rrbracket}, \end{aligned}$$

and in particular

$$\mathbb{M}_W(T) = \sum_{j=-\infty}^\infty \mathbb{M}_W(\partial \llbracket U_j \rrbracket) \quad \forall W \subset\subset U.$$

**3.19 Remark:** Let  $*$  :  $C_c^\infty(U; \mathbb{R}^{n+1}) \rightarrow \mathcal{D}^n(U)$  be defined by

$$*g = \sum_{j=1}^{n+1} (-1)^{j-1} g_j dx^1 \wedge \cdots \wedge dx^{j-1} \wedge dx^{j+1} \wedge \cdots \wedge dx^{n+1},$$

so that  $d * g = \operatorname{div} g dx^1 \wedge \cdots \wedge dx^{n+1}$ . Then for any  $\mathcal{L}^{n+1}$ -measurable  $A \subset U$  we have

$$\begin{aligned} \partial \llbracket A \rrbracket (*g) &= \llbracket A \rrbracket (d * g) \\ &= \int_U \chi_A \operatorname{div} g d\mathcal{L}^{n+1}, \end{aligned}$$

and hence by definition of  $|D\chi_A|$  (in §2 of Ch. 2) and  $\mathbb{M}(T)$  (in §2 of the present chapter) we see that

$$A \text{ has locally finite perimeter in } U \iff \mathbb{M}_W(\partial \llbracket A \rrbracket) < \infty \quad \forall W \subset\subset U,$$

and in this case

$$\begin{cases} \mathbb{M}_W(\partial \llbracket A \rrbracket) = \int_W |D\chi_A| \quad \forall W \subset\subset U \\ \overrightarrow{\partial \llbracket A \rrbracket} = *v_A, \quad |D\chi_A| \text{ a.e. in } U. \end{cases}$$

Here  $v_A$  is the inward unit normal function for  $A$  (defined on the reduced boundary  $\partial^*A$  as in 4.4 of Ch. 3).

**Proof of 3.18:**  $R$  must have the form

$$R = \underline{\tau}(V, \theta, \xi),$$

where  $V$  is an  $\mathcal{L}^{n+1}$ -measurable subset of  $U$  and  $\xi(x) = \pm e_1 \wedge \cdots \wedge e_{n+1}$  for each  $x \in V$ . Thus letting

$$\tilde{\theta}(x) = \begin{cases} \theta(x) & \text{when } x \in V \text{ and } \xi(x) = +e_1 \wedge \cdots \wedge e_{n+1} \\ -\theta(x) & \text{when } x \in V \text{ and } \xi(x) = -e_1 \wedge \cdots \wedge e_{n+1} \\ 0 & \text{when } x \notin V, \end{cases}$$

we have

$$(1) \quad R(\omega) = \int_V a \tilde{\theta} d\mathcal{L}^{n+1},$$

$\omega = a dx^1 \wedge \cdots \wedge dx^{n+1} \in \mathcal{D}^{n+1}(U)$  and (cf. 2.10)

$$(2) \quad \mathbb{M}_W(R) = \int_W |\tilde{\theta}| d\mathcal{L}^{n+1}, \quad \mathbb{M}_W(T) = \int_W |D\tilde{\theta}| \quad \forall W \subset\subset U$$

(and  $\tilde{\theta} \in BV_{\text{loc}}(U)$ ).

Define

$$U_j = \left\{ x \in U : \tilde{\theta}(x) \geq j \right\}, \quad j \in \mathbb{Z}$$

$$V_j = U \setminus U_j = \left\{ x \in U : \tilde{\theta}(x) \leq -1 - j \right\}, \quad j \leq 0.$$

Then one checks directly that

$$\tilde{\theta} = \sum_{j=1}^{\infty} \chi_{U_j} - \sum_{j=-\infty}^0 \chi_{V_j}$$

( $\chi_A$  = indicator function of  $A$ ,  $A \subset U$ ), and hence by (1)

$$(3) \quad R = \sum_{j=1}^{\infty} \llbracket U_j \rrbracket - \sum_{j=-\infty}^0 \llbracket V_j \rrbracket \text{ in } U.$$

Since  $T(\omega) = \partial R(\omega) = R(d\omega)$ ,  $\omega \in \mathcal{D}^n(U)$ , we then have

$$(4) \quad \begin{aligned} T &= \partial R = \sum_{j=1}^{\infty} \partial \llbracket U_j \rrbracket - \sum_{j=-\infty}^0 \partial \llbracket V_j \rrbracket \\ &= \sum_{j=-\infty}^{\infty} \partial \llbracket U_j \rrbracket, \end{aligned}$$

so we have the required decomposition, and it remains only to prove that each  $U_j$  has locally finite perimeter in  $U$  and that the corresponding measures add. To check this, take  $\psi_j \in C^1(\mathbb{R})$  with

$$\begin{cases} \psi_j(t) = 0 & \text{for } t \leq j - 1 + \varepsilon, \quad \psi_j(t) = 1, \quad t \geq j - \varepsilon \\ 0 \leq \psi_j \leq 1, \quad \sup |\psi_j'| \leq 1 + 3\varepsilon, \end{cases}$$

where  $\varepsilon \in (0, \frac{1}{2})$ . Then if  $a \in C_c^\infty(U)$  and  $g = (g^1, \dots, g^{n+1})$ ,  $g^j \in C_c^\infty(U)$ , with  $|g| \leq a$ , we have (since  $\chi_{U_j} = \psi_j \circ \tilde{\theta} \forall j$ ) that for any  $M \leq N$

$$(5) \quad \begin{aligned} \int_U \operatorname{div} g \sum_{j=M}^N \chi_{U_j} d\mathcal{L}^{n+1} &= \int_U \operatorname{div} g \sum_{j=M}^N \psi_j \circ \tilde{\theta} d\mathcal{L}^{n+1} \\ &= \lim_{\sigma \downarrow 0} \int_U \operatorname{div} g \sum_{j=M}^N \psi_j \circ \tilde{\theta}_\sigma d\mathcal{L}^{n+1} \\ &= - \lim_{\sigma \downarrow 0} \int_U g \cdot \nabla \tilde{\theta}_\sigma \psi_j'(\tilde{\theta}_\sigma) d\mathcal{L}^{n+1} \\ &\leq (1 + 3\varepsilon) \lim_{\sigma \downarrow 0} \int_U a |\nabla \tilde{\theta}_\sigma| d\mathcal{L}^{n+1}. \end{aligned}$$

On the other hand

$$(6) \quad \begin{cases} \int_U a |\nabla \tilde{\theta}_\sigma| = \sup_{g \in C_c^1(U), |g| \leq a} \int_U \operatorname{div} g \tilde{\theta}_\sigma d\mathcal{L}^n, \text{ and} \\ \int_U \operatorname{div} g \tilde{\theta}_\sigma d\mathcal{L}^n = \int_U \operatorname{div} g_\sigma \tilde{\theta} d\mathcal{L}^n = R(d\omega_\sigma) = T(\omega_\sigma) \leq \mathbb{M}(T)|\omega|, \end{cases}$$

where  $\omega_\sigma = \sum_{j=1}^n (-1)^{j-1} g_{j\sigma} dx^1 \wedge \cdots \wedge dx^{j-1} \wedge dx^{j+1} \wedge \cdots \wedge dx^n$ . Thus, taking  $M = N$ , we deduce from (5) and (6) that  $\mathbb{M}_W(\partial[U_j]) \leq \mathbb{M}_W(T) < \infty$  for each  $j$  and each open  $W \subset\subset U$ .

By taking  $M = -N$  in (5) and defining  $R_N = \sum_{j=1}^N \llbracket U_j \rrbracket - \sum_{j=-N}^0 \llbracket V_j \rrbracket$  we see that (with  $g$  as in 3.19)

$$|R_N(d * g)| \leq (1 + 3\varepsilon) \int_U a \, d\mu_T,$$

and hence, with  $T_N = \partial R_N$ ,

$$(7) \quad \int_U a \, d\mu_{T_N} \leq \int_U a \, d\mu_T \quad \forall N \geq 1,$$

$a \geq 0, a \in C_c^\infty(U)$ . On the other hand by 4.1 of Ch.3 we have

$$(8) \quad \begin{aligned} R_N(d * g) &= \sum_{j=-N}^N \int_U \operatorname{div} g \, \chi_{U_j} \, d\mathcal{L}^{n+1} \\ &= -\sum_{j=-N}^N \int_{\partial^* U_j} v_j \cdot g \, d\mathcal{H}^n, \end{aligned}$$

where  $v_j$  is the inward unit normal for  $U_j$  and  $\partial^* U_j$  is the reduced boundary for  $U_j$  (see §4 of Ch.3 and in particular 4.4 of Ch.3). By virtue of the fact that  $U_{j+1} \subset U_j$  we see from 4.4 (††) of Ch.3 that  $v_j \equiv v_k$  on  $\partial^* U_j \cap \partial^* U_k \, \forall j, k$ . Hence (8) can be written

$$T_N(*g) = - \int_U v \cdot g \, h_N \, d\mathcal{H}^n,$$

where  $h_N = \sum_{j=-N}^N \chi_{\partial^* U_j}$  and where  $v$  is defined on  $\cup_{j=-\infty}^\infty \partial^* U_j$  by  $v = v_j$  on  $\partial^* U_j$ . Since  $|v| \equiv 1$  on  $\cup_{j=-\infty}^\infty \partial^* U_j$  this evidently gives

$$\begin{aligned} \int_U a \, d\mu_{T_N} &= \int_U a \, h_N \, d\mathcal{H}^n \\ &= \sum_{j=-N}^N \int_{\partial^* U_j} a \, d\mathcal{H}^n \\ &= \sum_{j=-N}^N \int a \, d\mu_{\partial[U_j]}. \end{aligned}$$

Letting  $N \rightarrow \infty$  we thus have (by (7))

$$\mu_T \geq \sum_{j=-\infty}^\infty \mu_{\partial[U_j]}.$$

Since the reverse inequality follows directly from (4), the proof is complete.  $\square$

**3.20 Corollary.** *Let  $R$  be integer multiplicity  $\in \mathcal{D}_{n+1}(U)$ ,  $U \subset \mathbb{R}^P$ ,  $P \geq n+1$ , and suppose there is an  $(n+1)$ -dimensional  $C^1$  embedded submanifold  $N$  of  $\mathbb{R}^P$  with  $\operatorname{spt} R \subset N \cap U$ . Suppose further that  $T = \partial R$  and  $\mathbb{M}(T) < \infty \, \forall W \subset\subset U$ . Then  $T (\in \mathcal{D}_n(U))$  is*

integer multiplicity and for each point  $y \in N \cap U$  there is an open  $W_y \subset U$ ,  $y \in W_y$ , and  $\mathcal{H}^{n+1}$  measurable subset  $\{U_j\}_{j=-\infty}^{\infty}$  with  $U_{j+1} \subset U_j \subset N \cap U$ ,  $\mathbb{M}_{W_y}(\partial \llbracket U_j \rrbracket) < \infty \forall j$ , and with the following identities holding in  $W_y$ :

$$\begin{aligned} R &= \sum_{j=1}^{\infty} \llbracket U_j \rrbracket - \sum_{j=-\infty}^0 \llbracket U \setminus U_j \rrbracket \\ T &= \sum_{j=-\infty}^{\infty} \partial \llbracket U_j \rrbracket \\ \mu_T &= \sum_{j=-\infty}^{\infty} \mu_{\partial \llbracket U_j \rrbracket}. \end{aligned}$$

**Proof:** The proof is an easy consequence of 3.18 using local coordinate representations for  $N$ .  $\square$

## 4 Slicing

We first want to define the notion of slice for integer multiplicity currents. Preparatory to this we have the following lemma:

**4.1 Lemma.** *If  $M$  is  $\mathcal{H}^n$ -measurable, countably  $n$ -rectifiable,  $f$  is Lipschitz on  $\mathbb{R}^{n+k}$  and  $M_+ = \{x \in M : |\nabla^M f(x)| > 0\}$ , then for  $\mathcal{L}^1$ -almost all  $t \in \mathbb{R}$  the following statements hold:*

- (1)  $M_t \equiv f^{-1}(t) \cap M_+$  is countably  $\mathcal{H}^{n-1}$ -rectifiable
- (2) For  $\mathcal{H}^{n-1}$ -a.e.  $x \in M_t$ ,  $T_x M_t$  and  $T_x M$  both exist,  $T_x M_t$  is an  $(n-1)$ -dimensional subspace of  $T_x M$ , and in fact
 
$$(\ddagger) \quad T_x M = \{y + \lambda \nabla^M f(x) : y \in T_x M_t, \lambda \in \mathbb{R}\}.$$

Furthermore for any non-negative  $\mathcal{H}^n$ -measurable function  $g$  on  $M$  we have

$$\int_{-\infty}^{\infty} \left( \int_{M_t} g \, d\mathcal{H}^{n-1} \right) dt = \int_M |\nabla^M f| g \, d\mathcal{H}^n.$$

**Proof:** In fact (1) is just a restatement of 2.10(2) of Ch. 3, and (2) follows from 1.6 of Ch. 3 together with the facts that for  $\mathcal{L}^1$ -a.e.  $t \in \mathbb{R}$  and  $\mathcal{H}^{n-1}$ -a.e.  $x \in M_t$

$$\nabla^M f(x) \in T_x M \quad (\text{by definition of } \nabla^M f \text{ in §2 of Ch. 3})$$

and

$$\langle \nabla^M f(x), \tau \rangle = 0 \quad \forall \tau \in T_x M_t.$$

(This last follows for example from Definition 2.1 of Ch. 3.)

The last part of the lemma is just a restatement of the appropriate version of the co-area formula (discussed in §2 of Ch. 3).

**4.2 Remark:** Note that by replacing  $g$  (in 4.1 above) by  $g$  times the indicator function of  $\{x : f(x) < t\}$  we get the identity

$$\int_{M \cap \{f(x) < t\}} |\nabla^M f| g \, d\mathcal{H}^n = \int_{-\infty}^t \int_{M_s} g \, d\mathcal{H}^{n-1} \, ds$$

so that the left side as an absolutely continuous function of  $t$  and

$$\frac{d}{dt} \int_{M \cap \{f(x) < t\}} |\nabla^M f| g \, d\mathcal{H}^n = \int_{M_t} g \, d\mathcal{H}^{n-1}, \text{ a.e. } t \in \mathbb{R}.$$

Now let  $T = \underline{\tau}(M, \theta, \xi)$  be an integer multiplicity current in  $U$  ( $U$  open in  $\mathbb{R}^{n+k}$ ,  $M \subset U$ ), let  $f$  be Lipschitz in  $U$  and let  $\theta_+$  be defined  $\mathcal{H}^n$ -a.e. in  $M$  by

$$\theta_+(x) = \begin{cases} 0 & \text{if } \nabla^M f(x) = 0 \\ \theta(x) & \text{if } \nabla^M f(x) \neq 0. \end{cases}$$

For the ( $\mathcal{L}^1$ -almost all)  $t \in \mathbb{R}$  such that  $T_x M$ ,  $T_x M_t$  exist for  $\mathcal{H}^{n-1}$ -a.e.  $x \in M_t$  and such that 4.1 (2)( $\ddagger$ ) holds, we have

$$4.3 \quad \xi(x) \llcorner \frac{\nabla^M f(x)}{|\nabla^M f(x)|} \text{ is simple} \in \Lambda_{n-1}(T_x M_t) \subset \Lambda_{n-1}(T_x M)$$

and has unit length (for  $\mathcal{H}^{n-1}$ -a.e.  $x \in M_t$ ). Here we use the notation that if  $v \in \Lambda_n(T_x M)$  and  $w \in T_x M$ , then  $v \llcorner w \in \Lambda_{n-1}(T_x M)$  is defined by

$$\langle v \llcorner w, a \rangle = \langle v, w \wedge a \rangle, \quad a \in \Lambda_{n-1}(T_x M).$$

Using this notation we can now define the notion of a slice of  $T$  by  $f$ ; we continue to assume  $T \in \mathcal{D}_n(U)$  is given by  $T = \underline{\tau}(M, \theta, \xi)$  as above.

**4.4 Definition:** For the ( $\mathcal{L}^1$ -almost all)  $t \in \mathbb{R}$  since that  $T_x M$ ,  $T_x M_t$  exist and Lemma 4.1 (2)( $\ddagger$ ) holds  $\mathcal{H}^{n-1}$ -a.e.  $x \in M_t$ , with the notation introduced above (and bearing in mind 4.3) we define the integer multiplicity current  $\langle T, f, t \rangle \in \mathcal{D}_{n-1}(U)$  by

$$\langle T, f, t \rangle = \underline{\tau}(M_t, \theta_t, \xi_t),$$

where

$$\xi_t(x) = \xi(x) \llcorner \frac{\nabla^M f(x)}{|\nabla^M f(x)|}, \quad \theta_t = \theta_+|_{M_t}.$$

So defined,  $\langle T, f, t \rangle$  is called the slice of  $T$  by  $f$  at  $t$ .

**4.5 Lemma.** (1) For each open  $W \subset U$

$$\int_{-\infty}^{\infty} \mathbb{M}_W(\langle T, f, t \rangle) dt = \int_{M \cap W} |\nabla^M f| |\theta| d\mathcal{H}^n \leq (\text{ess sup}_{M \cap W} |\nabla^M f|) \mathbb{M}_W(T).$$

(2) If  $\mathbb{M}_W(\partial T) < \infty \forall W \subset\subset U$ , then for  $\mathcal{L}^1$ -a.e.  $t \in \mathbb{R}$

$$\langle T, f, t \rangle = \partial [T \llcorner \{f < t\}] - (\partial T) \llcorner \{f < t\}.$$

(3) If  $\partial T$  is integer multiplicity in  $\mathcal{D}_{n-1}(U)$ , then for  $\mathcal{L}^1$ -a.e.  $t \in \mathbb{R}$

$$\langle \partial T, f, t \rangle = -\partial \langle T, f, t \rangle.$$

**Proof:** (1) is a direct consequence of the last part of 4.1 (with  $g = \theta_+$ ).

To prove (2) we first recall that, since  $M$  is countably  $n$ -rectifiable, we can write (see Remark 1.3 of Ch.3)

$$M = \cup_{j=0}^{\infty} M_j,$$

where  $M_i \cap M_j = \emptyset \forall i \neq j$ ,  $\mathcal{H}^n(M_0) = 0$ , and  $M_j \subset N_j$   $j \geq 1$ , with  $N_j$  an embedded  $C^1$  submanifold of  $\mathbb{R}^{n+k}$ . By virtue of this decomposition and the definition of  $\nabla^M$  (in §2 of Ch.3) it easily follows that if  $h$  is Lipschitz on  $\mathbb{R}^{n+k}$  and if  $h^{(\sigma)}$  are the mollified functions (as in §2 of Ch.2) then, as  $\sigma \downarrow 0$ ,

$$(1) \quad \int_W v \cdot \nabla^M h^{(\sigma)} d\mu_T \rightarrow \int_W v \cdot \nabla^M h d\mu_T$$

for each  $W \subset\subset U$  and each fixed bounded  $\mathcal{H}^n$ -measurable  $v : U \rightarrow \mathbb{R}^{n+k}$ . (Indeed to check this, we have merely to check that (1) holds with  $N_j$  in place of  $M_j$  and with  $v$  vanishing on  $\mathbb{R}^{n+k} \setminus M_j$ ; since  $N_j$  is  $C^1$  this follows fairly easily by approximating  $v$  by smooth functions and using the fact that  $h^{(\sigma)}$  converges to  $h$  uniformly.)

Next let  $\varepsilon > 0$  and let  $\gamma$  be the Lipschitz function on  $\mathbb{R}$  defined by

$$\gamma(s) = \begin{cases} 1, & s < t - \varepsilon \\ \text{linear}, & t - \varepsilon \leq s \leq t \\ 0, & s > t \end{cases}$$

and apply the above to  $h = \gamma \circ f$ . Then letting  $\omega \in \mathcal{D}^n(U)$  we have

$$\begin{aligned} \partial T(h^{(\sigma)}\omega) &= T(d(h^{(\sigma)}\omega)) \\ &= T(dh^{(\sigma)} \wedge \omega) + T(h^{(\sigma)}d\omega). \end{aligned}$$

Then using the integral representations of the form 2.9 for  $\partial T$  we see that

$$(2) \quad (\partial T \llcorner h)(\omega) = \lim_{\sigma \downarrow 0} T(dh^{(\sigma)} \wedge \omega) + (T \llcorner h)(d\omega).$$

Since  $\xi(x)$  orients  $T_x M$ , we have

$$\begin{aligned} \langle \xi(x), dh^{(\sigma)} \wedge \omega \rangle &= \langle \xi(x), (dh^{(\sigma)}(x))^T \wedge \omega^T \rangle \\ &= \langle \xi(x), (dh^{(\sigma)}(x))^T \wedge \omega \rangle \end{aligned}$$

(where  $(\ )^T$  denotes the orthogonal projection of  $\Lambda^q(\mathbb{R}^{n+k})$  onto  $\Lambda^q(T_x M)$ ). Thus

$$\begin{aligned} T(dh^{(\sigma)} \wedge \omega) &= \int_M \langle \xi(x), (dh^{(\sigma)}(x))^T \wedge \omega \rangle \theta d\mathcal{H}^n \\ &= \int_M \langle \xi(x) \lrcorner \nabla^M h^{(\sigma)}(x), \omega \rangle \theta d\mathcal{H}^n \end{aligned}$$

so that by (1)

$$(3) \quad \lim_{\sigma \downarrow 0} T(dh^{(\sigma)} \wedge \omega) = \int_M \langle \xi(x) \lrcorner \nabla^M h(x), \omega \rangle \theta d\mathcal{H}^n.$$

By 2.1 of Ch. 3 and by the chain rule for the composition of Lipschitz functions we have

$$(4) \quad \nabla^M h = \gamma'(f) \nabla^M f \quad \mathcal{H}^n\text{-a.e. on } M$$

(where we set  $\gamma'(f) = 0$  when  $f$  takes the “bad” values  $t$  or  $t - \varepsilon$ ; note that  $\nabla^M h(x) = \nabla^M f(x) = 0$  for  $\mathcal{H}^n$ -a.e. in  $\{x \in M : f(x) = c\}$ ,  $c$  any given constant).

Using (3), (4) in (2), we thus deduce

$$(\partial T \lrcorner h)(\omega) = -\varepsilon^{-1} \int_M \langle \xi \lrcorner \nabla^M f, \omega \rangle \theta d\mathcal{H}^n + (T \lrcorner h)(d\omega).$$

Finally we let  $\varepsilon \downarrow 0$  and we use 4.2 with  $g = \theta \langle \xi \lrcorner \frac{\nabla^M f}{|\nabla^M f|}, \omega \rangle$  in order to complete the proof of (2); by considering a countable dense set of  $\omega \in \mathcal{D}^n(U)$  one can of course show that 4.2 is applicable with  $g = \theta \langle \xi \lrcorner \frac{\nabla^M f}{|\nabla^M f|}, \omega \rangle$  except for a set  $F$  of  $t$  having  $\mathcal{L}^1$ -measure zero, with  $F$  independent of  $\omega$ .

Finally to prove part (3) of the theorem, we first apply part (2) with  $\partial T$  in place of  $T$ . Since  $\partial^2 T = 0$ , this gives

$$\langle \partial T, f, t \rangle = \partial [(\partial T) \lrcorner \{f < t\}].$$

On the other hand, applying  $\partial$  to each side of the original identity (for  $T$ ) of (2), we get

$$\partial [(\partial T) \lrcorner \{f < t\}] = -\partial \langle T, f, t \rangle$$

and hence (3) is established.  $\square$

Motivated by the above discussions we are led to define slices for an arbitrary current  $\in \mathcal{D}_n(U)$  which, together with its boundary, has locally finite mass in  $U$ . Specifically, suppose  $\mathbb{M}_W(T) + \mathbb{M}_W(\partial T) < \infty \forall W \subset\subset U$ . Then we define “slices”

$$4.6 \quad \langle T, f, t_- \rangle = \partial(T \llcorner \{f < t\}) - (\partial T) \llcorner \{f < t\}$$

and

$$4.7 \quad \langle T, f, t_+ \rangle = -\partial(T \llcorner \{f > t\}) + (\partial T) \llcorner \{f > t\}.$$

Of course  $\langle T, f, t_+ \rangle = \langle T, f, t_- \rangle$  (and the common value is denoted  $\langle T, f, t \rangle$ ) for all but the countably many values of  $t$  such that  $\mathbb{M}(T \llcorner \{f = t\}) + \mathbb{M}((\partial T) \llcorner \{f = t\}) > 0$ .

The important properties of the above slices are that if  $f$  is Lipschitz on  $U$  (and if we continue to assume  $\mathbb{M}_W(T) + \mathbb{M}_W(\partial T) < \infty \forall W \subset\subset U$ ), then

$$4.8 \quad \text{spt } \langle T, f, t_+ \rangle \subset \text{spt } T \cap \{x : f(x) = t\}$$

and,  $\forall$  open  $W \subset U$ ,

$$4.9 \quad \begin{cases} \mathbb{M}_W(\langle T, f, t_+ \rangle) \leq \text{ess sup}_W |Df| \liminf_{h \downarrow 0} h^{-1} \mathbb{M}_W(T \llcorner \{t < f < t+h\}) \\ \mathbb{M}_W(\langle T, f, t_- \rangle) \leq \text{ess sup}_W |Df| \liminf_{h \downarrow 0} h^{-1} \mathbb{M}_W(T \llcorner \{t-h < f < t\}). \end{cases}$$

Notice that  $\mathbb{M}_W(T \llcorner \{f < t\})$  is increasing in  $t$ , hence is differentiable for  $\mathcal{L}^1$ -a.e.  $t \in \mathbb{R}$  and  $\int_a^b \frac{d}{dt} \mathbb{M}_W(T \llcorner \{f < t\}) dt \leq \mathbb{M}_W(T \llcorner \{a < f < b\})$ . Thus 4.9 gives

$$4.10 \quad \int_a^{*b} \mathbb{M}_W(\langle T, f, t_{\pm} \rangle) dt \leq \text{ess sup}_W |Df| \mathbb{M}(T \llcorner \{a < f < b\})$$

for every open  $W \subset U$ .

To prove 4.8 and 4.9 we consider first the case when  $f$  is  $C^1$  and take any smooth increasing function  $\gamma : \mathbb{R} \rightarrow \mathbb{R}_+$  and note that

$$4.11 \quad \begin{aligned} \partial(T \llcorner \gamma \circ f)(\omega) - ((\partial T) \llcorner \gamma \circ f)(\omega) &= (T \llcorner \gamma \circ f)(d\omega) - ((\partial T) \llcorner \gamma \circ f)(\omega) \\ &= T(\gamma \circ f d\omega) - T(d(\gamma \circ f)\omega) \\ &= -T(\gamma'(f) df \wedge \omega). \end{aligned}$$

Now let  $\varepsilon > 0$  be arbitrary and choose  $\gamma$  such that

$$\gamma(t) = 0 \text{ for } t < a, \quad \gamma(t) = 1 \text{ for } t > b, \quad 0 \leq \gamma'(t) \leq \frac{1+\varepsilon}{b-a} \text{ for } a < t < b.$$

Then the left side of 4.11 converges to  $\langle T, f, a_+ \rangle$  if we let  $b \downarrow a$ , and hence 4.8 follows because  $\text{spt } \gamma' \subset [a, b]$ . Furthermore the right side  $R$  of 4.11 evidently satisfies

$$|R| \leq (\sup_W |Df|) \left( \frac{1 + \varepsilon}{b - a} \right) \mathbb{M}_W(T \llcorner \{a < f < b\}) |\omega| \quad (\text{spt } \omega \subset W)$$

and so we also conclude the first part of 4.11 for  $f \in C^1$ . We similarly establish the second part for  $f \in C^1$ . To handle general Lipschitz  $f$  we simply use  $f^{(\sigma)}$  in place of  $f$  in 4.6, 4.7 and in the above proof, then let  $\sigma \downarrow 0$  where appropriate.

## 5 The Deformation Theorem

The deformation theorem, given below in 5.1 and 5.3, is a central result in the theory of currents, and was first proved by Federer and Fleming [FF60].

The special notation for this section is as follows:

$$n, k \in \{1, 2, \dots\},$$

$$C = [0, 1] \times \dots \times [0, 1] \quad (\text{standard unit cube in } \mathbb{R}^{n+k})$$

$$\mathbb{Z}^{n+k} = \{z = (z^1, \dots, z^{n+k}) : z^j \in \mathbb{Z}\}$$

$$F_\alpha = C \cap \text{span}\{e_{j_1}, \dots, e_{j_n}\} \text{ for } \alpha = (j_1, \dots, j_n) \in I_{n,n+k}$$

$$\mathcal{L}_j = \text{set of all } j\text{-faces} = \{z + F_\alpha : z \in \mathbb{Z}^{n+k}, \alpha \in I_{n,n+k}\}$$

$$\mathcal{L}_j = j\text{-skeleton of the decomposition} = \cup_{F \in \mathcal{L}_j} F$$

$$\mathcal{L}_j(\rho) = \{\rho F : F \in \mathcal{L}_j\}, \rho > 0$$

$$S_\alpha = \text{span}\{e_{i_1}, \dots, e_{i_{n+1}}\} \text{ for } \alpha = (i_1, \dots, i_{n+1}) \in I_{n+1,n+k}.$$

$$p_\alpha \text{ denotes the orthogonal projection of } \mathbb{R}^{n+k} \text{ onto } S_\alpha, \alpha \in I_{n+1,n+k}.$$

**5.1 (Deformation Theorem, unscaled version.)** *Suppose  $T$  is an  $n$ -current in  $\mathbb{R}^{n+k}$  (i.e.  $T \in \mathcal{D}_n(\mathbb{R}^{n+k})$ ) with  $\mathbb{M}(T) + \mathbb{M}(\partial T) < \infty$ . Then we can write*

$$T - P = \partial R + S$$

where  $P, R, S$  satisfy

$$P = \sum_{F \in \mathcal{L}_n} \beta_F \llbracket F \rrbracket \quad (\beta_F \in \mathbb{R}),$$

with

$$\mathbb{M}(P) \leq C\mathbb{M}(T), \quad \mathbb{M}(\partial P) \leq C\mathbb{M}(\partial T)$$

$$\mathbb{M}(R) \leq C\mathbb{M}(T), \quad \mathbb{M}(S) \leq C\mathbb{M}(\partial T)$$

$(C = C(n, k))$ , and

$$\begin{aligned} \text{spt } P \cup \text{spt } R &\subset \left\{ x : \text{dist}(x, \text{spt } T) < 2\sqrt{n+k} \right\} \\ \text{spt } \partial P \cup \text{spt } S &\subset \left\{ x : \text{dist}(x, \text{spt } \partial T) < 2\sqrt{n+k} \right\}. \end{aligned}$$

*In case  $T$  is an integer multiplicity current, then  $P, R$  can be chosen to be integer multiplicity currents (and the  $\beta_F$  appearing in the definition of  $P$  are integers). If in addition  $\partial T$  is integer multiplicity<sup>2</sup>, then  $S$  can be chosen to be integer multiplicity.*

**5.2 Remarks:** (1) Note that this is slightly sharper than the corresponding theorem in [FF60], [Fed69], because there is no term involving  $\mathbb{M}(\partial T)$  in the bound for  $\mathbb{M}(P)$ .

(2) It follows automatically from the other conclusions of the theorem that  $\mathbb{M}(\partial S) \leq C\mathbb{M}(\partial T)$ . Also, it follows from the inequalities  $\mathbb{M}(\partial P), \mathbb{M}(S) \leq C\mathbb{M}(\partial T)$  that  $S = 0$  and  $\partial P = 0$  when  $\partial T = 0$ .

The following “scaled version” of 5.1 is obtained from the above by first changing scale  $s \rightarrow \rho^{-1}s$ , then applying 5.1, then changing scale back by  $x \rightarrow \rho x$ .

**5.3 (Deformation Theorem, scaled version.)** *Suppose  $T, \partial T$  are as in 5.1, and  $\rho > 0$ . Then*

$$T - P = \partial R + S$$

where  $P, R, S$  satisfy

$$\begin{aligned} P &= \sum_{F \in \mathcal{L}_j(\rho)} \beta_F \llbracket F \rrbracket & (\beta_F \in \mathbb{R}) \\ \mathbb{M}(P) &\leq C\mathbb{M}(T), \quad \mathbb{M}(\partial P) \leq C\mathbb{M}(\partial T) \\ \mathbb{M}(R) &\leq C\rho\mathbb{M}(T), \quad \mathbb{M}(S) \leq C\rho\mathbb{M}(\partial T), \end{aligned}$$

and

$$\begin{aligned} \text{spt } P \cup \text{spt } R &\subset \left\{ x : \text{dist}(x, \text{spt } T) < 2\sqrt{n+k}\rho \right\} \\ \text{spt } \partial P \cup \text{spt } S &\subset \left\{ x : \text{dist}(x, \text{spt } \partial T) < 2\sqrt{n+k}\rho \right\}. \end{aligned}$$

*As in 5.1, in case  $T$  is integer multiplicity, so are  $P, R$ ; if  $\partial T$  is integer multiplicity then so is  $S$ .*

The main step in the proof of the deformation theorem will involve “pushing”  $T$  onto the  $n$ -skeleton  $L_n$  via a certain retraction map  $\psi$ . We first have to establish the existence of a suitable class of retraction maps. This is done in the following lemma, in which we

<sup>2</sup>Actually if  $\mathbb{M}(\partial T) < \infty$  then  $\partial T$  is automatically integer multiplicity if  $T$  is—see 6.3 below.

use the notation

$$\begin{aligned} q &= \text{center point of } C = (\tfrac{1}{2}, \tfrac{1}{2}, \dots, \tfrac{1}{2}), \\ L_{k-1}(a) &= a + L_{k-1} \text{ (} a \text{ a given point in } B_{1/4}(q)\text{),} \\ L_{k-1}(a; \rho) &= \left\{ x \in \mathbb{R}^{n+k} : \text{dist}(x, L_{k-1}(a)) < \rho \right\} \quad (\rho \in (0, \tfrac{1}{4})). \end{aligned}$$

Note that  $\text{dist}(L_{k-1}(a), L_n) \geq \frac{1}{4}$  for any point  $a \in B_{1/4}(q)$ .

**5.4 Lemma.** *For every  $a \in B_{1/4}(q)$  there is a locally Lipschitz map*

$$\psi : \mathbb{R}^{n+k} \setminus L_{k-1}(a) \rightarrow \mathbb{R}^{n+k} \setminus L_{k-1}(a)$$

*such that*

$$\begin{aligned} \psi(C \setminus L_{k-1}(a)) &= C \cap L_n, \quad \psi|_{C \cap L_n} = \text{id}_{C \cap L_n}, \\ |D\psi(x)| &\leq \frac{c}{\rho}, \quad \mathcal{L}^{n+k}\text{-a.e. } x \in C \setminus L_{k-1}(a; \rho), \quad \rho \in (0, \tfrac{1}{4}), \end{aligned}$$

*( $c = c(n, k)$ ), and such that*

$$\psi(z + x) = z + \psi(x), \quad x \in \mathbb{R}^{n+k} \setminus L_{k-1}(a), \quad z \in \mathbb{Z}^{n+k}.$$

**Proof:** We first construct a locally Lipschitz retraction  $\psi_0 : C \setminus L_{k-1}(a)$  onto the  $n$ -faces of  $C$ . This is done as follows:

Firstly for each  $j$ -face  $F$  of  $C$ ,  $j \geq n+1$ , let  $a_F \in F$  denote the orthogonal projection of  $a$  onto  $F$ , and let  $\psi_F$  denote the retraction of  $\overline{F} \setminus \{a_F\}$  onto  $\partial F$  which takes a point  $x \in \overline{F} \setminus \{a_F\}$  to the point  $y \in \partial F$  such that  $x \in \{a_F + \lambda(y - a_F) : \lambda \in (0, 1]\}$ . (Thus  $\psi_F$  is the “radial retraction” of  $F$  with  $a_F$  as origin.) Of course  $\psi_F|_{\partial F} = \text{id}_{\partial F}$ . Notice also that for any  $j$ -face  $F$  of  $C$ ,  $j \geq n+1$ , the line segment  $\overline{aa_F}$  is contained in  $L_{k-1}(a)$ ; in fact if  $J_F$  denotes the set of  $\ell$  such that  $S_\ell$  (see notation prior to 5.1) is parallel to an  $(n+1)$ -face of  $F$ , then (because  $\overline{aa_F}$  is orthogonal to  $F$ , hence orthogonal to each  $S_\ell$ ,  $\ell \in J_F$ ) we have

$$(1) \quad \overline{aa_F} \subset \cap_{\ell \in J_F} p_\ell^{-1}(p_\ell(a)),$$

and this is contained in  $L_{k-1}(a)$ , because (by definition)

$$(2) \quad L_{k-1}(a) = \cup_{\ell=1}^N \cup_{z \in \mathbb{Z}^{n+k}} (z + p_\ell^{-1}(p_\ell(a))).$$

Next, for each  $j \geq n+1$ , define

$$\psi^{(j)} : \cup \{ \overline{F} \setminus \{a_F\} : F \text{ is a } j\text{-face of } C \} \rightarrow \cup \{ \overline{G} : G \text{ is a } (j-1)\text{-face of } C \}$$

by setting

$$\psi^{(j)}|_{\overline{F} \setminus \{a_F\}} = \psi_F.$$

(Notice that then  $\psi^{(j)}$  is locally Lipschitz on its domain by virtue of the fact that each  $\psi_F$  is the identity on  $\partial F$ ,  $F$  any  $j$ -face of  $C$ .)

Then the composite  $\psi^{(n+1)} \circ \psi^{(n+2)} \circ \dots \circ \psi^{(n+k)}$  makes sense on  $C \setminus L_{k-1}(a)$  (by (1)), so we can set

$$\psi_0 = \psi^{(n+1)} \circ \psi^{(n+2)} \circ \dots \circ \psi^{(n+k)}|_{C \setminus L_{k-1}(a)}.$$

Notice that  $\psi_0$  has the additional property that if

$$z \in \mathbb{Z}^{n+k} \text{ and } x, z+x \in C, \text{ then } \psi_0(z+x) = z + \psi_0(x).$$

(Indeed  $x, z+x \in C$  means that either  $x, z+x$  are in  $L_n$  (where  $\psi_0$  is the identity) or else lie in the interior of parallel  $j$ -faces  $F_1, F_2 = z + F_1$  ( $j \geq n+1$ ) of  $C$  with  $z$  orthogonal to  $F_1$  and  $a_{F_2} = z + a_{F_1}$ .) It follows that we can then define a retraction  $\psi$  of all of  $C \setminus L_{k-1}(a)$  onto  $L_n$  by setting

$$\psi(z+x) = z + \psi_0(x), \quad x \in C \setminus L_{k-1}(a), \quad z \in \mathbb{Z}^{n+k}.$$

We now claim that

$$(3) \quad \sup |D\psi| \leq \frac{c}{\rho} \text{ on } \mathbb{R}^{n+k} \setminus L_{k-1}(a, \rho), \quad c = c(n, k).$$

(This will evidently complete the proof of the lemma.)

We can prove (3) by induction on  $k$  as follows. First note that (3) is evident from construction in case  $k = 1$ . Hence assume  $k \geq 2$  and assume (3) holds in case  $k-1$  replaces  $k$  in the above construction. Let  $x$  be any point of interior  $(C) \setminus L_{k-1}(a; \rho)$ , let  $y = \psi^{n+k}(x)$  ( $\psi^{n+k}$  is the radial retraction of  $C \setminus \{a\}$  onto  $\partial C$ , and let  $F$  be any closed  $(n+k-1)$ -face of  $C$  which contains  $y$ ).

Suppose now new coordinates are selected so that  $F \subset \mathbb{R}^{n+k-1} \times \{0\} \subset \mathbb{R}^{n+k}$ , and also let  $\tilde{L}_{k-2}(a) = L_{k-1}(a) \cap \mathbb{R}^{n+k-1} \times \{0\}$ . By virtue of (1) we have  $a_F \in L_{k-1}(a)$ , hence

$$(4) \quad |y - a_F| \geq \text{dist}(y, L_{k-1}(a)).$$

Let  $p_F$  be the orthogonal projection of  $\mathbb{R}^{n+k}$  onto  $\mathbb{R}^{n+k-1} \times \{0\} (\supset F)$ , so that  $a_F = p_F(a)$ . Evidently  $|p_F(x) - a_F| \geq \text{dist}(x, p_F^{-1}(p_F(a)))$  and hence by (2) we deduce

$$(5) \quad |p_F(x) - a_F| \geq \text{dist}(x, L_{k-1}(a)).$$

Furthermore by definition of  $y$  we know that  $y - a = \frac{|y-a|}{|x-a|}(x-a)$  and hence, applying  $p_F$ , we have

$$y - a_F = \frac{|y-a|}{|x-a|} p_F(x-a).$$

Hence since  $|y - a| \geq 3/4$ , we have

$$(6) \quad |y - a_F| \geq \frac{3}{4} \frac{|p_F(x - a)|}{|x - a|}.$$

Now let  $\tilde{\psi}$  be the retraction of  $F \setminus \tilde{L}_{k-2}(a)$  onto the  $n$ -faces of  $F$  ( $\tilde{\psi}$  defined as for  $\psi$  but with  $(k-1)$  in place of  $k$ ,  $a_F$  in place of  $a$ ,  $\mathbb{R}^{n+k-1}$  in place of  $\mathbb{R}^{n+k}$  and  $\tilde{L}_{k-2}(a) = L_{k-2}(a_F)$  in place of  $L_{k-1}(a)$ ). By the inductive hypothesis, together with (4), (5), (6) we have

$$(7) \quad \begin{aligned} |\bar{D}\tilde{\psi}(y)| &\leq \frac{c}{\text{dist}(y, \tilde{L}_{k-2}(a))}, \quad (|\bar{D}\tilde{\psi}(y)| = \limsup_{z \rightarrow y} \frac{|\tilde{\psi}(z) - \tilde{\psi}(y)|}{|z - y|}) \\ &\leq \left(\frac{4}{3}\right)c \frac{1}{|y - a_F|} \frac{|x - a|}{|p_F(x - a)|} \\ &\leq \left(\frac{4}{3}\right)c \frac{|x - a|}{\text{dist}(x, L_{k-1}(a))}, \end{aligned}$$

when  $k = 2$ . For general  $k$ , we label  $L = L_{k-2}(a_F)$  and note that  $\frac{\text{dist}(y, L)}{\text{dist}(x, p_F^{-1}(L))} = \frac{|y - a|}{|x - a|}$  by similarity, and  $p_F^{-1}(L) \subset L_{k-1}(a)$ . So  $|\bar{D}\tilde{\psi}(y)| \leq \frac{c|x - a|}{\text{dist}(x, L_{k-1}(a))}$  as required. Also, by the definition of  $\psi^{n+k}$  we have that

$$(8) \quad |\bar{D}\psi^{n+k}(x)| \leq \frac{c}{|x - a|}, \quad |\bar{D}\psi^{n+k}(x)| = \limsup_{y \rightarrow x} \frac{|\psi^{n+k}(y) - \psi^{n+k}(x)|}{|y - x|}.$$

Since  $\psi(x) = \tilde{\psi} \circ \psi^{n+k}(x)$ , we have by (7), (8) and the chain rule that

$$\begin{aligned} |\bar{D}\psi(x)| &\leq |\bar{D}\tilde{\psi}(y)| |\bar{D}\psi^{n+k}(x)| \leq \frac{c}{|x - a|} \frac{|x - a|}{\text{dist}(x, L_{k-1}(a))} \\ &= \frac{c}{\text{dist}(x, L_{k-1}(a))}. \quad \square \end{aligned}$$

## Proof of the Deformation Theorem:

We use the subspaces  $S_1, \dots, S_N$  and projections  $p_1, \dots, p_N$  introduced at the beginning of the section. Let  $F_j = C \cap S_j$  (so that  $F_j$  is a closed  $(n+1)$ -dimensional face of  $C$ ), let  $x_j$  be the central point of  $F_j$ , and for each  $j = 1, \dots, N$  define a “good” subset  $G_j \subset F_j \cap B_{\frac{1}{4}}(x_j)$  by  $g \in G_j \iff g \in F_j \cap B_{\frac{1}{4}}(x_j)$  and

$$(1) \quad \mathbb{M}(T \sqcup \cup_{z \in \mathbb{Z}^{n+k} \cap S_j} p_j^{-1}(B_\rho(g + z))) \leq \beta \rho^{n+1} \mathbb{M}(T) \quad \forall \rho \in (0, \frac{1}{4})$$

( $\beta$  to be chosen,  $G_j = G_j(\beta)$ ).

We now claim that the “bad” set  $B_j = F_j \cap B_{\frac{1}{4}}(x_j) \setminus G_j$  in fact has  $\mathcal{L}^{n+1}$ -measure (taken in  $S_j$ ) small; in fact we claim

$$(2) \quad \mathcal{L}^{n+1}(B_j) \leq 20^{n+1} \beta^{-1} \omega_{n+1} \left(\frac{1}{4}\right)^{n+1} \quad (\omega_{n+1} = \mathcal{L}^{n+1}(B_1(0))),$$

which is indeed small if we choose large  $\beta$ . To see (2), we argue as follows. For each  $b \in B_j$  there is (by definition) a  $\rho_b \in (0, \frac{1}{4})$  such that

$$(3) \quad \mathbb{M}(T \llcorner \cup_{z \in \mathbb{Z}^{n+k} \cap S_j} p_j^{-1}(B_{\rho_b}(b+z))) > \beta \rho_b^{n+1} \mathbb{M}(T),$$

and by the 5-times Covering Lemma 3.4 of Ch. 1 there is a pairwise disjoint subcollection  $\{B_{\rho_\ell}(b_\ell)\}_{\ell=1,2,\dots}$  ( $\rho_\ell = \rho_{b_\ell}$ ) of the collection  $\{B_{\rho_b}(b)\}_{b \in B_j}$  such that

$$(4) \quad B_j \subset \cup_\ell B_{5\rho_\ell}(b_\ell).$$

But then, setting  $b = b_\ell$  in (3) and summing, we get

$$\beta(\sum_\ell \rho_\ell^{n+1}) \mathbb{M}(T) \leq \mathbb{M}(T) \quad (\text{i.e. } \sum_\ell \rho_\ell^{n+1} \leq \beta^{-1}),$$

(using the fact that  $\{p_j^{-1}B_{\rho_\ell}(b_\ell+z)\}_{\ell=1,2,\dots,z \in \mathbb{Z}^{n+k} \cap S_j}$  is a pairwise disjoint collection for fixed  $j$ ). Thus by (4) we conclude

$$\mathcal{L}^{n+1}(B_j) \leq \beta^{-1} 5^{n+1} \omega_{n+1},$$

which after trivial re-arrangement gives (2) as required. Thus we have

$$\mathcal{L}^{n+1}(G_j) \geq (1 - 20^{n+1} \beta^{-1}) \omega_{n+1} \left(\frac{1}{4}\right)^{n+1},$$

and it follows that

$$(5) \quad \mathcal{L}^{n+k}(p_j^{-1}(G_j) \cap B_{\frac{1}{4}}(q)) \geq (1 - \frac{\omega_{n+1}}{\omega_{n+k}} 20^{n+1} \beta^{-1}) \omega_{n+k} \left(\frac{1}{4}\right)^{n+k},$$

where  $q$  is the center point  $(\frac{1}{2}, \dots, \frac{1}{2})$  of  $C$ . (So  $p_j(q) = x_j$ .)

Then selecting  $\beta$  large enough so that  $20^{n+1} \omega_{n+1} N \beta^{-1} < \omega_{n+k}/(n+k)$ , we see from (5) that we can choose a point  $a \in \cap_{j=1}^N p_j^{-1}(G_j) \cap B_{\frac{1}{4}}(q)$ . Next let  $L_{k-1}(a) = a + L_{k-1}$ ,  $L_{k-1}(a; \rho) = \{x \in \mathbb{R}^{n+k} : \text{dist}(x, L_{k-1}(a)) < \rho\}$  (as in the proof of 5.4) and note that in fact

$$L_{k-1}(a; \rho) = \cup_{j=1}^N \cup_{z \in \mathbb{Z}^{n+k} \cap S_j} p_j^{-1}(B_\rho(p_j(a) + z)).$$

Then since  $p_j(a) \in G_j$  we have (by definition of  $G_j$ )

$$(6) \quad \mathbb{M}(T \llcorner L_{k-1}(a; \rho)) \leq N \beta \rho^{n+1} \mathbb{M}(T) \quad \forall \rho \in (0, \frac{1}{4}).$$

Indeed let us suppose that we take  $\beta$  such that  $20^{n+1}\omega_{n+1}N\beta^{-1} < \omega_{n+k}/2(n+k)$ . Then more than half the ball  $B_{\frac{1}{4}}(q)$  is in the set  $\cap_{j=1}^N p_j^{-1}(G_j)$  and hence, repeating the whole argument above with  $\partial T$  in place of  $T$ , we can actually select  $a$  so that, *in addition to* (6), we also have

$$(7) \quad \mathbb{M}(\partial T \llcorner L_{k-1}(a; \rho)) \leq N\beta\rho^{n+1}\mathbb{M}(\partial T) \quad \forall \rho \in (0, \tfrac{1}{4}).$$

Now let  $\psi$  be the retraction of  $\mathbb{R}^{n+k} \setminus L_{k-1}(a)$  onto  $L_n$  given in 5.4, and let

$$(8) \quad T_\rho = T \llcorner L_{k-1}(a; \rho), \quad (\partial T)_\rho = \partial T \llcorner L_{k-1}(a; \rho),$$

so that by (6), (7)

$$(9) \quad \mathbb{M}(T_\rho) \leq c\rho^{n+1}\mathbb{M}(T), \quad \mathbb{M}((\partial T)_\rho) \leq c\rho^{n+1}\mathbb{M}(\partial T), \quad \forall \rho \in (0, \tfrac{1}{4}).$$

Furthermore by 4.10 we know that for each  $\rho \in (0, \frac{1}{4})$  we can find  $\rho^* \in (\rho/2, \rho)$  such that

$$(10) \quad \mathbb{M}(\langle T, d, \rho^* \rangle) \leq \frac{c}{\rho} \mathbb{M}(T_\rho - T_{\rho/2}) \leq c\rho^n \mathbb{M}(T),$$

where  $d$  is the (Lipschitz) distance function to  $L_{k-1}(a)$  ( $d(x) = \text{dist}(x, L_{k-1}(a))$ ,  $\text{Lip}(d) = 1$ ) and  $\langle T, d, \rho^* \rangle$  is the slice of  $T$  by  $d$  at  $\rho^*$ . (Notice that we can choose  $\rho^*$  such that (10) holds and such that  $\langle T, d, \rho^* \rangle$  is integer multiplicity—see 4.5 and the following discussion.)

We now want to apply the homotopy formula 2.32 to the case when  $h(x, t) = x + t(\psi(x) - x)$ ,  $\in \mathbb{R}^{n+k} \setminus L_{k-1}(a; \sigma)$ ,  $\sigma > 0$ . Notice that  $h$  is Lipschitz on  $\mathbb{R}^{n+k} \setminus L_{k-1}(a; \sigma)$ , so we can define  $h_\#$ ,  $\psi_\#$  as in 2.36. (We shall apply  $h_\#$ ,  $\psi_\#$  only to currents supported away from  $[0, 1] \times L_{k-1}(a)$  and  $L_{k-1}(a)$  respectively.)

Keeping this in mind we note that by 5.4, (6) and (7) we have

$$(11) \quad \mathbb{M}(\psi_\#(T_\rho - T_{\rho/2})) \leq \frac{c}{\rho^n} \rho^{n+1} \mathbb{M}(T) \leq c\rho \mathbb{M}(T)$$

and

$$(12) \quad \mathbb{M}(\psi_\#((\partial T)_\rho - (\partial T)_{\rho/2})) \leq \frac{c}{\rho^{n-1}} \rho^{n+1} \mathbb{M}(\partial T) \leq c\rho \mathbb{M}(\partial T).$$

Similarly by the homotopy formula 2.32, together with 2.34 and (6), (7) above, we have

$$(13) \quad \mathbb{M}(h_\#(\llbracket(0, 1)\rrbracket \times (T_\rho - T_{\rho/2}))) \leq c\rho \mathbb{M}(T)$$

and

$$(14) \quad \mathbb{M}(h_\#(\llbracket(0, 1)\rrbracket \times ((\partial T)_\rho - (\partial T)_{\rho/2}))) \leq c\rho \mathbb{M}(\partial T).$$

Notice also that by (6), (10) and 2.34 we have

$$(15) \quad \mathbb{M}(\psi_{\#} \langle T, d, \rho^* \rangle) \leq c\rho \mathbb{M}(T)$$

and

$$(16) \quad \mathbb{M}(h_{\#}(\llbracket(0, 1)\rrbracket \times \langle T, d, \rho^* \rangle)) \leq c\rho \mathbb{M}(T).$$

Next note that by iteration (11), (12) imply

$$(17) \quad \begin{cases} \mathbb{M}(\psi_{\#}(T_{\rho} - T_{\rho/2^{\nu}})) \leq 2c\rho \mathbb{M}(T) \\ \mathbb{M}(\psi_{\#}((\partial T)_{\rho} - (\partial T)_{\rho/2^{\nu}})) \leq 2c\rho \mathbb{M}(\partial T) \end{cases}$$

for each integer  $\nu \geq 1$ , where  $c$  is as in (11), (12) ( $c$  independent of  $\nu$ ). Selecting  $\rho = \frac{1}{4}$  and using the arbitrariness of  $\nu$ , it follows that

$$(18) \quad \begin{cases} \mathbb{M}(\psi_{\#}(T - T_{\sigma})) \leq c\mathbb{M}(T) \\ \mathbb{M}(\psi_{\#}(\partial T - (\partial T)_{\sigma})) \leq c\mathbb{M}(\partial T) \end{cases}$$

for each  $\sigma \in (0, 1)$  (with  $c$  independent of  $\sigma$ ).

Now select  $\rho = \rho_{\nu} = 2^{-\nu}$  and  $\rho_{\nu}^* \in [2^{-\nu-1}, 2^{-\nu}]$  such that (10), (15), (16) hold with  $\rho_{\nu}^*$  in place of  $\rho^*$ ; then by (15), (16), (17), (18) we have that

$$\begin{aligned} & \psi_{\#}(T - T_{\rho_{\nu}^*}), \quad h_{\#}(\llbracket(0, 1)\rrbracket \times (T - T_{\rho_{\nu}^*})), \\ & \psi_{\#}(\partial T - \partial T_{\rho_{\nu}^*}), \quad h_{\#}(\llbracket(0, 1)\rrbracket \times \partial(T - T_{\rho_{\nu}^*})) \end{aligned}$$

are Cauchy sequences relative to  $\mathbb{M}$ , and  $\mathbb{M}(\langle T, d, \rho_{\nu}^* \rangle) + \mathbb{M}(\psi_{\#} \langle T, d, \rho_{\nu}^* \rangle) \rightarrow 0$ . Hence there are currents  $Q, S_1 \in \mathcal{D}_n(\mathbb{R}^{n+k})$  and  $R_1 \in \mathcal{D}_{n+1}(\mathbb{R}^{n+k})$  such that

$$(19) \quad \begin{cases} \lim \mathbb{M}(Q - \psi_{\#}(T - T_{\rho_{\nu}^*})) = 0 \\ \lim \mathbb{M}(S_1 - h_{\#}(\llbracket(0, 1)\rrbracket \times \partial(T - T_{\rho_{\nu}^*}))) = 0 \\ \lim \mathbb{M}(R_1 - h_{\#}(\llbracket(0, 1)\rrbracket \times (T - T_{\rho_{\nu}^*}))) = 0. \end{cases}$$

Furthermore by the homotopy formula and 2.34 we have for each  $\nu$

$$(20) \quad \begin{aligned} T - T_{\rho_{\nu}^*} - \psi_{\#}(T - T_{\rho_{\nu}^*}) &= \partial(h_{\#}(\llbracket(0, 1)\rrbracket \times (T - T_{\rho_{\nu}^*}))) \\ &\quad - h_{\#}(\llbracket(0, 1)\rrbracket \times \partial(T - T_{\rho_{\nu}^*})). \end{aligned}$$

Since  $\partial T_{\rho_{\nu}^*} = (\partial T)_{\rho_{\nu}^*} - \langle T, d, \rho_{\nu}^* \rangle$  (by the definition 4.6, 4.7 of slice) we thus get that

$$(21) \quad T - Q = \partial R_1 + S_1.$$

(Notice that  $Q$ ,  $R_1$  are integer multiplicity by (19), 4.4, 4.5 and 3.17 in case  $T$  is integer multiplicity; similarly  $S_1$  is integer multiplicity if  $\partial T$  is.)

Using the fact that  $\psi$  retracts  $\mathbb{R}^{n+k} \setminus L_{k-1}(a)$  onto  $L_n$  we know (by 2.34) that  $\text{spt } \psi_\#(T - T_{\rho_v}^*) \subset L_n$ , and hence

$$(22) \quad \text{spt } Q \subset L_n.$$

We also have (since  $\psi(z + C) \subset z + C \ \forall z \in \mathbb{Z}^{n+k}$ ) that

$$(23) \quad \begin{cases} \text{spt } R_1 \cup \text{spt } Q \subset \{x : \text{dist}(x, \text{spt } T) < \sqrt{n+k}\} \\ \text{spt } S_1 \subset \{x : \text{dist}(x, \text{spt } \partial T) < \sqrt{n+k}\} \end{cases}$$

and, by (18), (19), we have

$$(24) \quad \mathbb{M}(Q) \leq c\mathbb{M}(T), \ \mathbb{M}(R_1) \leq c\mathbb{M}(T), \ \mathbb{M}(S_1) \leq c\mathbb{M}(\partial T).$$

Also by (18) and the semi-continuity of  $\mathbb{M}$  under weak convergence, we have

$$(25) \quad \begin{aligned} \mathbb{M}(\partial Q) &\leq \liminf \mathbb{M}(\partial \psi_\#(T - T_{\rho_v}^*)) \\ &= \liminf \mathbb{M}(\psi_\# \partial(T - T_{\rho_v}^*)) \\ &\leq c\mathbb{M}(\partial T). \end{aligned}$$

Now let  $F$  be a given face of  $L_n$  (i.e.  $F \in \mathcal{L}_n$ ) and let  $\check{F}$  = interior of  $F$ . Assume for the moment that  $F \subset \mathbb{R}^n \times \{0\} (\subset \mathbb{R}^{n+k})$ , and let  $p$  be the orthogonal projection onto  $\mathbb{R}^n \times \{0\}$ . By construction of  $\psi$  we know that  $p \circ \psi = \psi$  in a neighborhood of any point  $y \in \check{F}$ . We therefore have (since  $Q$  is given by (18)) that

$$(26) \quad p_\#(Q \llcorner \check{F}) = Q \llcorner \check{F}.$$

It then follows, by the obvious modifications of the arguments in the proof of the Constancy Theorem 2.41 and in 2.42, that

$$(27) \quad Q \llcorner \check{F} = \int_{\check{F}} \langle e_1 \wedge \cdots \wedge e_n, \omega(x) \rangle \theta_F(x) d\mathcal{L}^n(x)$$

$\forall \omega \in \mathcal{D}^n(\mathbb{R}^{n+k})$ , for some  $BV_{\text{loc}}(\mathbb{R}^n)$  function  $\theta_F$ , and

$$(28) \quad \mathbb{M}(Q \llcorner \check{F}) = \int_{\check{F}} |\theta_F| d\mathcal{L}^n, \ \mathbb{M}((\partial Q) \llcorner \check{F}) = \int_{\check{F}} |D\theta_F|.$$

Furthermore, since

$$(29) \quad (Q \llcorner \check{F} - \beta \llbracket F \rrbracket)(\omega) = \int_{\check{F}} (\theta_F - \beta) \langle e_1 \wedge \cdots \wedge e_n, \omega(x) \rangle d\mathcal{L}^n(x)$$

(by (27)), we have (again using the reasoning of 2.42)

$$(30) \quad \begin{cases} \mathbb{M}(Q \llcorner \check{F} - \beta \llbracket F \rrbracket) = \int_{\check{F}} |\theta_F - \beta| d\mathcal{L}^n \\ \mathbb{M}(\partial(Q \llcorner \check{F} - \beta \llbracket F \rrbracket)) = \int_{\mathbb{R}^n} |D(\chi_{\check{F}}(\theta_F - \beta))|, \end{cases}$$

where  $\chi_{\check{F}}$  = characteristic function of  $\check{F}$ . Thus taking  $\beta = \beta_F$  such that

$$(31) \quad \min \left\{ \mathcal{L}^n \left\{ x \in \check{F} : \theta_F \geq \beta \right\}, \mathcal{L}^n \left\{ x \in \check{F} : \theta_F(x) \leq \beta \right\} \right\} \geq \frac{1}{2}$$

(which we can do because  $\mathcal{L}^n(\check{F}) = 1$ ; notice that we can, and we do, take  $\beta_F \in \mathbb{Z}$  if  $\theta_F$  is integer-valued), we have by 2.7 and 2.9 of Ch. 2, (28) and (30) that

$$(32) \quad \begin{cases} \mathbb{M}(Q \llcorner \check{F} - \beta \llbracket F \rrbracket) \leq c \int_{\check{F}} |D\theta_F| = c\mathbb{M}(\partial Q \llcorner \check{F}) \\ \mathbb{M}(\partial(Q \llcorner \check{F} - \beta \llbracket F \rrbracket)) \leq c \int_{\check{F}} |D\theta_F| = c\mathbb{M}(\partial Q \llcorner \check{F}). \end{cases}$$

We also have by 2.45(1)

$$(33) \quad Q \llcorner \partial F = 0.$$

Then summing over  $F \in \mathcal{L}_n$  and using (32), (33) we have, with  $P = \sum_{F \in \mathcal{L}_n} \beta_F \llbracket F \rrbracket$ , that

$$(34) \quad \begin{cases} \mathbb{M}(Q - P) \leq c\mathbb{M}(\partial Q) \\ \mathbb{M}(\partial Q - \partial P) \leq c\mathbb{M}(\partial Q). \end{cases}$$

Actually by (31) we have

$$(35) \quad |\beta_F| \leq 2 \int_{\check{F}} |\theta_F| d\mathcal{L}^n,$$

and hence (using again the first part of (28)), since  $\mathbb{M}(P) = \sum_F |\beta_F|$ ,

$$(36) \quad \mathbb{M}(P) \leq c\mathbb{M}(Q).$$

Notice that the second part of (34) gives

$$(37) \quad \mathbb{M}(\partial P) \leq c\mathbb{M}(\partial Q).$$

Finally we note that (21) can be written

$$(38) \quad T - P = \partial R_1 + (S_1 + (Q - P)).$$

Setting  $R = R_1$ ,  $S = S_1 + (Q - P)$ , the theorem now follows immediately from (23), (24), (25) and (34), (36), (37), (38); the fact that  $P, R$  are integer multiplicity if  $T$  is should be evident from the remarks during the course of the above proof, as should be the fact that  $S$  is integer multiplicity if  $T, \partial T$  are.  $\square$

## 6 Applications of the Deformation Theorem

We here establish a couple of simple (but very important) applications of the deformation theorem, namely the isoperimetric theorem and the weak polyhedral approximation theorem. This latter theorem, when combined with the compactness 3.15, implies the important “Boundary Rectifiability Theorem” (6.3 below), which asserts that if  $T$  is an integer multiplicity current in  $\mathcal{D}_n(U)$  and if  $\mathbb{M}_W(\partial T) < \infty \forall W \subset\subset U$ , then  $\partial T (\in \mathcal{D}_{n-1}(U))$  is integer multiplicity. (Notice that in the case  $k = 0$ , this has already been established in 3.18.)

**6.1 (Isoperimetric Theorem.)** *Suppose  $T \in \mathcal{D}_{n-1}(\mathbb{R}^{n+k})$  is integer multiplicity,  $n \geq 2$ ,  $\text{spt } T$  is compact and  $\partial T = 0$ . Then there is an integer multiplicity current  $R \in \mathcal{D}_n(\mathbb{R}^{n+k})$  with  $\text{spt } R$  compact,  $\partial R = T$ , and*

$$\mathbb{M}(R)^{\frac{n-1}{n}} \leq c \mathbb{M}(T),$$

where  $c = c(n, k)$ .

**Proof:** The case  $T = 0$  is trivial, so assume  $T \neq 0$ . Let  $P, R, S$  be integer multiplicity currents as in 5.3, where for the moment  $\rho > 0$  is arbitrary, and note that  $S = 0$  because  $\partial T = 0$ . Evidently (since  $\mathcal{H}^{n-1}(F) = \rho^{n-1} \forall F \in \mathcal{F}_{n-1}(\rho)$ ) we have

$$(1) \quad \mathbb{M}(P) = N(\rho) \rho^{n-1}$$

for some non-negative integer  $N(\rho)$ . But since  $\mathbb{M} \leq c \mathbb{M}(T)$  (from 5.3) we see that necessarily  $N(\rho) = 0$  in (1) if we choose  $\rho = (2c \mathbb{M}(T))^{\frac{1}{n-1}}$ . Then  $P = 0$ , and 5.3 gives  $T = \partial R$  for some integer multiplicity current  $R$  with  $\text{spt } R$  compact and  $\mathbb{M}(R) \leq c \rho \mathbb{M}(T) = c' (\mathbb{M}(T))^{\frac{1}{n-1}}$ .  $\square$

**6.2 (Weak Polyhedral Approximation Theorem.)** *Given any integer multiplicity  $T \in \mathcal{D}_n(U)$  with  $\mathbb{M}_W(T), \mathbb{M}_W(\partial T) < \infty \forall W \subset\subset U$ , there is a sequence  $\{P_k\}$  of current of the form*

$$(\ddagger) \quad P_k = \sum_{F \in \mathcal{F}_n(\rho_k)} \beta_R^{(k)} \llbracket F \rrbracket, \quad (\beta_R^{(k)} \in \mathbb{Z}), \quad \rho_k \downarrow 0,$$

such that  $P_k \rightharpoonup T$  (and hence also  $\partial P_k \rightharpoonup \partial T$ ) in  $U$  (in the sense of 2.15).

**Proof:** First consider the case  $U = \mathbb{R}^{n+k}$  and  $\mathbb{M}(T), \mathbb{M}(\partial T) < \infty$ . In this case we simply use the deformation theorem: for any sequence  $\rho_k \downarrow 0$ , the scaled version of the deformation theorem (with  $\rho = \rho_k$ ) gives  $P_k$  as in  $(\ddagger)$  such that

$$(1) \quad T - P_k = \partial R_k + S_k$$

for some  $R_k, S_k$  such that

$$(2) \quad \begin{cases} \mathbb{M}(R_k) \leq c\rho_k \mathbb{M}(T) \rightarrow 0 \\ \mathbb{M}(S_k) \leq c\rho_k \mathbb{M}(\partial T) \rightarrow 0 \end{cases}$$

and

$$\mathbb{M}(P_k) \leq c\mathbb{M}(T), \quad \mathbb{M}(\partial P_k) \leq c\mathbb{M}(\partial T).$$

Evidently (1), (2) give  $P_k(\omega) \rightarrow T_k(\omega) \quad \forall \omega \in \mathcal{D}^n(\mathbb{R}^{n+k})$ , and  $\partial P_k = 0$  if  $\partial T = 0$ , so the theorem is proved in case  $U = \mathbb{R}^{n+k}$  and  $T, \partial T$  are of finite mass.

In the general case we take any Lipschitz function  $\varphi$  on  $\mathbb{R}^{n+k}$  such that  $\varphi > 0$  in  $U$ ,  $\varphi \equiv 0$  in  $\mathbb{R}^{n+k} \setminus U$  and such that  $\{x : \varphi(x) > \lambda\} \subset\subset U \quad \forall \lambda > 0$ . For  $\mathcal{L}^1$ -a.e.  $\lambda > 0$ , 4.5 implies that  $T_\lambda = T \llcorner \{x : \varphi(x) > \lambda\}$  is such that  $\mathbb{M}(\partial T_\lambda) < \infty$ . Since  $\text{spt } T_\lambda \subset\subset U$ , we can apply the argument above to approximate  $T_\lambda$  for any such  $\lambda$ . Taking a suitable sequence  $\lambda_j \downarrow 0$ , the required approximation then immediately follows.  $\square$

**6.3 (Boundary Rectifiability Theorem.)** *Suppose  $T$  is an integer multiplicity current in  $\mathcal{D}_n(U)$  with  $\mathbb{M}(\partial T) < \infty \quad \forall W \subset\subset U$ . Then  $\partial T (\in \mathcal{D}_{n-1}(U))$  is an integer multiplicity current.*

**Proof:** A direct consequence of 6.2 above and the Compactness 3.15 (applied with  $\partial T_j$  in place of  $T_j$ ).  $\square$

**6.4 Remark:** The above proof used the Compactness Theorem 3.15 applied to the sequence  $\partial T_j$  rather than to  $T_j$ , so used only the special case of 3.15 when the given sequence  $\{T_j\}$  has  $\partial T_j = 0 \quad \forall j$ .

## 7 The Flat Metric Topology

The main result to be proved here is the equivalence of weak convergence and “flat metric”<sup>3</sup> convergence (see below for terminology) for a sequence of integer multiplicity currents  $\{T_j\} \subset \mathcal{D}_n(U)$  such that  $\sup_{j \geq 1} (\mathbb{M}_W(T_j) + \mathbb{M}_W(\partial T_j)) < \infty \quad \forall W \subset\subset U$ .

We let  $U$  denote (as usual) an arbitrary open subset of  $\mathbb{R}^{n+k}$ ,

$$\mathcal{I} = \{T \in \mathcal{D}_n(U) : T \text{ is integer multiplicity and } \mathbb{M}_W(\partial T) < \infty \quad \forall W \subset\subset U\}.$$

and

$$\mathcal{I}_{M,W} = \{T \in \mathcal{I} : \text{spt } T \subset \overline{W}, \mathbb{M}(T) + \mathbb{M}(\partial T) \leq M\}.$$

<sup>3</sup>Note that the word “flat” here has *no* physical or geometric significance, but relates rather to Whitney’s use of the symbol  $\flat$  (the “flat” symbol in musical notation) in his work. We mention this because it is often a source of confusion.

for any  $M > 0$  and open  $W \subset\subset U$ .

On  $\mathcal{I}$  we define a family of pseudometrics  $\{d_W\}_{W \subset\subset U}$  by

$$7.1 \quad d_W(T_1, T_2) = \inf \{ \mathbb{M}_W(S) + \mathbb{M}_W(R) : T_1 - T_2 = \partial R + S, \text{ where} \\ R \in \mathcal{D}_{n+1}(U), S \in \mathcal{D}_n(U) \text{ are integer multiplicity} \}.$$

We henceforth assume  $\mathcal{I}$  is equipped with the topology given (in the usual way) by the family  $\{d_W\}_{W \subset\subset U}$  of pseudometrics. This topology is called the “flat metric topology” for  $\mathcal{I}$ : there is a countable base of neighborhoods at each point, and  $T_j \rightarrow T$  in this topology if and only if  $d_W(T_j, T) \rightarrow 0 \forall W \subset\subset U$ .

**7.2 Theorem.** *Let  $T, \{T_j\} \subset \mathcal{D}_n(U)$  be integer multiplicity with*

$$\sup_{j \geq 1} (\mathbb{M}_W(T_j) + \mathbb{M}_W(\partial T_j)) < \infty \quad \forall W \subset\subset U.$$

*Then  $T_j \rightarrow T$  (in the sense of 2.15) if and only if  $d_W(T_j, T) \rightarrow 0$  for each  $W \subset\subset U$ .*

**7.3 Remark:** Notice that no use is made of the Compactness 3.15 in this theorem; however if we combine the compactness theorem with it, then we get the statement that for any family of positive (finite) constants  $\{c(W)\}_{W \subset\subset U}$  the set  $\{T \in \mathcal{I} : \mathbb{M}_W(T_j) + \mathbb{M}_W(\partial T_j) \leq c(W) \quad \forall W \subset\subset U\}$  is sequentially compact when equipped with the flat metric topology.

**Proof of 7.2:** First note that the “if” part of the theorem is trivial (indeed for a given  $W \subset\subset U$ , the statement  $d_W(T_j, T) \rightarrow 0$  evidently implies  $(T_j - T)(\omega) \rightarrow 0$  for any fixed  $\omega \in \mathcal{D}^n(U)$  with  $\text{spt } \omega \subset W$ ).

For the “only if” part of the theorem, the main difficulty is to establish the appropriate “total boundedness” property; specifically we show that for any given  $\varepsilon > 0$  and  $W \subset\subset \widetilde{W} \subset\subset U$ , we can find  $N = N(\varepsilon, W, \widetilde{W}, M)$  and integer multiplicity currents  $P_1, \dots, P_N \in \mathcal{D}_n(U)$  such that

$$(1) \quad \mathcal{I}_{M,W} \subset \sum_{j=1}^N B_{\varepsilon, \widetilde{W}}(P_j),$$

where, for any  $P \in \mathcal{I}$ ,  $B_{\varepsilon, \widetilde{W}}(P) = \{S \in \mathcal{I} : d_{\widetilde{W}}(S, P) < \varepsilon\}$ . This is an easy consequence of the Deformation Theorem: in fact for any  $\rho > 0$ , 5.3 guarantees that for  $T \in \mathcal{I}_{M,W}$  we can find integer multiplicity  $P, R, S$  such that

$$(2) \quad \begin{cases} T - P = \partial R + S \\ P = \sum_{F \in \mathcal{F}_n(\rho)} \beta_F \llbracket F \rrbracket, \beta_F \in \mathbb{Z} \\ \text{spt } P \subset \{x : \text{dist}(x, \text{spt } T) < 2\sqrt{n+k}\rho\} \end{cases}$$

$$(3) \quad \begin{cases} \mathbb{M}(P) (\equiv \sum_{F \in \mathcal{F}_n(\rho)} |\beta_F| \rho^n) \leq c \mathbb{M}(T) \leq cM \\ \text{spt } R \cup \text{spt } S \subset \{x : \text{dist}(x, \text{spt } T) < 2\sqrt{n+k}\rho\} \\ \mathbb{M}(R) + \mathbb{M}(S) \leq c\rho \mathbb{M}(T) \leq c\rho M. \end{cases}$$

Then for  $\rho$  small enough to ensure  $2\sqrt{n+k}\rho < \text{dist}(W, \widetilde{W})$ , we see from (2),(3) that

$$d_{\widetilde{W}}(T, P) \leq c\rho M.$$

Hence, since there are only finitely many  $P_1, \dots, P_N$  currents  $P$  as in (2) ( $N$  depends only on  $M, W, n, k, \rho$ ), we have (1) as required.

Next note that (by 4.5(1),(2) and an argument as in 7.7(2) of Ch.2) we can find a subsequence  $\{T_{j'}\} \subset \{T_j\}$  and a sequence  $\{W_i\}$ ,  $W_i \subset\subset W_{i+1} \subset\subset U$ ,  $\cup_{i=1}^{\infty} W_i = U$ , such that  $\sup_{j' \geq 1} \mathbb{M}(\partial(T_{j'} \llcorner W_i)) < \infty \forall i$ . Thus from now on we can assume without loss of generality that  $W \subset\subset U$  and

$$(4) \quad \text{spt } T_j \subset \overline{W} \quad \forall j.$$

Then take any  $\widetilde{W}$  such that  $W \subset\subset \widetilde{W} \subset\subset U$  and apply (1) with  $\varepsilon = 1, \frac{1}{2}, \frac{1}{4}$  etc. to extract a subsequence  $\{T_{j_r}\}_{r=1,2,\dots}$  from  $\{T_j\}$  such that

$$d_{\widetilde{W}}(T_{j_{r+1}}, T_{j_r}) < 2^{-r}$$

and hence

$$(5) \quad T_{j_{r+1}} - T_{j_r} = \partial R_r + S_r$$

where  $R_r, S_r$  are integer multiplicity,

$$\begin{aligned} \text{spt } R_r \cup \text{spt } S_r &\subset \widetilde{W} \\ \mathbb{M}(R_r) + \mathbb{M}(S_r) &\leq \frac{1}{2^r}. \end{aligned}$$

Therefore by 3.17 we can define integer multiplicity  $R^{(\ell)}, S^{(\ell)}$  by the  $\mathbb{M}$ -absolutely convergent series

$$R^{(\ell)} = \sum_{r=\ell}^{\infty} R_r, \quad S^{(\ell)} = \sum_{r=\ell}^{\infty} S_r;$$

then

$$\mathbb{M}(R^{(\ell)}) + \mathbb{M}(S^{(\ell)}) \leq 2^{-\ell+1}$$

and (from (5))

$$T - T_{j_\ell} = \partial R^{(\ell)} + S^{(\ell)}.$$

Thus we have a subsequence  $\{T_{j_\ell}\}$  of  $\{T_j\}$  such that  $d_{\widetilde{W}}(T, T_{j_\ell}) \rightarrow 0$ . Since we can thus extract a subsequence converging relative to  $d_{\widetilde{W}}$  from any given subsequence of  $\{T_j\}$ , we then have  $d_{\widetilde{W}}(T, T_j) \rightarrow 0$ ; since this can be repeated with  $W = W_i$ ,  $\widetilde{W} = W_{i+1} \forall i$  ( $W_i$  as above), the required result evidently follows.  $\square$

## 8 The Rectifiability and Compactness Theorems

Here we prove the important Rectifiability Theorem for currents  $T$  which, together with  $\partial T$ , have locally finite mass and which have the additional property that  $\Theta^{*n}(\mu_T, x) > 0$  for  $\mu_T$ -a.e.  $x$ . The main tool of the proof is the Structure Theorem 3.7 of Ch. 3. Having established the Rectifiability Theorem, we show (in 8.2) that it is then straightforward to establish the Compactness 3.15. Although this proof of the Compactness Theorem has the advantage of being conceptually straightforward, it is rather lengthy if one takes into account the effort needed to prove the Structure Theorem. Recently B. Solomon [Sol82] showed that it is possible to prove the Compactness Theorem (and to develop the whole theory of integer multiplicity currents) without use of the structure theorem.

**8.1 (Rectifiability Theorem.)** *Suppose  $T \in \mathcal{D}_n(U)$  is such that  $\mathbb{M}_W(T) + \mathbb{M}_W(\partial T) < \infty$  for all  $W \subset\subset U$ , and  $\Theta^{*n}(\mu_T, x) > 0$  for  $\mu_T$ -a.e.  $x \in U$ . Then  $T$  is rectifiable; that is*

$$T = \underline{\tau}(M, \theta, \xi),^4$$

where  $M$  is countably  $n$ -rectifiable,  $\mathcal{H}^n$ -measurable,  $\theta$  is a positive locally  $\mathcal{H}^n$ -integrable function on  $M$ , and  $\xi(x)$  orients the approximate tangent space  $T_x M$  of  $M$  for  $\mathcal{H}^n$ -a.e.  $x \in M$  (i.e.  $\xi|_x$  is a measurable function of  $x$  and  $\xi|_x = \pm \tau_1 \wedge \cdots \wedge \tau_n$ , where  $\tau_1, \dots, \tau_n$  is an orthonormal basis for the approximate tangent space  $T_x M$  of  $M$ , for  $\mathcal{H}^n$ -a.e.  $x \in M$ .)

**Proof:** First note that for any locally finite Borel regular measure  $\nu$  on  $\mathbb{R}^{n+k}$ , by The Comparison Theorem 3.3 of Ch. 1, for any  $d > 0$  and any open  $W \subset \mathbb{R}^{n+k}$

$$(1) \quad \mathcal{H}^d \left\{ x \in W : \Theta^{*d}(\nu, x) > t \right\} \leq t^{-1} \nu(W), \quad \forall t > 0.$$

In particular

$$(2) \quad \mathcal{H}^d \left\{ x \in W : \Theta^{*d}(\nu, x) = \infty \right\} = 0$$

for any open  $W \subset\subset \mathbb{R}^{n+k}$ .

Now let

$$M = \{x \in U : \Theta^{*n}(\mu_T, x) > 0\}$$

and let  $W_j$ ,  $j = 1, 2, \dots$ , be open with  $W_j \subset W_{j+1} \subset\subset U \forall j$  and  $\cup_j W_j = U$ . Then  $M = \cup_{j=1}^{\infty} M_j$ , where  $M_j = \{x \in M \cap W_j : \Theta^{*n}(\mu_T, x) > 1/j\}$ , so we see from (1), with  $\nu = \mu_T$  and  $d = n$ , that  $\mathcal{H}^n(M_j) < \infty$  for each  $j$ , so  $M$  is  $\mathcal{H}^n$   $\sigma$ -finite.

<sup>4</sup> The notation here is as for integer multiplicity rectifiable currents as in §3 of the present chapter. That is,  $\underline{\tau}(M, \theta, \xi)(\omega) = \int_M \langle \xi, \omega \rangle \theta d\mathcal{H}^n$ , although  $\theta$  is not assumed to be integer-valued here.

Suppose  $P$  is an  $\mathcal{H}^n$ -measurable purely unrectifiable subset of  $M$ . Then

$$(3) \quad \mu_T(P) = 0$$

by Remark 2.45(2). For any open  $V \supset P \cap M_j$ , by (1), again with  $\nu = \mu_T$  and  $d = n$ ,

$$\mathcal{H}^n(P \cap M_j) \leq j \mu_T(V).$$

Since  $\inf_{V \text{ open}, V \supset P \cap M_j} \mu_T(V) = \mu_T(P)$  by 1.22 of Ch. 1, we thus conclude by (3) that

$$(4) \quad \mathcal{H}^n(P) = 0 \quad \forall \text{ purely unrectifiable } P \subset M.$$

Then by Lemma 3.2 of Ch. 3 we conclude  $M$  is countably  $n$ -rectifiable. Thus we have proved that

$$(5) \quad \mu_T = \mu_T \llcorner M,$$

with  $M \subset U$  is countably  $n$ -rectifiable.

By Theorem 2.43,  $\mu_T$  is absolutely continuous with respect to  $\mathcal{H}^n$  and

$$(6) \quad \mu_T = \mathcal{H}^n \llcorner \theta,$$

where  $\theta$  is a non-negative locally  $\mathcal{H}^n$ -integrable function on  $M$  and  $\theta = 0$  on  $U \setminus M$ . Thus, with  $\xi = \tilde{T}$ , the identity  $T(\omega) = \int_U \langle \omega, \tilde{T} \rangle d\mu_T$  can be written

$$(7) \quad T(\omega) = \int_M \langle \omega, \xi \rangle \theta d\mathcal{H}^n,$$

It thus remains only to prove that  $\xi(x)$  orients  $T_x M$  for  $\mathcal{H}^n$ -a.e.  $x \in M$  (i.e.  $\xi(x) = \pm \tau_1 \wedge \cdots \wedge \tau_n$  for  $\mathcal{H}^n$ -a.e.  $x \in M$ , where  $\tau_1, \dots, \tau_n$  is any orthonormal basis for the approximate tangent space  $T_x M$  of  $M$  with respect to  $\theta$  (such  $T_x M$  exists for  $\mathcal{H}^n$ -a.e.  $x \in M$  as discussed in Theorem 1.9 of Ch. 3).

To check that indeed  $\xi(x)$  orients  $T_x M$ , write  $M = \cup_{j=0}^{\infty} M_j$ ,  $M_j$  pairwise disjoint,  $\mathcal{H}^n(M_0) = 0$ ,  $M_j \subset N_j$ ,  $N_j$  a  $C^1$  embedded submanifold of  $\mathbb{R}^{n+k}$ ,  $j \geq 1$ . By the Upper Density Theorem 3.8 of Ch. 1, for  $j \geq 1$  we have

$$(8) \quad \Theta^{*n}(\mu_T \llcorner ((N_j \cup M) \setminus M_j), x) = 0, \quad \mathcal{H}^n\text{-a.e. } x \in M_j.$$

$\omega = \sum_{\alpha \in I_{n,n+k}} \omega_{\alpha} dx^{\alpha} \in \mathcal{D}^n(\mathbb{R}^{n+k})$ , and, as usual, take  $\eta_{x,\lambda}(y) = \lambda^{-1}(y - x)$ . Then

$\eta_{x,\lambda}^\# \omega = \lambda^{-n} \sum_{\alpha \in I_{n,n+k}} \omega_\alpha \circ \eta_{x,\lambda} dx^\alpha$  has support in  $U$  for small enough  $\lambda$ , and

$$\begin{aligned} \eta_{x,\lambda}^\# T(\omega) &= T(\eta_{x,\lambda}^\# \omega) \\ &= \int_M \langle \eta_{x,\lambda}^\# \omega, \xi \rangle \theta d\mathcal{H}^n \\ &= \int_{N_j} \langle \eta_{x,\lambda}^\# \omega, \xi \rangle \theta d\mathcal{H}^n \\ &\quad + \int_{M \setminus M_j} \langle \eta_{x,\lambda}^\# \omega, \xi \rangle \theta d\mathcal{H}^n - \int_{N_j \setminus M_j} \langle \eta_{x,\lambda}^\# \omega, \xi \rangle \theta d\mathcal{H}^n \\ &= \int_{N_j} \langle \eta_{x,\lambda}^\# \omega, \xi \rangle \theta d\mathcal{H}^n + \varepsilon(\omega, x, \lambda) \end{aligned}$$

where  $\varepsilon(\omega, x, \lambda) \rightarrow 0$  as  $\lambda \downarrow 0$  for  $\mathcal{H}^n$ -a.e.  $x \in M_j$  by (8). That is, after the change of variable  $z = \eta_{x,\lambda}(y)$  (i.e.  $y = x + \lambda z$ ),

$$\eta_{x,\lambda}^\# T(\omega) = \int_{\eta_{x,\lambda}(N_j)} \langle \omega(z), \xi(x + \lambda z) \rangle \theta(x + \lambda z) d\mathcal{H}^n(z) + \varepsilon(\lambda)$$

$\mathcal{H}^n$ -a.e.  $x \in M_j$ . Since  $N_j$  is  $C^1$ , this gives

$$(9) \quad \lim_{\lambda \downarrow 0} \eta_{x,\lambda}^\# T(\omega) = \theta(x) \int_L \langle \omega(z), \xi(x) \rangle d\mathcal{H}^n(z)$$

for  $\mathcal{H}^n$ -a.e.  $x \in M_j$  (independent of  $\omega$ ), where  $L$  is the tangent space  $T_x N_j$  of  $N_j$  at  $x$ . Thus (by Definition 1.7 in Ch.3 of  $T_x M$ ) we have (9) with  $L = T_x M$  for  $\mathcal{H}^n$ -a.e.  $x \in M_j$ . On the other hand, provided  $\text{spt } \omega \subset B_R(0)$ ,

$$\begin{aligned} (10) \quad \partial \eta_{x,\lambda}^\# T(\omega) &= \eta_{x,\lambda}^\# \partial T(\omega) = \partial T(\eta_{x,\lambda}^\# \omega) = \int_{B_{\lambda R}(x)} \langle \omega|_{\eta_{x,\lambda}(y)}, \eta_{x,\lambda}^\# \overrightarrow{\partial T} \rangle d\mu_{\partial T} \\ &\leq C |\omega| \lambda^{1-n} \mu_{\partial T}(B_{\lambda R}(x)) \rightarrow 0 \text{ as } \lambda \downarrow 0 \end{aligned}$$

for  $\mathcal{H}^n$ -a.e.  $x \in M_j$  (independent of  $\omega$ ), because by applying (2) with  $d = n$  and  $\nu = \mu_{\partial T}$  we have

$$\Theta^{*n}(\mu_{\partial T}, x) = \limsup_{\lambda \downarrow 0} \lambda^{-n} \mu_{\partial T}(B_\lambda(x)) < \infty \text{ for } \mathcal{H}^n\text{-a.e. } x \in M_j.$$

Thus by (9) and (10), for any sequence  $\lambda_\ell \downarrow 0$ ,

$$\eta_{x,\lambda_\ell}^\# T \rightharpoonup S_x, \quad \partial S_x = 0 \text{ for } \mathcal{H}^n\text{-a.e. } x \in M,$$

where  $S_x \in \mathcal{D}_n(\mathbb{R}^{n+k})$  is defined by

$$(11) \quad S_x(\omega) = \theta(x) \int_L \langle \omega(z), \xi(x) \rangle d\mathcal{H}^n(z),$$

$\omega \in \mathcal{D}^n(\mathbb{R}^{n+k})$ ,  $L = T_x M$ . We now claim that (11), taken together with the fact that  $\partial S_x = 0$ , implies that  $\xi(x)$  orients  $L$  (i.e.  $\xi|_x = \pm \tau_1 \wedge \cdots \wedge \tau_n$  with  $\tau_1, \dots, \tau_n$

an orthonormal basis for  $L$ ). To see this, assume (without loss of generality) that  $L = \mathbb{R}^n \times \{0\} \subset \mathbb{R}^{n+k}$  and select  $\omega \in \mathcal{D}^{n-1}(\mathbb{R}^{n+k})$  so that  $\omega(y) = y^j \varphi(y) dy^{i_1} \wedge \cdots \wedge dy^{i_{n-1}}$ , where  $y = (y^1, \dots, y^{n+k})$ ,  $j \geq n+1$ ,  $\{i_1, \dots, i_{n-1}\} \subset \{1, \dots, n+k\}$ , and  $\varphi \in C_c^\infty(\mathbb{R}^{n+k})$ . Then since  $y_j = 0$  on  $\mathbb{R}^n \times \{0\}$  we deduce, from (11) and the fact that  $\partial S_x = 0$ ,

$$\begin{aligned} 0 &= \partial S_x(\omega) = S_x(d\omega) = \theta(x) \int_L \varphi(y) \langle dy^j \wedge dy^{i_1} \wedge \cdots \wedge dy^{i_{n-1}}, \xi(x) \rangle d\mathcal{H}^n(y) \\ &= \theta(x) \int_L \varphi(y) \xi(x) \cdot (e_j \wedge e_{i_1} \wedge \cdots \wedge e_{i_{n-1}}) d\mathcal{H}^n(y). \end{aligned}$$

That is, since  $\varphi \in C_c^\infty(\mathbb{R}^{n+k})$  is arbitrary, we deduce that  $\xi(x) \cdot (e_j \wedge e_{i_1} \wedge \cdots \wedge e_{i_{n-1}}) = 0$  whenever  $j \geq n+1$  and  $\{i_1, \dots, i_{n-1}\} \subset \{1, \dots, n+k\}$ . Thus we must have (since  $|\xi(x)| = 1$ ),  $\xi(x) = \pm e_1 \wedge \cdots \wedge e_n$  as required.  $\square$

We can now give the proof of the Compactness Theorem 3.15. For convenience we first re-state the theorem.

**8.2 Theorem. (Federer-Fleming Compactness Theorem.)** *If  $\{T_j\} \subset \mathcal{D}_n(U)$  is a sequence of integer multiplicity currents with*

$$(\ddagger) \quad \sup_{j \geq 1} (\mathbb{M}_W(T_j) + \mathbb{M}_W(\partial T_j)) < \infty \quad \forall W \subset\subset U,$$

*then there is an integer multiplicity  $T \in \mathcal{D}_n(U)$  and a subsequence  $\{T_{j'}\}$  such that  $T_{j'} \rightarrow T$  in  $U$ .*

**Proof of 8.2:** First note that the theorem is trivial in case  $n = 0$ . Then assume  $n \geq 1$  and, as in inductive hypothesis, suppose the theorem is true with  $n-1$  in place of  $n$ . The Weak Polyhedral Approximation Theorem 6.2 applied to  $T_j$  also gives an integer multiplicity polyhedral weak approximation to  $\partial T_j$ , hence by applying the inductive hypothesis to  $\partial T_j$ , we see that  $\partial T_j$  is an integer multiplicity current for each  $j = 1, 2, \dots$

Note that if  $B_{\rho_0}(\xi) \subset U$ , then by 4.5(1), (2) and an argument like that in 7.7(2) of Ch. 2, we know that, for  $\mathcal{L}^1$ -a.e.  $r \in (0, \rho_0]$ ,  $\partial(T_j \llcorner B_r(\xi))$  are integer multiplicity and 8.2 ( $\ddagger$ ) holds with  $T_{j'} \llcorner B_r(\xi)$  in place of  $T_j$  for some subsequence  $\{j'\} \subset \{j\}$  (depending on  $r$ ) (which we can take to be the original sequence  $\{j\}$ ), and, again by the inductive hypothesis,

$$(1) \quad \partial(T \llcorner B_r(\xi)) \quad \text{is an integer multiplicity current for } \mathcal{L}^1\text{-a.e. } r > 0.$$

In particular (in view of the arbitrariness of the ball  $B_{\rho_0}(\xi)$ ), we can (and we shall) assume without loss of generality that  $\text{spt } T_j \subset B_R(0)$  for some fixed compact  $R > 0$ , and that  $U = \mathbb{R}^P$ .

Also, since we are now assuming  $U = \mathbb{R}^P$ ,  $\text{spt } T_j \subset B_R(0)$ , we know that  $0 \ll \partial T - T$  has zero boundary and is the weak limit of  $0 \ll \partial T_j - T_j$ ; since  $0 \ll \partial T$  is integer multiplicity (by the inductive hypothesis) we thus see that the general case of the theorem follows from the special case when  $\partial T = 0$ . We shall therefore henceforth also assume  $\partial T = 0$ .

Next, define (for  $\xi \in \mathbb{R}^P$  fixed)

$$f(r) = \mathbb{M}(T \llcorner B_r(\xi)), \quad r > 0.$$

By virtue of 4.9 we have (since  $\partial T = 0$ )

$$(2) \quad \mathbb{M}(\partial(T \llcorner B_r(\xi))) \leq f'(r), \quad \mathcal{L}^1\text{-a.e. } r > 0.$$

(Notice that  $f'(r)$  exists a.e.  $r > 0$  because  $f(r)$  is increasing.) On the other hand if  $\Theta^{*n}(\mu_T, \xi) < \eta^n / \omega_n$  ( $\eta > 0$  a given constant), then  $\limsup_{\rho \downarrow 0} \frac{f(\rho)}{\omega_n \rho^n} < \eta^n$ , and hence for each  $\delta > 0$  we can arrange

$$(3) \quad \frac{d}{dr}(f^{1/n}(r)) \leq 2\eta$$

for a set of  $r \in (0, \delta)$  of positive  $\mathcal{L}^1$ -measure. (Because  $\frac{1}{\delta} \int_0^\delta \frac{d}{dr}(f^{1/n}(r)) dr \leq \delta^{-1} f^{1/n}(\delta) \leq \eta$  for all sufficiently small  $\delta > 0$ .)

Now by (1) and the Isoperimetric Theorem, we can find an integer multiplicity  $S_r \in \mathcal{D}_n(\mathbb{R}^P)$  such that  $\partial S_r = \partial(T \llcorner B_r(\xi))$  and

$$(4) \quad \begin{aligned} \mathbb{M}(S_r)^{\frac{n-1}{n}} &\leq c \mathbb{M}(\partial(T \llcorner B_r(\xi))) \\ &\leq c \eta \mathbb{M}(T \llcorner B_r(\xi))^{\frac{n-1}{n}} \quad (\text{by (2), (3)}) \end{aligned}$$

for a set of  $r$  of positive  $\mathcal{L}^1$ -measure in  $(0, \delta)$ .<sup>5</sup> Since  $\delta$  was arbitrary we then have both (1), (4) for a sequence of  $r \downarrow 0$ . But then (since we can repeat this for any  $\xi$  such that  $\Theta^{*n}(\mu_T, \xi) < \eta$ ) if  $K$  is any compact subset of  $\{x \in \mathbb{R}^P : \Theta^{*n}(\mu_T, x) < \eta\}$ , by Remark 4.4(3) of Ch. 1, for each given  $\rho > 0$  we get a pairwise disjoint family  $B_j = B_{\rho_j}(\xi_j)$  of closed balls covering  $\mu_T$ -almost all of  $K$ , with

$$\cup_j B_j \subset \{x : \text{dist}(x, K) < \rho\}$$

and with

$$(5) \quad \mathbb{M}(S_j^{(\rho)}) \leq c \eta \mathbb{M}(T \llcorner B_j)$$

<sup>5</sup>In case  $n = 1$ , (1), (2), (3) (for  $\eta < \frac{1}{4}$ ) imply  $\partial(T \llcorner B_r(\xi)) = 0$ , hence we get, in place of (4),  $\mathbb{M}(S_r) \leq \mathbb{M}(T \llcorner B_r(\xi))$  trivially by taking  $S_r = 0$ .

for some integer multiplicity  $S_j^{(\rho)}$  with

$$(6) \quad \partial S_j^{(\rho)} = \partial(T \llcorner B_j).$$

Now because of (6) we have  $S_j^{(\rho)} - T \llcorner B_j = \partial(\{\xi_j\} \times (S_j^{(\rho)} - T \llcorner B_j))$ , and hence (by 2.34, 2.37) we have for  $\omega \in \mathcal{D}^n(\mathbb{R}^P)$

$$\begin{aligned} 8.3 \quad | (S_j^{(\rho)} - T \llcorner B_j)(\omega) | &\leq c\rho \mathbb{M}(S_j^{(\rho)} - T \llcorner B_j) |d\omega| \\ &\leq c\rho \mathbb{M}(T \llcorner B_j) |d\omega| \quad (\text{by (5)}). \end{aligned}$$

Therefore we have  $\sum_j (S_j^{(\rho)} - T \llcorner B_j) \rightarrow 0$  as  $\rho \downarrow 0$ , and since the series  $\sum_j S_j^{(\rho)}$  and  $\sum_j T \llcorner B_j$  are  $\mathbb{M}$ -absolutely convergent (by (5) and the fact that the  $B_j$  are disjoint) we thus have

$$(7) \quad (T - \sum_j T \llcorner B_j) + \sum_j S_j^{(\rho)} \rightarrow T$$

as  $\rho \downarrow 0$ . Using (5) again, together with the lower semi-continuity of  $\mathbb{M}_W$  ( $W$  open) under weak convergence, we then have

$$\begin{aligned} (8) \quad \mu_T(\{x : \text{dist}(x, K) < \rho\}) \\ \leq \mu_T(\{x : \text{dist}(x, K) < \rho\} \setminus K) + c\eta \mu_T(\{x : \text{dist}(x, K) < \rho\}). \end{aligned}$$

Choosing  $\eta$  such that  $c\eta \leq \frac{1}{2}$ , this gives

$$\mu_T(\{x : \text{dist}(x, K) < \rho\}) \leq 2\mu_T(\{x : \text{dist}(x, K) < \rho\} \setminus K)$$

Letting  $\rho \downarrow 0$ , we get  $\mu_T(K) = 0$ .

Thus we have shown that  $\Theta^{*n}(\mu_T, x) > 0$  for  $\mu_T$ -a.e.  $x \in \mathbb{R}^P$ . We can therefore apply 8.1 in order to conclude that  $T = \tau(M, \theta, \xi)$  as in 8.1. It thus remains only to prove that  $\theta$  is integer-valued. This is achieved as follows:

First note that for  $\mathcal{L}^n$ -a.e.  $x \in M$  we have (Cf. the argument leading to (9) in the proof of Theorem 8.1)

$$(9) \quad \eta_{x, \lambda \#} T \rightarrow \theta(x) \llbracket T_x M \rrbracket \quad \text{as } \lambda \downarrow 0,$$

where  $\llbracket T_x \rrbracket$  is oriented by  $\xi(x)$ . Assuming without loss of generality that  $T_x M = \mathbb{R}^n \times \{0\} \subset \mathbb{R}^P$  and setting  $d(y) = \text{dist}(y, \mathbb{R}^n \times \{0\})$ , by 4.5(1) we can find a sequence  $\lambda_j \downarrow 0$  and a  $\rho > 0$  such that the slice  $\langle \eta_{x, \lambda_j \#} T, d, \rho \rangle$  (notation as in §4) is integer multiplicity with

$$(10) \quad \mathbb{M}_W(\langle \eta_{x, \lambda_j \#} T, d, \rho \rangle) \leq c \quad (\text{independent of } j)$$

where  $W = \check{B}_1^n(0) \times \mathbb{R}^{P-n} \subset \mathbb{R}^P$ . Next, we choose  $\{j'\} \subset \{j\}$  and  $\rho > 0$  so that  $\eta_{x, \lambda_j \# T_{j'}} \rightharpoonup \theta(x) \llbracket T_x M \rrbracket$  (which is possible by (9) and the fact that  $T_j \rightharpoonup T$ ), and so that (10) remains valid with  $T_{j'}$  instead of  $T$  (which is justified by 4.5(1) and a selection arguments in 7.7(2) of Ch. 2). Then by 4.5(2) we have  $S_j \equiv (\eta_{x, \lambda_j \# T_j}) \llcorner \{y : d(y) < \rho\}$  is such that

$$(11) \quad \sup_{j \geq 1} (\mathbb{M}_W(S_j) + \mathbb{M}_W(\partial S_j)) < \infty$$

with  $W = \check{B}_1^n(0) \times \mathbb{R}^{P-n} \subset \mathbb{R}^P$ . Now let  $p$  denote the restriction to  $W$  of the orthogonal projection of  $\mathbb{R}^P$  onto  $\mathbb{R}^n$ ; and let  $\tilde{S}_j$  be the current in  $\mathcal{D}_n(W)$  obtained by setting  $\tilde{S}_j(\omega) = S_j(\tilde{\omega})$ ,  $\omega \in \mathcal{D}^n(W)$ ,  $\tilde{\omega} \in \mathcal{D}^n(\mathbb{R}^P)$  such that  $\tilde{\omega} = \omega$  in  $W$  and  $\tilde{\omega} \equiv 0$  on  $\mathbb{R}^P \setminus W$ . Then we have  $p_\# \tilde{S}_j \in \mathcal{D}_n(B_1^n(0))$ , and hence, by 2.42 and (11) above,

$$p_\# \tilde{S}_j(\omega) = \int_{B_1^n(0)} a \theta_j d\mathcal{L}^n, \quad \omega = a dx^1 \wedge \cdots \wedge dx^n, \quad a \in C_c^\infty(\mathbb{R}^n),$$

for some integer-valued  $BV_{\text{loc}}(B_1^n(0))$  function  $\theta_j$  with

$$(12) \quad \begin{cases} \mathbb{M}_{\check{B}_1^n(0)}(p_\# \tilde{S}_j) = \int_{B_1^n(0)} |\theta_j| d\mathcal{L}^n \\ \mathbb{M}_{\check{B}_1^n(0)}(\partial p_\# \tilde{S}_j) = \int_{B_1^n(0)} |D\theta_j|. \end{cases}$$

Then by (11), (12) we deduce  $\int_{B_1^n(0)} |D\theta_j| + \int_{B_1^n(0)} |\theta_j| d\mathcal{L}^n \leq c$ ,  $c$  independent of  $j$ , and hence by the Compactness Theorem 2.6 of Ch. 2 we know  $\theta_j$  converges strongly in  $L^1$  in  $B_1^n(0)$  to an integer-valued  $BV$  function  $\theta_*$ . On the other hand  $S_j \rightharpoonup \theta(x) \llbracket \mathbb{R}^n \times \{0\} \rrbracket$  by (9), and hence  $p_\# \tilde{S}_j \rightharpoonup \theta(x) p_\# \llbracket \mathbb{R}^n \times \{0\} \rrbracket = \theta(x) \llbracket \mathbb{R}^n \rrbracket$  in  $B_1^n(0)$ . We thus deduce that  $\theta_* \equiv \theta(x)$  in  $B_1^n(0)$ ; thus  $\theta(x) \in \mathbb{Z}$  as required.  $\square$

## CHAPTER 6 PROBLEMS

6.1 Let  $\gamma, t(\tau), \tilde{\gamma}$  be as in Q3.1 of Ch.3 problems.

(i) If  $h : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is continuous, prove that the change of variable formula

$$\int_a^b h(\gamma(x)) \cdot \gamma'(x) dx = \int_0^L h(\tilde{\gamma}(\tau)) \cdot \tilde{\gamma}'(\tau) d\tau$$

is valid at least when  $\exists \varepsilon > 0$  with  $|\gamma'(x)| \geq \varepsilon$  for a.e.  $x \in [a, b]$ .

Hint: Show in this case that  $\tau \in [0, L] \mapsto t(\tau) \in [a, b]$  is a 1:1, increasing, Lipschitz map of  $[0, L]$  onto  $[a, b]$ , and hence (by the 1-dimensional area formula) we have  $\int_a^b f(x) dx = \int_0^L f(t(\tau)) t'(\tau) d\tau$  for each  $L^1([a, b])$  function  $f$ .

(ii) Prove the formula in (i) is valid without the assumption  $|\gamma'(x)| \geq \varepsilon$ .

Hint: To start, apply (i) to  $\gamma_\varepsilon(t) = (\gamma(t), \varepsilon t) : [a, b] \rightarrow \mathbb{R}^{n+1}$  and note that  $\tilde{\gamma}_\varepsilon(\tau) \rightarrow \tilde{\gamma}(\tau)$  on  $[0, L]$ —because if  $t_\varepsilon(\tau), \gamma_\varepsilon(\tau), L_\varepsilon$  are the analogues of  $t(\tau), \gamma(\tau), L$  when we use  $\gamma_\varepsilon$  in place of  $\gamma$ , then, as  $\varepsilon \downarrow 0$ ,  $L_\varepsilon \downarrow L$  and  $t_\varepsilon(\tau)$  increases to some limit  $s(\tau)$  with  $\text{length}(\gamma|[a, s(\tau)]) = \tau$ , hence  $s(\tau) \leq t(\tau)$ ,  $\gamma[s(\tau), t(\tau)] = \text{const.}$ , and  $\tilde{\gamma}_\varepsilon(\tau) = \gamma_\varepsilon(t_\varepsilon(\tau)) \rightarrow \gamma(s(\tau)) = \gamma(t(\tau)) = \tilde{\gamma}(\tau)$ .

6.2 Using the result of 6.1(ii) above, prove that if  $\gamma_j : [a_j, b_j] \rightarrow \mathbb{R}^n$  is a sequence as in Q2.1 of Ch.2 problems, then there is a Lipschitz  $\gamma : [0, 1] \rightarrow \mathbb{R}^n$  and a subsequence  $\gamma_{j'}$  such that

$$\int_{a_{j'}}^{b_{j'}} \langle \omega|_{\gamma_{j'}(x)}, \gamma_{j'}'(x) \rangle dx \rightarrow \int_0^1 \langle \omega|_{\gamma(x)}, \gamma'(x) \rangle dx$$

for each continuous 1-form  $\omega = \sum_{i=1}^n \omega_i dx^i$  on  $\mathbb{R}^n$ , where  $\langle \omega, v \rangle$  denotes the dual pairing between 1-forms and vectors in  $\mathbb{R}^n$ .

Note: The above is a version of the 1-dimensional case of the Federer-Fleming compactness theorem for integral currents, because  $\int_{a_{j'}}^{b_{j'}} \langle \omega|_{\gamma_{j'}(x)}, \gamma_{j'}'(x) \rangle dx = T_{j'}(\omega)$ , where  $T_j$  is the 1-dimensional integer multiplicity current given by  $T_j = \gamma_j \# \llbracket [a_j, b_j] \rrbracket$ ; by the area formula this can be represented  $T_j(\omega) = \int_{\Gamma_j} \langle \omega(y), \eta_j(y) \rangle N(\gamma_j, y) d\mathcal{H}^1(y)$ , where  $\Gamma_j = \gamma_j([a_j, b_j])$  and  $\eta_j$  is a Borel measurable unit vector function on  $\Gamma_j$  which orients the approximate tangent space of  $\Gamma_j$  for  $\mathcal{H}^1$ -a.e.  $y \in \Gamma_j$ .

6.3 Let  $V$  be a convex open subset of  $\mathbb{R}^Q$ ,  $U$  is open in  $\mathbb{R}^P$ ,  $f, g : U \rightarrow V$  are proper  $C^\infty$  maps, and  $h : [0, 1] \times U \rightarrow V$  is the affine homotopy  $h(t, x) = tg(x) + (1-t)f(x)$ .

(i) Show that  $h$  is not proper if  $g(x) = -f(x)$  for each  $x \in U$ .

(ii) If, in addition to the assumptions above,  $U$  is bounded and  $f = \bar{f}|U$ ,  $g = \bar{g}|U$  with  $\bar{f}, \bar{g}$  continuous on  $\bar{U}$ , prove that  $h$  is proper if  $\bar{f} = \bar{g}$  on  $\partial U$ .

Hint: By definition of proper (i.e. the preimage of each compact set in  $V$  is a compact set in  $U$ ),  $f : U \rightarrow V$  is proper if and only if the following property holds: Whenever  $\{x_k\} \subset U$  with either  $|x_k| \rightarrow \infty$  or  $\text{dist}(x_k, \partial U) \rightarrow 0$ , then either  $|f(x_k)| \rightarrow \infty$  or  $\text{dist}(f(x_k), \partial V) \rightarrow 0$ .

6.4 In lecture we proved that if  $T \in \mathcal{D}_n(U)$  with  $\mathbb{M}(T) < \infty, \mathbb{M}(\partial T) < \infty$ , if  $f, g : U \rightarrow V$  are  $C^1$  maps such that  $f|_{\text{spt } T}, g|_{\text{spt } T}$  are proper, and if  $f|_{\text{spt } T} = g|_{\text{spt } T}$ , then  $f_{\#}T = g_{\#}T$ .

Give an example to show that this may fail without the condition  $\mathbb{M}(\partial T) < \infty$ .

Hint: Let  $T \in \mathcal{D}_1(\mathbb{R})$  be defined by  $T(\omega) = a(0)$  for any 1-form  $\omega = a dx^1 \in \mathcal{D}^1(\mathbb{R})$ .

6.5 Check the claim 2.20 of Ch.6 of the text: That by applying 2.17, 2.18, one can check that  $\mathbb{M}_{W_1 \times W_2}(S \times T) = \mathbb{M}_{W_1}(S) \mathbb{M}_{W_2}(T)$ , assuming  $S, T \in \mathcal{D}_n(U)$  have locally finite mass in  $U$ .

6.6 Suppose  $R \in \mathcal{D}_n(\mathbb{R}^n)$  (i.e. we are in the setting  $P = n$  and  $U$  = the whole Euclidean space), and suppose  $R$  is an integer multiplicity current of finite mass.

(i) Prove that there are pairwise disjoint Lebesgue measurable subsets  $U_j, j = \pm 1, \pm 2, \dots$ , of  $\mathbb{R}^n$  such that  $R = \sum_{j=1}^{\infty} j \llbracket U_j \rrbracket - \sum_{j=1}^{\infty} j \llbracket U_{-j} \rrbracket$  and  $\mathbb{M}(R) = \sum_{j=1}^{\infty} j (\mathcal{L}^n(U_j) + \mathcal{L}^n(U_{-j}))$ .

(ii) If  $V_j = \cup_{k=j}^{\infty} U_k$  for  $j = 1, 2, \dots$  and  $W_j = \cup_{k=1+j}^{\infty} U_{-k}$  for  $j = 0, 1, \dots$  (note that then  $V_{j+1} \subset V_j$  and  $W_j \subset W_{j-1}$  for each  $j = 1, 2, \dots$ ), prove that  $R = \sum_{j=1}^{\infty} \llbracket V_j \rrbracket - \sum_{j=0}^{\infty} \llbracket W_j \rrbracket$  and  $\mathbb{M}(R) = \sum_{j=1}^{\infty} (\mathcal{L}^n(V_j) + \mathcal{L}^n(W_j))$ .

Hint: If  $a_j \geq 0$  for each  $j = 1, 2, \dots$  and  $b_i = \sum_{j=i}^{\infty} a_j$ , then  $\sum_{i=1}^{\infty} b_i = \sum_{j=1}^{\infty} j a_j$ .

(iii) If  $V_j, j = 1, 2, \dots, W_j, j = 0, 1, 2, \dots$  are as in (ii), and if  $V_{-j} = \mathbb{R}^n \setminus W_j$  for  $j = 0, 1, 2, \dots$  (so now  $V_{j+1} \subset V_j$  for all  $j = 0, \pm 1, \dots$ ) prove that  $\partial R = \sum_{j=-\infty}^{\infty} \partial \llbracket V_j \rrbracket$  (and the sum makes sense—i.e.  $\omega \in \mathcal{D}^{n-1}(\mathbb{R}^n) \Rightarrow \sum_{j=-\infty}^{\infty} \llbracket V_j \rrbracket(d\omega)$  is a convergent series).

6.7 If  $T \in \mathcal{D}_{n-1}(\mathbb{R}^n)$  is an integer multiplicity current with  $\text{spt } T$  compact,  $\mathbb{M}(T) < \infty$  and  $\partial T = 0$ , prove that the cone  $0 \times T$  (defined as in § 2 of Ch.6 of the text) is also integer multiplicity of finite mass. Hence, using the result of Q.6.6 above, prove that there is a sequence  $V_j, j = 0, \pm 1, \pm 2, \dots$ , of Lebesgue measurable sets with  $V_{j+1} \subset V_j$  for each  $j$  such that  $T = \sum_{j=-\infty}^{\infty} \partial \llbracket V_j \rrbracket$  and  $\mathbb{M}(T) \leq \sum_j \mathbb{M}(\partial \llbracket V_j \rrbracket)$ .

Remark: Actually equality holds in the last inequality, i.e.  $\mathbb{M}(T) = \sum_j \mathbb{M}(\partial \llbracket V_j \rrbracket)$ , but we are not quite in a position to prove this because we skipped the discussion of sets of locally finite perimeter (§ 4 of Ch.3 of the text).

Suggestion: Take a look at a few examples of the case when  $T = \gamma_{\#}([0, 1])$ , where  $\gamma$  is a  $C^1$  immersion of  $[0, 1]$  into  $\mathbb{R}^2$  with  $\gamma(0) = \gamma(1)$  (which ensures  $\partial T = 0$ ); it is interesting to see how the sets  $V_j$  work out in such cases, when  $\gamma$  has a few self-intersections.

6.8 (Degree of a mapping.) Suppose  $U$  is bounded open in  $\mathbb{R}^n$  and  $f : \bar{U} \rightarrow \mathbb{R}^n$  is continuous with  $f|_U \in C^1(U, \mathbb{R}^n)$ , let  $\tilde{U} = U \setminus f^{-1}(f(\partial U)) \neq \emptyset$ , and  $\tilde{f} = f|_{\tilde{U}}$ . Also, let  $W_1, W_2, \dots, W_N$  or  $W_1, W_2, \dots$  denote the connected components of  $\mathbb{R}^n \setminus f(\partial U)$  (the latter alternative in the event there are infinitely many connected components).

(i) Prove that  $\tilde{f} : \tilde{U} \rightarrow \mathbb{R}^n \setminus f(\partial U)$  is proper.

(ii) Prove  $\tilde{f}_\#[\tilde{U}] = \sum_j d_j [W_j]$ , where (a)  $d_j \in \mathbb{Z}$  for each  $j$ , and (b)  $d_j \neq 0 \Rightarrow W_j \subset f(U)$ .

(iii) Prove furthermore that (a)  $|d_j| \leq \mathcal{H}^0(f^{-1}y)$  for  $\mathcal{L}^n$ -a.e.  $y \in W_j$ , (b)  $|d_j|$  is congruent to  $\mathcal{H}^0(f^{-1}y) \bmod 2$  for  $\mathcal{L}^n$ -a.e.  $y \in W_j$ , and (c)  $d_j = \mathcal{H}^0(f^{-1}y)$  for any point  $y \in W_j$  such that  $J_f(x) > 0$  for each  $x \in f^{-1}y$ .

Hint:  $\tilde{f}_\#[\tilde{U}](\omega) = \int_{\tilde{U}} \langle df|_x \omega|_{f(x)}, e_1 \wedge \dots \wedge e_n \rangle d\mathcal{L}^n = \int_{\tilde{U}} \langle \omega|_{f(x)}, df|_{x\#}(e_1 \wedge \dots \wedge e_n) \rangle d\mathcal{L}^n$ ; compute using area formula.

Note: For  $y \in W_j$ ,  $d_j$  is called the (topological) degree of the map  $f$  at  $y$ , denoted  $d(f, U, y)$ .

6.9 (Invariance of degree under homotopy.)

(i) If  $U, f, W_j, d_j$  are as in Q.6.8,  $K$  compact  $\supset f(\partial U)$ , and  $V = U \setminus f^{-1}K \neq \emptyset$ , then  $f_\#[V] = \sum_i c_i [E_i]$ , where  $E_i$  are the connected components of  $\mathbb{R}^n \setminus K$ , and each  $E_i \subset$  some  $W_j$  with  $c_i = d_j$ .

(ii) If  $U$  is bounded open in  $\mathbb{R}^n$  and  $h : [0, 1] \times \bar{U} \rightarrow \mathbb{R}^n$  is continuous, if  $f_t : U \rightarrow \mathbb{R}^n$  is  $C^1$  for each  $t \in [0, 1]$ , with  $f_t(x) = h(t, x)$  for  $x \in U$ , prove that  $d(f_0, U, y) = d(f_1, U, y) \forall y \in \mathbb{R}^n \setminus h([0, 1] \times \partial U)$ .

Hint: Show  $d(f_t, U, y)$  is a continuous function of  $t$  for  $y \in \mathbb{R}^n \setminus h([0, 1] \times \partial U)$ .

6.10 Let  $P(z) = z^n + \sum_{j=0}^{n-1} a_j z^j$ , where  $a_0, \dots, a_{n-1}$  are given complex constants. Prove, using Q.6.9 above, the ‘‘Fundamental Theorem of Algebra’’ that  $P(z) = 0$  has a root (hence  $n$  roots by the Remainder Theorem and induction on  $n$ ).

Hint: View  $P$  as a map of  $\mathbb{R}^2 \rightarrow \mathbb{R}^2$  and take  $U = \{(x, y) : |(x, y)| < R\}$  (i.e.  $\{z : |z| < R\}$ ). Let  $P_0(z) = z^n$  (as a map  $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ ) and prove (by direct computation with the aid of the area formula) that  $P_{0\#}[U] = n[U]$ . Then show problem 6.9 is applicable with homotopy  $h(t, x, y) = z^n + t \sum_{j=0}^{n-1} a_j z^j$ ,  $z = x + iy$ , if  $R$  is large enough.

6.11 (i) (Constancy theorem on a submanifold.) Suppose  $M$  is a connected oriented  $n$ -dimensional submanifold of  $\mathbb{R}^P$  and let  $U \subset \mathbb{R}^P$  be open with  $M \subset U$ . If  $T \in \mathcal{D}_n(U)$  with  $\text{spt } T \subset M$ ,  $\mathbb{M}(T) < \infty$ ,  $\vec{T}|_x \in \Lambda_n(T_x M)$  for  $\mu_T$ -a.e.  $x \in M$ , and  $\partial T = 0$ , prove that  $T = c [M]$  for some real constant  $c$ .

Hint: Use local coordinates and the constancy theorem (proved in lecture) in  $\mathbb{R}^n$ .

(ii) Give an example to show that the above is false without the hypothesis  $\vec{T}|_x \in \Lambda_n(T_x M)$ .

Hint: Consider  $M = \{0\} \times \mathbb{R} \subset \mathbb{R}^2$ .

6.12 (Degree of a map between  $n$ -dimensional manifolds.) (i) If  $M, N$  are compact oriented connected  $C^1$  submanifolds (without boundary) of  $\mathbb{R}^P$  and  $\mathbb{R}^Q$  respectively, and if  $f : M \rightarrow N$  is  $C^1$ , prove (using 6.11(i) above)  $f_{\#} \llbracket M \rrbracket = d \llbracket N \rrbracket$  for some  $d \in \mathbb{Z}$ .

Note:  $d = d(f)$  is called the degree of the map  $f : M \rightarrow N$ .

(ii) If  $M, N$  are as in (i) and if  $h : [0, 1] \times M \rightarrow N$  is continuous with  $f_t = M \rightarrow N$  of class  $C^1$  for each  $t \in [0, 1]$ , where  $f_t(x) = h(t, x)$  for  $x \in M$ , prove that  $d(f_0) = d(f_1)$ .

6.13 Give an example to show that the Federer-Fleming compactness theorem (Theorem 3.11 of Ch.6 of the text) fails without the hypothesis that  $\sup_j \mathbb{M}_W(\partial T_j) < \infty$ .

Hint: Modify the example in Q 2.11 of Ch.2; instead of vertical cylinders use the vertical strips  $\{(x, y, z) \in \mathbb{R}^3 : |x - i/N| < 1/N^2, y = j/N, z \in \mathbb{R}\}$  with orienting 2-vector  $e_1 \wedge e_3, i, j \in \{0, \pm 1, \pm 2, \dots\}$ .

Note: Even better, deleting the  $z$  coordinate in the same example (so the vertical strips become line segments with orienting 1-vector  $e_1$ ) gives a 1-dimensional sequence in  $\mathbb{R}^2$ .



# Chapter 7

## Area Minimizing Currents

§ 1	Basic Concepts .....	211
§ 2	Existence and Compactness Results .....	213
§ 3	Tangent Cones and Densities .....	217
§ 4	Some Regularity Results (Arbitrary Codimension) .....	223
§ 5	Codimension 1 Theory .....	226

### 1 Basic Concepts

Suppose  $A$  is any subset of  $\mathbb{R}^{n+k}$ ,  $A \subset U$ ,  $U$  open in  $\mathbb{R}^{n+k}$ , and  $T \in \mathcal{D}_n(U)$  an integer multiplicity current.

**1.1 Definition:** We say that  $T$  is minimizing in  $A$  if

$$\mathbb{M}_W(T) \leq \mathbb{M}_W(S)$$

whenever  $W \subset\subset U$ ,  $\partial S = \partial T$  (in  $U$ ) and  $\text{spt}(S - T)$  is a compact subset of  $A \cap W$ .

There are two especially important cases (in fact the only cases we are interested in here) of this definition:

- (1) when  $A = U$
- (2) when  $A = N$  with  $N \subset U$  an  $(n + k)$ -dimensional embedded  $C^2$  submanifold of  $\mathbb{R}^{n+k}$  (in the sense of §4 of Ch. 2).

Corresponding to the current  $T = \underline{\tau}(M, \theta, \xi) \in \mathcal{D}_n(U)$  we have the integer multiplicity varifold  $V = \underline{\nu}(M, \theta)$ . As one would expect,  $V$  is stationary in  $U$  if  $T$  is minimizing in  $U$  and  $\partial T = 0$ :

**1.2 Lemma.** *Suppose  $T$  is minimizing in  $N$ , where  $N$  is an  $(n + k)$ -dimensional  $C^2$  embedded submanifold of  $\mathbb{R}^{n+L}$  ( $L \geq k$ , so  $N = U$ , an open subset of  $\mathbb{R}^{n+k}$ , is an important special case) and suppose  $\partial T = 0$  in  $N$ . Then  $V$  is stationary in  $N$  in the sense of 2.6 of Ch. 4, so that in particular  $V$  has locally bounded generalized mean curvature in  $N$  (in the sense of 3.15 of Ch. 4).*

*In fact  $V$  is minimizing in  $N$  in the sense that*

$$(1) \quad \mathbb{M}_W(V) \leq \mathbb{M}_W(\varphi_\# V),$$

*whenever  $W$  is open in  $N$  with  $W \subset\subset N$  and  $\varphi$  is a  $C^1$  diffeomorphism of  $N$  such that  $\varphi(N) \subset N$  and  $\varphi|_{N \setminus K} = \text{id}_{N \setminus K}$  for some compact  $K \subset W$ .*

**1.3 Remark:** In view of 1.2 (together with the fact that  $\theta \geq 1$ ) we can represent  $T = \underline{\tau}(M_*, \theta_*, \xi)$  where  $M_*$  is a relatively closed countably  $n$ -rectifiable subset of  $U$ , and  $\theta_*$  is an upper semi-continuous function on  $M_*$  with  $\theta_* \geq 1$  everywhere on  $M_*$  (and  $\theta_*$  integer-valued  $\mathcal{H}^n$ -a.e. on  $M_*$ ).

**Proof of 1.2:** Evidently (in view of the discussion of §2 of Ch. 4) the first claim in 1.2 follows from 1.2 (1) (by taking  $\varphi = \varphi_t$  in 1.2 (1),  $\varphi_t$  is in 2.5 of Ch. 4).

To prove 1.2 (1) we first note that, for any  $W$ ,  $\varphi$  as in the statement of the theorem,

$$(1) \quad \mathbb{M}_W(\varphi_\# V) = \mathbb{M}_W(\varphi_\# T)$$

by 3.2(3) of Ch. 6. Also, since  $\partial T = 0$  (in  $U$ ), we have

$$(2) \quad \partial \varphi_\# T = \varphi_\# \partial T = 0.$$

Finally,

$$(3) \quad \text{spt}(T - \varphi_\# T) \subset K \subset W.$$

By virtue of (2), (3) we are able to use the inequality of 1.1 with  $S = \varphi_\# T$ . This gives 1.2 (1) as required by virtue of (1).  $\square$

We conclude this section with the following useful decomposition lemma:

**1.4 Lemma.** *Suppose  $T_1, T_2 \in \mathcal{D}_n(U)$  are integer multiplicity and suppose  $T_1 + T_2$  is minimizing in  $A$ ,  $A \subset U$ , and*

$$\mathbb{M}_W(T_1 + T_2) = \mathbb{M}_W(T_1) + \mathbb{M}_W(T_2)$$

*for each  $W \subset\subset U$ . Then  $T_1, T_2$  are both minimizing in  $A$ .*

**Proof:** Let  $X \in \mathcal{D}_n(U)$  be integer multiplicity with  $\text{spt } X \subset K$ ,  $K$  a compact subset of  $A \cap W$ , and with  $\partial X = 0$ . Because  $T_1 + T_2$  is minimizing in  $A$  we have (by 1.1)

$$\mathbb{M}_W(T_1 + T_2 + X) \geq \mathbb{M}_W(T_1 + T_2).$$

However since  $\mathbb{M}_W(T_1 + T_2) = \mathbb{M}_W(T_1) + \mathbb{M}_W(T_2)$ , and  $\mathbb{M}(T_1 + T_2 + X) \leq \mathbb{M}_W(T_1 + X) + \mathbb{M}_W(T_2)$ , this gives

$$\mathbb{M}_W(T_1) \leq \mathbb{M}_W(T_1 + X).$$

In view of the arbitrariness of  $X$ , this establishes that  $T_1$  is minimizing in  $A \cap W$  (in accordance with 1.1). Interchanging  $T_1, T_2$  in the above argument, we likewise deduce that  $T_2$  is minimizing in  $A \cap W$ .  $\square$

## 2 Existence and Compactness Results

We begin with a result which establishes the rich abundance of area minimizing currents in Euclidean space.

**2.1 Lemma.** *Let  $S \in \mathcal{D}_{n-1}(\mathbb{R}^{n+k})$  be integer multiplicity with  $\text{spt } S$  compact and  $\partial S = 0$ . Then there is an integer multiplicity current  $T \in \mathcal{D}_n(\mathbb{R}^{n+k})$  such that  $\text{spt } T$  is compact and  $\mathbb{M}(T) \leq \mathbb{M}(R)$  for each integer multiplicity  $R \in \mathcal{D}_n(\mathbb{R}^{n+k})$  with  $\text{spt } R$  compact and  $\partial R = S$ .*

**2.2 Remarks:** (1) Of course  $T$  is minimizing in  $\mathbb{R}^{n+k}$  in the sense of 1.1.

(2) By virtue of 1.2 and the convex hull property (Theorem 7.2 of Ch. 4) we have automatically that  $\text{spt } T \subset \text{convex hull of spt } S$ .

(3)  $\mathbb{M}(T)^{\frac{n-1}{n}} \leq c\mathbb{M}(S)$  by virtue of the Isoperimetric Inequality 6.1 of Ch. 6.

**Proof of 2.1:** Let

$$\mathcal{I}_S = \left\{ R \in \mathcal{D}_n(\mathbb{R}^{n+k}) : R \text{ is integer multiplicity, } \text{spt } R \text{ compact, } \partial R = S \right\}.$$

Evidently  $\mathcal{I}_S \neq \emptyset$ . (e.g.  $0 \# S \in \mathcal{I}_S$ .) Take any sequence  $\{R_q\} \subset \mathcal{I}_S$  with

$$(1) \quad \lim_{q \rightarrow \infty} \mathbb{M}(R_q) = \inf_{R \in \mathcal{I}_S} \mathbb{M}(R),$$

let  $B_R(0)$  be any ball in  $\mathbb{R}^{n+k}$  such that  $\text{spt } S \subset B_R(0)$ , and let  $f : \mathbb{R}^{n+k} \rightarrow B_R(0)$  be the nearest point (radial) retract of  $\mathbb{R}^{n+k}$  onto  $B_R(0)$ . Then  $\text{Lip } f = 1$  and hence

$$(2) \quad \mathbb{M}(f \# R_q) \leq \mathbb{M}(R_q).$$

On the other hand  $\partial f \# R_q = f \# \partial R_q = f \# S = S$ , because  $f|_{B_R(0)} = \text{id}_{B_R(0)}$  and  $\text{spt } S \subset B_R(0)$ . Thus  $f \# R_q \in \mathcal{I}_S$  and by (1), (2) we have

$$(3) \quad \lim_{q \rightarrow \infty} \mathbb{M}(f \# R_q) = \inf_{R \in \mathcal{I}_S} \mathbb{M}(R).$$

Now by the Compactness Theorem 3.15 of Ch. 6 there is a subsequence  $\{q'\} \subset \{q\}$  and an integer multiplicity current  $T \in \mathcal{D}_n(\mathbb{R}^{n+k})$  such that  $f_{\#}R_{q'} \rightarrow T$  and (by (3) and lower semi-continuity of mass with respect to weak convergence)

$$(4) \quad \mathbb{M}(T) \leq \inf_{R \in \mathcal{I}_S} \mathbb{M}(R).$$

However  $\text{spt } T \subset B_R(0)$  and  $\partial T = \lim \partial f_{\#}R_{q'} = \lim f_{\#}\partial R_{q'} = S$ , so that  $T \in \mathcal{I}_S$ , and the lemma is established (by (4)).  $\square$

The proof of the following lemma is similar to that of 2.1 (and again based on 3.15 of Ch. 6), and its proof is left to the reader.

**Lemma.** *Suppose  $N$  is an  $(n+k)$ -dimensional compact  $C^1$  embedded submanifold in  $\mathbb{R}^{n+k}$  and suppose  $R_1 \in \mathcal{D}_n(\mathbb{R}^{n+k})$  is given such that  $\partial R_1 = 0$ ,  $\text{spt } R_1 \subset N$  and*

$$\mathcal{I}_{R_1} = \left\{ R \in \mathcal{D}_n(\mathbb{R}^{n+k}) : R - R_1 = \partial S \text{ for some integer multiplicity } S \in \mathcal{D}_{n+1}(\mathbb{R}^{n+k}) \text{ with } \text{spt } S \subset N \right\} \neq \emptyset.$$

*Then there is  $T \in \mathcal{I}_{R_1}$  such that*

$$\mathbb{M}(T) = \inf_{R \in \mathcal{I}_{R_1}} \mathbb{M}(R).$$

**2.3 Remarks:** (1)  $R - R_1 = \partial S$  with  $S$  integer multiplicity and  $\text{spt } S \subset N$  means that  $R, R_1$  represents homologous cycles in the  $n$ -th singular homology class (with integer coefficients) of  $N$  (See [Fed69] or [FF60] for discussion.)

(2) It is quite easy to see that  $T$  is *locally* minimizing in  $N$ ; thus for each  $\xi \in \text{spt } T$  there is a neighborhood  $U$  of  $\xi$  such that  $T$  is minimizing in  $N \cap U$ .

We conclude this section with the following important compactness theorem for minimizing currents:

**2.4 Theorem.** *Suppose  $\{T_j\}$  is a sequence of minimizing currents in  $U$  with*

$$\sup_{j \geq 1} (\mathbb{M}_W(T_j) + \mathbb{M}_W(\partial T_j)) < \infty \text{ for each } W \subset\subset U,$$

*and suppose  $T_j \rightarrow T \in \mathcal{D}_n(U)$ . Then  $T$  is minimizing in  $U$  and  $\mu_{T_j} \rightarrow \mu_T$  (in the usual sense of Radon measures in  $U$ ).*

**2.5 Remarks:** (1) Note that  $\mu_{T_j} \rightarrow \mu_T$  means the corresponding sequence of varifolds converge in the measure theoretic sense of §1 of Ch. 4 to the varifold associated with  $T$ . ( $T$  is automatically integer multiplicity by 3.15 of Ch. 6.)

(2) If the hypotheses are as in the theorem, except that  $\text{spt } T_j \subset N_j \subset U$  and  $T_j$  is minimizing in  $N_j$ ,  $\{N_j\}$  a sequence of  $C^1$  embedded  $(n+k)$ -dimensional submanifolds

of  $\mathbb{R}^{n+k}$  converging in the  $C^1$  sense to  $N$ ,  $N \subset U$  an embedded  $(n+k)$ -dimensional  $C^1$  submanifold of  $\mathbb{R}^{n+k}$ ,<sup>1</sup> then  $T$  minimizes in  $N$  (and we still have  $\mu_{T_j} \rightarrow \mu_T$  in the sense of Radon measures in  $U$ ). We leave this modification of 2.4 to the reader. (It is easily checked by using suitable local representations for the  $N_j$  and by obvious modifications of the proof of 2.4 given below.)

**Proof of 2.4:** Let  $K \subset U$  be an arbitrary compact set and choose a smooth  $\varphi : U \rightarrow [0, 1]$  such that  $\varphi \equiv 1$  in some neighborhood of  $K$ , and  $\text{spt } \varphi \subset \{x \in U : \text{dist}(x, K) < \varepsilon\}$ , where  $0 < \varepsilon < \text{dist}(K, \partial U)$  is arbitrary. For  $0 < \lambda < 1$ , let

$$W_\lambda = \{x \in U : \varphi(x) > \lambda\}.$$

Then

$$(1) \quad K \subset W_\lambda \subset\subset U$$

for each  $\lambda$ ,  $0 \leq \lambda < 1$ .

By virtue of 7.2 of Ch. 6 we know that  $d_W(T_j, T) \rightarrow 0$  for each  $W \subset\subset U$ , hence in particular we have

$$(2) \quad T - T_j = \partial R_j + S_j, \quad \mathbb{M}_{W_0}(R_j) + \mathbb{M}_{W_0}(S_j) \rightarrow 0$$

$$(W_0 = \{x \in U : \varphi(x) > 0\}).$$

By the slicing theory (and in particular by 4.5 of Ch. 6) we can choose  $0 < \alpha < 1$  and a subsequence  $\{j'\} \subset \{j\}$  (subsequently denoted simply by  $\{j\}$ ) such that

$$(3) \quad \partial(R_j \llcorner W_\alpha) = (\partial R_j) \llcorner W_\alpha + P_j$$

where  $\text{spt } P_j \subset \partial W_\alpha$ ,  $P_j$  is integer multiplicity, and

$$(4) \quad \mathbb{M}(P_j) \rightarrow 0.$$

We can also of course choose  $\alpha$  to be such that

$$(5) \quad \mathbb{M}(T_j \llcorner \partial W_\alpha) = 0 \quad \forall j \text{ and } \mathbb{M}(T \llcorner \partial W_\alpha) = 0.$$

Thus, combining (2), (3), (4) we have

$$(6) \quad T \llcorner W_\alpha = T_j \llcorner W_\alpha + \partial \widetilde{R}_j + \widetilde{S}_j$$

---

<sup>1</sup>Thus  $\exists \psi_j : U \rightarrow U$ ,  $\psi_j|_{N_j}$  in a diffeomorphism onto  $N$ , and  $\psi_j \rightarrow \text{id}_U$  locally in  $U$  with respect to the  $C^1$  metric.

with  $\tilde{R}_j, \tilde{S}_j$  integer multiplicity ( $\tilde{R}_j = R_j \llcorner W_\alpha, \tilde{S}_j = S_j \llcorner W_\alpha + P_j$ ) with

$$(7) \quad \mathbb{M}(\tilde{R}_j) + \mathbb{M}(\tilde{S}_j) \rightarrow 0.$$

Now let  $X \in \mathcal{D}_n(U)$  be any integer multiplicity current with  $\partial X = 0$  and  $\text{spt } X \subset K$ . We want to prove

$$(8) \quad \mathbb{M}_{W_\alpha}(T) \leq \mathbb{M}_{W_\alpha}(T + X).$$

(In view of the arbitrariness of  $K, X$  this will evidently establish the fact that  $T$  is minimizing in  $U$ .)

By (6), we have

$$(9) \quad \begin{aligned} \mathbb{M}_{W_\alpha}(T + X) &= \mathbb{M}_{W_\alpha}(T_j + X + \partial \tilde{R}_j + \tilde{S}_j) \\ &\geq \mathbb{M}_{W_\alpha}(T_j + X + \partial \tilde{R}_j) - \mathbb{M}(\tilde{S}_j). \end{aligned}$$

Now since  $T_j$  is minimizing and  $\partial(X + \partial \tilde{R}_j) = 0$  with  $\text{spt}(X + \partial \tilde{R}_j) \subset \bar{W}_\alpha$ , we have

$$(10) \quad \mathbb{M}_{W_\lambda}(T_j + X + \partial \tilde{R}_j) \geq \mathbb{M}_{W_\lambda}(T_j)$$

for  $\lambda > \alpha$ . But by (3) we have  $\mathbb{M}(\partial \tilde{R}_j \llcorner \partial W_\alpha) = \mathbb{M}(P_j) \rightarrow 0$ , and by (5)  $\mathbb{M}(T_j \llcorner \partial W_\alpha) = 0, (T \llcorner \partial W_\alpha) = 0$ . Hence letting  $\lambda \downarrow 0$  in (10) we get

$$(11) \quad \mathbb{M}_{W_\alpha}(T_j + X + \partial \tilde{R}_j) \geq \mathbb{M}_{W_\alpha}(T_j) - \mathbb{M}(P_j),$$

and therefore from (9) we obtain

$$(12) \quad \mathbb{M}_{W_\alpha}(T + X) \geq \mathbb{M}_{W_\alpha}(T_j) - \varepsilon_j, \quad \varepsilon_j \downarrow 0.$$

In particular, setting  $X = 0$ , we have

$$(13) \quad \mathbb{M}_{W_\alpha}(T) \geq \mathbb{M}_{W_\alpha}(T_j) - \varepsilon_j, \quad \varepsilon_j \downarrow 0.$$

Using the lower semi-continuity of mass with respect to weak convergence in (12), we then have (8) as required.

It thus remains only to prove that  $\mu_{T_j} \rightarrow \mu_T$  in the sense of Radon measures in  $U$ . First note that by (13) we have

$$\limsup \mathbb{M}_{W_\alpha}(T_j) \leq \mathbb{M}_{W_\alpha}(T),$$

so that (since  $K \subset W_\alpha \subset \{x : \text{dist}(x, K) < \varepsilon\}$  by construction)

$$\limsup \mu_{T_j}(K) \leq \mathbb{M}_{\{x : \text{dist}(x, K) < \varepsilon\}}(T).$$

Hence, letting  $\varepsilon \downarrow 0$

$$(14) \quad \limsup \mu_{T_j}(K) \leq \mu_T(K).$$

(We actually only proved this for some subsequence, but we can repeat the argument for a subsequence of any given subsequence, hence it holds for the original sequence  $\{T_j\}$ .)

By the lower semi-continuity of mass with respect to weak convergence we have

$$(15) \quad \mu_T(W) \leq \liminf \mu_{T_j}(W) \quad \forall \text{ open } W \subset\subset U.$$

Since (14), (15) hold for arbitrary compact  $K$  and open  $W \subset U$ , it now easily follows (by a standard approximation argument) that  $\int f \, d\mu_{T_j} \rightarrow \int f \, d\mu_T$  for each continuous  $f$  with compact support in  $U$ , as required.  $\square$

### 3 Tangent Cones and Densities

In this section we prove the basic results concerning tangent cones and densities of area minimizing currents. All results depend on the fact that (by virtue of 1.2 the varifold associated with a minimizing current is stationary. This enables us to bring into play the important monotonicity results of §4 of Ch.4.

Subsequently we take  $N$  to be a smooth (at least  $C^2$ )  $(n+k)$ -dimensional embedded submanifold of  $\mathbb{R}^{n+L}$  ( $L \geq k$ ),  $U$  open in  $\mathbb{R}^{n+k}$  and  $(\bar{N} \setminus N) \cap U = \emptyset$ . Notice that an important case is when  $N = U$  (when  $L = k$ ).

**3.1 Theorem.** *Suppose  $T \in \mathcal{D}_n(U)$  is minimizing in  $U \cap N$ ,  $\text{spt } T \subset U \cap N$ , and  $x \in \text{spt } T \setminus \text{spt } \partial T$ . Then*

- (1)  $\Theta^n(\mu_T, x)$  exists everywhere in  $U$  and is an upper semi-continuous function of  $x \in U$ ;
- (2) For each  $x \in \text{spt } T$  and each sequence  $\{\lambda_j\} \downarrow 0$ , there is a subsequence  $\{\lambda_{j'}\}$  such that  $\eta_{x, \lambda_{j'} \# T} \rightarrow C$  and  $\mu_{\eta_{x, \lambda_{j'} \# T}} \rightarrow \mu_C$  in  $\mathbb{R}^{n+k}$ , where  $C \in \mathcal{D}_n(\mathbb{R}^{n+k})$  is integer multiplicity and minimizing in  $\mathbb{R}^{n+k}$ ,  $\eta_{0, \lambda \# C} = C \, \forall \lambda > 0$ , and  $\Theta^n(\mu_C, 0) = \Theta^n(\mu_T, x)$ .

**3.2 Remarks:** (1) If  $C$  is as in 3.1(2) above, we say that  $C$  is a *tangent cone* for  $T$  at  $x$ . If  $\text{spt } C$  is an  $n$ -dimensional subspace  $P$ . Notice that since  $C$  is integer multiplicity and  $\partial C = 0$ , it then follows from 2.41 of Ch.6 that, assuming we chose an appropriate (constant) orientation for  $P$ ,  $C = m \llbracket P \rrbracket$  for some  $m \in \{1, 2, \dots\}$ . In this case we call  $C$  a *multiplicity  $m$  tangent plane* for  $T$  at  $x$ .

(2) Notice that is *not* clear whether or not there is an *unique* tangent cone for  $T$  at  $x$ ; thus it is an open question whether or not  $C$  depends on the particular sequence  $\{\lambda_j\}$  or

subsequence  $\{\lambda_{j'}\}$  use in its definition. The work of [Sim83] shows that if  $C$  is a tangent cone of  $T$  at  $x$  such that  $\Theta^n(\mu_C, x) = 1$  for all  $x \in \text{spt } C \setminus \{0\}$ , then  $C$  is the unique tangent cone for  $T$  at  $x$ , and hence  $\eta_{x, \lambda\#} T \rightarrow C$  as  $\lambda \downarrow 0$ . Also B. White [Whi82] has shown in case  $n = 2$  that  $C$  is always unique with  $\text{spt } C$  consisting of a union of 2-planes.

**Proof of 3.1:** By virtue of 1.2 we can apply the monotonicity formula of 4.3 of Ch. 4 (with  $\alpha = 1$ ) and 4.7 of Ch. 4 in order to deduce that  $\Theta^n(\mu_T, x)$  exists for every  $x \in U$  and is an upper semi-continuous function of  $x$  in  $U$ .

Thus in particular

$$(1) \quad (\omega_n R^n)^{-1} \mathbb{M}_{\check{B}_R(0)}(\eta_{x, \lambda_j\#} T) = (\omega_n \lambda_j^n R^n)^{-1} \mathbb{M}_{\check{B}_{\lambda_j R}(x)}(T) \rightarrow \Theta^n(\mu_T, x)$$

for each  $R > 0$ , and hence  $\sup_j \mathbb{M}_{\check{B}_R(0)}(\eta_{x, \lambda_j\#} T) < \infty$  for each  $R > 0$ , while  $\partial \eta_{x, \lambda_j\#} T = 0$  in  $\check{B}_R(0)$  for sufficiently large  $j$  (because  $x \notin \text{spt } \partial T$ ), so we can apply the compactness theorem 2.4 to give a subsequence  $j'$  such  $\eta_{x, \lambda_{j'}\#} T \rightarrow C$  in  $\mathbb{R}^{n+k}$  with  $C$  integer multiplicity minimizing<sup>2</sup>, so

$$(2) \quad C = \tau(\text{spt } C, \xi, \Theta^n(\mu_C, \cdot)),$$

$\mu_{\eta_{x, \lambda_{j'}\#} T} \rightarrow \mu_C$  in  $\mathbb{R}^{n+k}$ , and (by Lemma 1.2) the rectifiable varifold

$$(3) \quad V_C = \underline{v}(\text{spt } C, \Theta^n(\mu_C, \cdot))$$

is stationary in  $\mathbb{R}^{n+k}$ . In particular for any  $\rho > 0$  with  $\mu_C(\partial B_\rho(0)) = 0$  (which is true except for at most a countable set of  $\rho$ ) we have

$$(4) \quad \mu_{\eta_{x, \lambda_{j'}\#} T}(B_\rho(0)) \rightarrow \mu_C(B_\rho(0)),$$

and together with (1) gives  $(\omega_n \rho^n)^{-1} \mu_C(B_\rho(0)) = \Theta^n(\mu_T, x)$  for each  $\rho > 0$ . Then by the monotonicity formula 3.8 of Ch. 4, applied to the stationary varifold  $V_C$  of (3), we have  $D^\perp r = 0$   $\mu_C$ -a.e., where  $r = |x|$ , and  $D^\perp r$  is orthogonal projection of  $Dr = r^{-1}x$  onto the normal space  $(T_x \text{spt } C)^\perp$ . That is  $x \in T_x \text{spt } C$  for  $\mu_C$ -a.e.  $x$ , so in particular  $x \wedge \vec{C} = 0$   $\mu_C$ -a.e. and hence we can apply Lemma 2.40 to deduce that  $C$  is a cone.  $\square$

**3.3 Theorem.**<sup>3</sup> Suppose  $T \in \mathcal{D}_n(U)$  is minimizing in  $U \cap N$ ,  $\text{spt } T \subset U \cap N$ , and  $\partial T = 0$  (in  $U$ ). Then

$$(1) \quad \Theta^n(\mu_T, x) \in \mathbb{Z} \text{ for all } x \in U \setminus E, \text{ where } \mathcal{H}^{n-3+\alpha}(E) = 0 \quad \forall \alpha > 0;$$

<sup>2</sup>See Remark 2.5; notice this establishes first that  $C$  is minimizing only in the  $(n+k)$ -dimensional subspace  $T_x N \subset \mathbb{R}^{n+k}$ . However since orthogonal projection of  $\mathbb{R}^{n+k}$  onto  $T_x N$  does not increase area, and since  $\text{spt } C \subset T_x N$ , it then follows that  $C$  is area minimizing in  $\mathbb{R}^{n+k}$  as claimed.

<sup>3</sup>Cf. Almgren [Alm84]

- (2) *There is a set  $F \subset E$  ( $E$  as in (1)) with  $\mathcal{H}^{n-2+\alpha}(F) = 0 \ \forall \alpha > 0$  and such that for each  $x \in \text{spt } T \setminus F$  there is a tangent plane (see 3.2(1) above for terminology) for  $T$  at  $x$ .*

**Note:** We do not claim  $E, R$  are *closed*.

The proof of both parts is based on the abstract dimension reducing argument of Appendix A. In order to apply this in the context of currents we need the observation of the following remark.

**3.4 Remark:** Given an integer multiplicity current  $S \in \mathcal{D}_n(\mathbb{R}^{n+k})$ , there is an associated function  $\varphi_S = (\varphi_S^0, \varphi_S^1, \dots, \varphi_S^N) : \mathbb{R}^{n+k} \rightarrow \mathbb{R}^{N+1}$ , where  $N = \binom{n+k}{n}$ , such that (writing  $\theta_S(x) = \theta^{*n}(\mu_S, x)$ )

$$\varphi_S^0(x) = \theta_S(x), \quad \varphi_S^j(x) = \theta_S(x) \xi_S^j(x), \quad j = 1, \dots, N,$$

where  $\xi_S^j(x)$  in the  $j$ -th component of the orientation  $\vec{S}(x)$  relative to the usual orthonormal basis  $e_{i_1} \wedge \dots \wedge e_{i_n}$ ,  $1 \leq i_1 < i_2 < \dots < i_n \leq n+k$  for  $\Lambda_n(\mathbb{R}^{n+k})$  (ordered in any convenient manner). Evidently, for any  $x \in \mathbb{R}^{n+k}$ ,

$$\varphi_S(x + \lambda y) = \varphi_{\eta_{x, \lambda\#} S}(y), \quad y \in \mathbb{R}^{n+k},$$

and, given a sequence  $\{S_i\} \subset \mathcal{D}_n(I + \mathbb{R}^{n+k})$  of such integer multiplicity currents, we trivially have

$$\varphi_{S_i}^j d\mathcal{H}^n \rightarrow \varphi_S^j d\mathcal{H}^n \quad \forall j \in \{1, \dots, N\} \iff S_i \rightharpoonup S$$

and

$$\varphi_{S_i}^0 d\mathcal{H}^n \rightarrow \varphi_S^0 d\mathcal{H}^n \iff \mu_{S_i} \rightarrow \mu_S$$

(where  $\psi_i d\mathcal{H}^n \rightarrow \psi d\mathcal{H}^n$  means  $\int f \psi_i d\mathcal{H}^n \rightarrow \int f \psi d\mathcal{H}^n \ \forall f \in C_c(\mathbb{R}^{n+k})$ ).

We shall also need the following simple lemma, the proof of which is left to the reader.

**3.5 Lemma.** *Suppose  $S$  is minimizing in  $\mathbb{R}^{n+k}$ ,  $\partial S = 0$ , and*

$$\eta_{x, 1\#} S = S \quad \forall x \in \mathbb{R}^m \times \{0\} \subset \mathbb{R}^{n+k}$$

*for some positive integer  $m < n$ . (Recall  $\eta_{x, 1} : y \mapsto y - x$ ,  $y \in \mathbb{R}^{n+k}$ .) Then*

$$S = [\mathbb{R}^m] \times S_0,$$

*where  $\partial S_0 = 0$  and  $S_0$  is minimizing in  $\mathbb{R}^{n+k-m}$ .*

*Furthermore if  $S$  is a cone (i.e.  $\eta_{0, \lambda\#} S = S$  for each  $\lambda > 0$ ), then so is  $S_0$ .*

**Proof of 3.3(1):** For each positive integer  $m$  and  $\beta \in (0, \frac{1}{2})$  let

$$(1) \quad U_{m, \beta} = \{x \in U : \Theta^n(\mu_{T, x}) < m - \beta\}.$$

Now  $T$  is minimizing in  $U \cap N$ , so by the monotonicity of 4.3 of Ch. 4 (which can be applied by virtue of 1.2) we have, firstly, that  $U_{m,\beta}$  is open, and secondly that for each  $x \in U_{m,\beta}$ , there is some ball  $B_{2\rho}(x) \subset U_{m,\beta}$  such that

$$(2) \quad \frac{\mu_T(B_\sigma(y))}{\omega_n \sigma^n} \leq m - \frac{\beta}{2} \quad \forall \sigma < \rho, \quad y \in B_\rho(x).$$

We ultimately want to prove

$$(3) \quad \mathcal{H}^{n-3+\alpha}(\cup_{m=1}^\infty \{x \in U_{m,\beta} : m - 1 + \beta < \Theta^n(\mu_T, x) < m - \beta\}) = 0$$

for each sufficiently small  $\alpha, \beta > 0$  and, in view of (2), by a rescaling and translation it will evidently suffice to assume

$$(4) \quad B_2(0) = U, \quad \frac{\mu_T(B_\sigma(y))}{\omega_n \sigma^n} \leq m - \beta \quad \forall \sigma < 1, \quad y \in B_1(0),$$

and then prove

$$(5) \quad \mathcal{H}^{n-3+\alpha} \{x \in B_1(0) : \Theta^n(\mu_T, x) \geq m - 1 + \beta\} = 0.$$

We consider the set  $\mathcal{T}$  of weak limit points of sequences  $S_i = \eta_{x_i, \lambda_i \#} T$  where  $|x_i| < 1 - \lambda_i$ ,  $0 < \lambda_i < 1$ , with  $\lim x_i \in B_1(0)$  and  $\lim \lambda_i = \lambda \geq 0$  both existing. For any such sequence  $S_i$  we have (by (4))

$$(6) \quad \limsup \mathbb{M}_W(S_i) < \infty$$

for each  $W \subset \subset \eta_{x,\lambda}(U)$  in case  $\lambda > 0$ , and for each  $W \subset \subset \mathbb{R}^{n+k}$  in case  $\lambda = 0$ . Hence we can apply the Compactness 2.4 to conclude that each element  $S$  of  $\mathcal{T}$  is integer multiplicity and

$$(7) \quad S \text{ minimizes in } \eta_{x,\lambda}U \cap \eta_{x,\lambda}N \text{ in case } S = \lim \eta_{x_i, \lambda_i \#} T$$

with  $\lim x_i = x$  and  $\lim \lambda_i = \lambda > 0$ , and

$$(8) \quad S \text{ minimizes in all of } \mathbb{R}^{n+k} \text{ in case } S = \lim \eta_{x_i, \lambda_i \#} T$$

with  $\lim x_i = x$  and  $\lim \lambda_i = 0$ . (Cf. the discussion in the proof of 3.1(2).)

For convenience we define

$$(9) \quad U_S = \begin{cases} \eta_{x,\lambda}U & \text{in case } \lim \lambda_i > 0 \text{ (as in (7))} \\ \mathbb{R}^{n+k} & \text{in case } \lim \lambda_i = 0 \text{ (as in (8))} \end{cases}$$

so that  $S \in \mathcal{D}_n(U_S)$  for each  $S \in \mathcal{T}$ .

Now by definition one readily checks that

$$(10) \quad \eta_{x, \lambda\#} \mathcal{T} = \mathcal{T}, \quad 0 < \lambda < 1, \quad |x| < 1 - \lambda,$$

and, by (4),

$$(11) \quad \Theta^n(\mu_S, y) \leq m - \beta \quad \forall y \in U_S, \quad S \in \mathcal{T}.$$

Furthermore by using the compactness theorem 2.4 together with the monotonicity 4.3 of Ch. 4, one readily checks that if  $S_i \rightarrow S$  ( $S_i, S \in \mathcal{T}$ ) and if  $y, y_i \in B_1(0)$  with  $\lim y_i = y$ , then

$$(12) \quad \Theta^n(\mu_S, y) \geq \limsup \Theta^n(\mu_{S_i}, y_i).$$

It now follows from (10), (11), (12) and 2.4 that all the hypotheses of Theorem A.4 (of Appendix A) are satisfied with

$$(13) \quad \mathcal{F} = \{\varphi_S : S \in \mathcal{T}\} \quad (\text{using notation of Remark 3.4})$$

and with  $\text{sing}$  defined by

$$(14) \quad \text{sing } \varphi_S = \{x \in U_S : \Theta^n(\mu_S, \cdot) \geq m - 1 + \beta\}$$

for  $S \in \mathcal{T}$ . We claim that in this case the additional hypothesis is satisfied with  $d = n - 3$ . Indeed suppose  $d \geq n - 2$ ; then there is  $S \in \mathcal{T}$  and  $\eta_{y, \lambda\#} S = S \quad \forall y \in L, \lambda > 0$  with  $L$  an  $(n - 2)$ -dimensional subspace of  $\mathbb{R}^{n+k}$ ,  $L \subset \text{sing } \varphi_S$ . Since we can make a rotation of  $\mathbb{R}^{n+k}$  to bring  $L$  into coincidence with  $\mathbb{R}^{n-2} \times \{0\}$ , we assume that  $L = \mathbb{R}^{n-2} \times \{0\}$ . Then by 3.5 we have

$$(15) \quad S = \llbracket \mathbb{R}^{n-2} \rrbracket \times S_0,$$

where  $S_0 \in \mathcal{D}_2(\mathbb{R}^N)$ ,  $N = 2 + k$ , with  $S_0$  a 2-dimensional area minimizing cone in  $\mathbb{R}^N$ . Then  $\text{spt } S_0$  is contained in a finite union  $\cup_{i=1}^q P_i$  of 2-planes, with  $P_i \cap P_j = \{0\} \quad \forall i \neq j$ . (For a formal proof of this characterization of 2 dimensional area minimizing cones, see for example [Whi82].) In particular, since  $\Theta^n(\mu_S, \cdot)$  is constant on  $P_i \setminus \{0\}$  (by the Constancy 2.41 of Ch. 6), we have that  $\Theta^n(\mu_S, y) \in \mathbb{Z}$  for every  $y \in \mathbb{R}^{n+k}$ , and by (11) it follows that  $\Theta^n(\mu_S, y) \leq m - 1 \quad \forall y \in \mathbb{R}^{n+k}$ . That is,  $\text{sing } \varphi_S = \emptyset$ , a contradiction, hence we can take  $d = n - 3$  as claimed. We have thus established (5) as required.  $\square$

**Proof of 3.3(2):** The proof goes similarly to 3.3(1). This time we assume (again without loss of generality) that

$$(1) \quad U = B_2(0),$$

and we prove that  $T$  has a tangent plane at all points of  $\text{spt } T \cap B_1(0)$  except for a set  $F \subset \text{spt } T \cap B_1(0)$  with

$$(2) \quad \mathcal{H}^{n-2+\alpha}(F) = 0 \quad \forall \alpha > 0.$$

$\mathcal{T}$  is as described in the proof of 3.3(1), and for any  $S \in \mathcal{T}$  and  $\beta > 0$  we let

$$R_\beta(S) = \{x \in \text{spt } S : B_\rho(x) \subset U_S \text{ and } h(\text{spt } S, L, \rho, x) < \beta\rho \\ \text{for some } \rho > 0 \text{ and some } n\text{-dimensional subspace } L \text{ of } \mathbb{R}^{n+k}\},$$

where  $U_S$  is as in the proof of 3.3(1) (so that  $S \in \mathcal{D}_n(U_S)$ ), and where we define

$$(3) \quad h(\text{spt } S, L, \rho, x) = \sup_{y \in \text{spt } S \cap B_\rho(x)} |q(y - x)|,$$

with  $q$  the orthogonal projection of  $\mathbb{R}^{n+k}$  onto  $L^\perp$ .

Now notice that (Cf. the proof of 3.3(1))

$$(4) \quad \eta_{x,\lambda\#}\mathcal{T} = \mathcal{T} \quad \forall 0 < \lambda < 1, |x| < 1 - \lambda,$$

and

$$(5) \quad \eta_{x,\lambda} R_\beta(S) = R_\beta(\eta_{x,\lambda\#}S), \quad S \in \mathcal{T}.$$

Furthermore if  $S_j \rightarrow S$ ,  $S_j, S \in \mathcal{T}$ , then by the monotonicity 4.3 of Ch. 4 it is quite easy to check that if  $y \in R_\beta(S)$  and if  $y_j \in \text{spt } S_j$  with  $y_j \rightarrow y$ , then  $y_j \in R_\beta(S_j)$  for all sufficiently large  $j$ . Because of this, and because of (4), (5) above, it is now straightforward to check that the hypotheses of A.4 hold with (again in notation of 3.4)

$$(6) \quad \mathcal{F} = \{\varphi_S : S \in \mathcal{T}\}$$

and

$$(7) \quad \text{sing } \varphi_S = \text{spt } \Theta^n(\mu_S, \cdot) \cap U_S \setminus R_\beta(S).$$

(Notice that  $R_\beta(S)$  is completely determined by  $\Theta^n(\mu_S, \cdot)$ , and hence this makes sense.) In this case we claim that  $d \leq n - 2$ . Indeed if  $d > n - 2$  (i.e.  $d = n - 1$ ) then  $\exists S \in \mathcal{T}$  such that

$$(8) \quad \eta_{x,\lambda\#}S = S \quad \forall x \in L, \lambda > 0, \text{ and } L \subset \text{sing } \varphi_S$$

where  $L$  is an  $(n - 1)$ -dimensional subspace. Then, supposing with loss of generality that  $L = \mathbb{R}^{n-1} \times \{0\}$ , we have by 3.5 that

$$(9) \quad S = \llbracket \mathbb{R}^{n-1} \rrbracket \times S_0,$$

where  $S_0$  is a 1-dimensional minimizing cone in  $\mathbb{R}^{k+1}$ . However it is easy to check that such a 1-dimensional minimizing cone necessarily has the form

$$(10) \quad S_0 = m\llbracket \ell \rrbracket,$$

where  $m \in \mathbb{Z}$  and  $\ell$  is a 1-dimensional subspace of  $\mathbb{R}^{k+1}$ . Thus (9) gives that  $S = m\llbracket L \rrbracket$  where  $L$  is an  $n$ -dimensional subspace and hence  $\text{sing } \varphi_S = \emptyset$ , a contradiction, so  $d \leq n - 2$  as claimed.

We therefore conclude from A.4 that for each  $S \in \mathcal{T}$

$$(11) \quad \mathcal{H}^{n-2+\alpha}(\text{spt } S \setminus R_\beta(S) \cap B_1(0)) = 0 \quad \forall \alpha > 0.$$

If  $\beta_j \downarrow 0$  we thus conclude in particular that

$$(12) \quad \mathcal{H}^{n-2+\alpha}(\text{spt } T \setminus \bigcup_{j=1}^{\infty} R_{\beta_j}(T) \cap B_1(0)) = 0 \quad \forall \alpha > 0.$$

However by (1) we see that

$$(13) \quad x \in \bigcup_{j=1}^{\infty} R_{\beta_j}(T) \iff T \text{ has a tangent plane at } x,$$

and therefore (12) gives (2) as required.  $\square$

## 4 Some Regularity Results (Arbitrary Codimension)

In this section, for  $T \in \mathcal{D}_n(U)$  any integer multiplicity current, we define a relatively closed subset  $\text{sing } T$  of  $U$  by

$$4.1 \quad \text{sing } T = \text{spt } T \setminus \text{reg } T,$$

where  $\text{reg } T$  denotes the set of points  $\xi \in \text{spt } T$  such that for some  $\rho > 0$  there is a  $m \in \mathbb{Z} \setminus \{0\}$  and an  $n$ -dimensional oriented  $C^1$  embedded submanifold  $M$  of  $\mathbb{R}^{n+k}$  with  $T = m\llbracket M \rrbracket$  in  $B_\rho(\xi)$ .

F.J. Almgren [Alm84] has proved the very important theorem that

$$\mathcal{H}^{n-2+\alpha}(\text{sing } T) = 0 \quad \forall \alpha > 0$$

in case  $\text{spt } T \subset N$ ,  $\partial T = 0$  and  $T$  is minimizing in  $N$ , where  $N$  is a smooth  $(n+k)$ -dimensional embedded submanifold of  $\mathbb{R}^{n+L}$  (where  $L \geq k$ ). The proof is very non-trivial and requires development of a whole new range of results for minimizing currents. We here restrict ourselves to more elementary results.

Firstly, the following theorem is an immediate consequence of The Allard Theorem 5.2 of Ch. 5 and Lemma 1.2 of the present chapter.

**4.2 Theorem.** *Suppose  $T \in \mathcal{D}_n(U)$  is integer multiplicity and minimizing in  $U \cap N$  for some embedded  $C^2$   $(n+k)$ -dimensional submanifold  $N$  of  $\mathbb{R}^{n+k}$ ,  $(\bar{N} \setminus N) \cap U = \emptyset$ , and suppose  $\text{spt } T \subset U \cap N$ ,  $\partial T = 0$  (in  $U$ ). Then  $\text{reg } T$  is dense in  $\text{spt } T$ .*

(Note that by definition  $\text{reg } T$  is relatively open in  $\text{spt } T$ .)

The following is a useful fact; however its applicability is limited by the hypothesis that  $\Theta^n(\mu_T, y) = 1$ .

**4.3 Theorem.** *Suppose  $\{T_i\} \subset \mathcal{D}_n(U)$ ,  $T \in \mathcal{D}_n(U)$  are integer multiplicity currents with  $T_i$  minimizing in  $U \cap N_i$ ,  $T$  minimizing in  $U \cap N$ ,  $N, N_i$  embedded  $(n+k)$ -dimensional  $C^2$  submanifolds, and  $\text{spt } T_i \subset N_i$ ,  $\text{spt } T \subset N$ ,  $\partial T_i = \partial T = 0$  (in  $U$ ). Suppose also that  $N_i$  converges to  $N$  in the  $C^2$  sense in  $U$ ,  $T_j \rightarrow T$  in  $\mathcal{D}_n(U)$ , and suppose  $y \in N \cap U$  with  $\Theta^n(\mu_T, y) = 1$ ,  $y = \lim y_j$ , where  $y_j$  is a sequence such that  $y_j \in \text{spt } T_j \forall j$ . Then  $y \in \text{reg } T$  and  $y_j \in \text{reg } T_j$  for all sufficiently large  $j$ .*

**Proof:** By virtue of the monotonicity formula 4.3 of Ch. 4 (which is applicable by 1.2) it is easily checked that

$$\limsup \Theta^n(\mu_{T_j}, y_j) \leq \Theta^n(\mu_T, y) = 1,$$

hence (since  $\Theta^n(\mu_{T_j}, y_j) \geq 1$  by 4.5 of Ch. 4) we conclude that  $\Theta^n(\mu_{T_j}, y_j) \rightarrow \Theta^n(\mu_T, y) = 1$ . Hence by Allard's Theorem 5.2 of Ch. 5 we have  $y \in \text{reg } T$  and  $y_j \in \text{reg } T_j$  for all sufficiently large  $j$ . (1.2 justifies the use of 5.2 of Ch. 5.)

Next we have the following consequences of A.4 of Appendix A.

**4.4 Theorem.** *Suppose  $T$  is as in 4.2, and in addition suppose  $\xi \in \text{spt } T$  is such that  $\Theta^n(\mu_T, \xi) < 2$ . Then there is a  $\rho > 0$  such that*

$$\mathcal{H}^{n-2+\alpha}(\text{sing } T \cap B_\rho(\xi)) = 0 \quad \forall \alpha > 0.$$

**Proof:** Let  $\alpha = 2 - \Theta^n(\mu_T, \xi)$  and let  $B_\rho(\xi)$  be such that  $B_{2\rho}(\xi) \subset U$  and

$$(1) \quad (\omega_n \sigma^n)^{-1} \mu_T(B_\sigma(\xi)) < 2 - \alpha/2$$

$\forall \zeta \in \text{spt } T \cap B_\rho(\xi)$ ,  $0 < \sigma < \rho$ . (Notice that such  $\rho$  exists by virtue of the monotonicity 4.3 of Ch. 4, which can be applied by Lemma 1.2.) Assume without loss of generality that  $\xi = 0$ ,  $\rho = 1$  and  $U = B_2(0)$ , and define  $\mathcal{T}$  to be the set of weak limits  $S$  of sequences  $\{S_i\}$  of the form  $S_i = \eta_{x_i, \lambda_i} T$ ,  $|x_i| < 1 - \lambda_i$ ,  $0 < \lambda_i < 1$ , where  $\lim x_i$  and  $\lim \lambda_i = \lambda$  are assumed to exist. Notice that

$$(2) \quad \limsup \mathbb{M}_W(S_i) < \infty$$

for each  $W \subset\subset \eta_{x,\lambda}(U)$  in case  $\lambda > 0$  and for each  $W \subset\subset \mathbb{R}^{n+k}$  in case  $\lambda = 0$ . Hence by the Compactness 2.4 any such  $S$  is integer multiplicity in  $U_S$

$$(3) \quad (U_S = \eta_{x,\lambda}U \text{ in case } \lambda > 0, \ U_S = \mathbb{R}^{n+k} \text{ in case } \lambda = 0)$$

and (Cf. the proof of 3.3(2))

$$(4) \quad S \text{ minimizes in } \eta_{x,\lambda}U \cap \eta_{x,\lambda}N \text{ in case } \lambda > 0$$

$$(5) \quad S \text{ minimizes in } \mathbb{R}^{n+k} \text{ in case } \lambda = 0.$$

One readily checks that, by definition of  $\mathcal{T}$ ,

$$(6) \quad \eta_{y,\tau\#}\mathcal{T} = \mathcal{T}, \ 0 < \tau < 1, \ |y| < 1 - \tau$$

Furthermore we note that (by (1))

$$(7) \quad \Theta^n(\mu_S, x) = 1, \ \mu_S\text{-a.e. } x \in U_S,$$

and by Allard's 5.2 of Ch. 5 there is  $\delta > 0$  such that

$$(8) \quad \text{sing } S = \{x \in U_S : \Theta^n(\mu_S, x) \geq 1 + \delta\}, \ S \in \mathcal{T}.$$

Now in view of (4), (5), (6), (7), (8) and the upper semi-continuity of  $\Theta^n$  as in (12) in the proof of 3.3(1), all the hypotheses of A.4 of A are satisfied with  $\mathcal{F} = \{\varphi_S : S \in \mathcal{T}\}$  (notation as in 3.4) and with  $\text{sing } \varphi_S = \{x \in U_S : \Theta^n(\mu_S, x) \geq 1 + \delta\}$  ( $\equiv \text{sing } S$  by (8)). In fact we claim that in this case we may take  $d = n - 2$ , because if  $d = n - 1$   $\exists S \in \mathcal{T}$  and  $\eta_{x,\lambda\#}S = S \ \forall x \in L, \lambda > 0$ , where  $L \subset \text{sing } S$  is an  $(n - 1)$ -dimensional subspace of  $\mathbb{R}^{n+k}$ , then (Cf. the last part of the proof of 3.3(2)) we have  $S = m\llbracket Q \rrbracket$  for some  $n$ -dimensional subspace  $Q$ . Hence  $\text{sing } S = \emptyset$ , a contradiction.  $\square$

The following theorem is often useful:

**4.5 Theorem.** *Suppose  $C \in \mathcal{D}_n(\mathbb{R}^{n+k})$  is minimizing in  $\mathbb{R}^{n+k}$ ,  $\partial C = 0$ , and  $C$  is a cone:  $\eta_{0,\lambda\#}C = C \ \forall \lambda > 0$ . Suppose further that  $\text{spt } C \subset \bar{H}$  where  $H$  is an open  $\frac{1}{2}$ -space of  $\mathbb{R}^{n+k}$  with  $0 \in \partial H$ . Then  $\text{spt } C \subset \partial H$ .*

**4.6 Remark:** The reader will see that the theorem here is actually valid with any stationary rectifiable varifold  $V$  in  $\mathbb{R}^{n+k}$  satisfying  $\eta_{0,\lambda\#}V = V$  in place of  $C$ .

**Proof of 4.5:** Since the varifold  $V$  associated with  $C$  is stationary (by 1.2) in  $\mathbb{R}^{n+k}$  we have 6.1 of Ch. 4 (Since  $(Dr)^\perp = 0$  by virtue of the fact that  $C$  is a cone),

$$(1) \quad \frac{d}{d\rho}(\rho^{-n} \int_{\mathbb{R}^{n+k}} h\varphi(r/\rho) d\mu_C) = \rho^{-n-1} \int_{\mathbb{R}^{n+k}} x \cdot (\nabla^C h)\varphi(r/\rho) d\mu_C$$

for each  $\rho > 0$ , where  $r = |x|$  and  $\varphi$  is a non-negative  $C^1$  function on  $\mathbb{R}$  with compact support, and  $h$  is an arbitrary  $C^1(\mathbb{R}^{n+k})$  function. ( $\nabla^C h(x)$  denotes the orthogonal projection of  $\nabla_{\mathbb{R}^{n+k}} h(x)$  onto the tangent space  $T_x V$  of  $V$  at  $x$ .)

Now suppose without loss of generality that  $H = \{x = (x^1, \dots, x^{n+k}) : x^1 > 0\}$  and select  $h(x) \equiv x^1$ . Then  $x \cdot \nabla^C h = e_1^T \cdot x = e_1 \cdot x^T = r e_1 \cdot \nabla^C r$ , where  $v^T$  denotes orthogonal projection of  $v$  onto  $T_x V$ . Thus the term on the right side of (1) can be written  $-\int_{\mathbb{R}^{n+k}} (e_1 \cdot \nabla^C r)(r\varphi(r/\rho)) d\mu_C$ , which in turn can be written  $-\int_{\mathbb{R}^{n+k}} e_1 \cdot \nabla^C \psi_\rho d\mu_C$ , where  $\psi_\rho(x) = \int_{|x|}^\infty r\varphi(r/\rho) dr$ . (Thus  $\psi_\rho$  has compact support in  $\mathbb{R}^{n+k}$ .) But  $e_1 \cdot \nabla^C \psi_\rho \equiv \operatorname{div}_V(\psi_\rho e_1)$ , and hence the term on the right of (1) actually vanishes by virtue of the fact that  $V$  is stationary. Thus (1) gives

$$\rho^{-n} \int_{\mathbb{R}^{n+k}} x_1 \varphi(r/\rho) d\mu_C = \text{const.}, \quad 0 < \rho < \infty.$$

In view of the arbitrariness of  $\varphi$ , this implies

$$\rho^{-n} \int_{B_\rho(0)} x_1 d\mu_C \equiv \text{const.}$$

However trivially we have  $\lim_{\rho \downarrow 0} \rho^{-n} \int_{B_\rho(0)} x_1 d\mu_C = 0$ , and hence we deduce

$$\rho^{-n} \int_{B_\rho(0)} x_1 d\mu_C = 0 \quad \forall \rho > 0.$$

Thus since  $x_1 \geq 0$  on  $\operatorname{spt} C (\subset \overline{H})$ , we conclude  $\operatorname{spt} C \subset \partial H (= \{x : x^1 = 0\})$ .  $\square$

The following corollary of 4.5 follows directly by combining 4.5 and 3.1(2).

**4.7 Corollary.** *If  $T$  is as in 4.2, if  $\xi \in \operatorname{spt} T$ , if  $Q$  is a  $C^1$  hypersurface in  $\mathbb{R}^{n+k}$  such that  $\xi \in Q$  and if  $\operatorname{spt} T$  is locally on one side of  $Q$  near  $\xi$ , then all tangent cones  $C$  of  $T$  at  $\xi$  satisfy  $\operatorname{spt} C \subset T_\xi Q \cap T_\xi N$ .*

## 5 Codimension 1 Theory

We begin by looking at those integer multiplicity currents  $T \in \mathcal{D}_n(U)$  with  $\operatorname{spt} T \subset N \cap U$ ,  $N$  an  $(n+1)$ -dimensional oriented embedded submanifold of  $\mathbb{R}^{n+k}$  with  $(\overline{N} \setminus N) \cap U = \emptyset$  and such that

$$5.1 \quad T = \partial[[E]]$$

(in  $U$ ), where  $E$  is an  $\mathcal{H}^{n+1}$ -measurable subset of  $N$ . (We know by 3.20 of Ch. 6, 1.4 that all minimizing currents  $T \in \mathcal{D}_n(U)$  with  $\partial T = 0$  and  $\operatorname{spt} T$  in  $N$  can be locally decomposed into minimizing currents of this special form.)

**5.2 Remark:** The fact that  $T$  has the form 5.1 and  $T$  is integer multiplicity evidently is equivalent to the requirement that if  $V \subset U$  is open, and if  $\varphi$  is a  $C^2$  diffeomorphism of  $V$  onto an open subset of  $\mathbb{R}^{n+k}$  such that  $\varphi(V \cap N) = G$ ,  $G$  open in  $\mathbb{R}^{n+1}$ , then  $\varphi(E)$  has locally finite perimeter in  $G$ . This is an easy consequence of 2.42 of Ch. 6, and in fact we see from this and 4.4 of Ch. 3 that any  $T$  of the form 5.1 with  $\mathbb{M}_W(T) < \infty \forall W \subset\subset U$  is automatically integer multiplicity with

$$(\ddagger) \quad \Theta^n(T, x) = 1, \mu_T\text{-a.e. } x \in U.$$

We shall here develop the theory of minimizing currents of the form 5.1; indeed we show this is naturally done using only the more elementary facts about currents. In particular we shall not in this section have any need of the Compactness 3.15 of Ch. 6 (instead we use only the elementary BV Compactness Theorem 2.6 of Ch. 2), nor shall we need the Deformation Theorem and the subsequent material of Chapter 6.

The following theorem could be derived from the general Compactness 2.4, but here (as we mentioned above) we can give a more elementary treatment. In this theorem, and subsequently, we take  $U \subset \mathbb{R}^{n+k}$  to be open and  $\mathcal{O}$  will denote the collection of  $(n+1)$ -dimensional oriented embedded  $C^2$  submanifolds  $N$  of  $\mathbb{R}^{n+k}$  with  $(\bar{N} \setminus N) \cap U = \emptyset$ ,  $N \cap U \neq \emptyset$ . A sequence  $\{N_j\} \subset \mathcal{O}$  is said to converge to  $N \in \mathcal{O}$  in the  $C^2$  sense in  $U$  if there are orientation preserving  $C^2$  embeddings  $\psi_j : N \cap U \rightarrow N_j$  with  $\psi_j \rightarrow \mathbf{1}_{N \cap U}$  then  $\eta_{x,\lambda} N$  converges to  $T_x N$  in the  $C^2$  sense in  $W$  as  $\lambda \downarrow 0$ , for each  $W \subset\subset \mathbb{R}^{n+k}$ .

In the following theorem  $p$  is a proper  $C^2$  map  $U \rightarrow N \cap U$  such that in some neighborhood  $V \subset U$  of  $N \cap U$ ,  $p$  coincides with the nearest point projection of  $V$  onto  $N$ . (Since the nearest point projection is  $C^2$  in some neighborhood of  $N \cap U$  it is clear that such  $p$  exists.)

**5.3 (Compactness Theorem for minimizing  $T$  as in 5.1).** *Suppose  $T_j \in \mathcal{D}_n(U)$ ,  $T_j = \partial[E_j]$  (in  $U$ ),  $E_j$   $\mathcal{H}^{n+1}$ -measurable subsets of  $N_j \cap U$ ,  $N_j \in \mathcal{O}$ ,  $N_j \rightarrow N \in \mathcal{O}$  in the  $C^2$  sense described above, and suppose  $T_j$  is integer multiplicity and minimizing in  $U \cap N_j$ .*

*Then there is a subsequence  $\{T_{j'}\}$  with  $T_{j'} \rightarrow T$  in  $\mathcal{D}_n(U)$ ,  $T$  integer multiplicity,  $T = \partial[E]$  (in  $U$ ),  $\chi_{p(E_{j'})} \rightarrow \chi_E$  in  $L^1_{\text{loc}}(\mathcal{H}^{n+1}, U)$ ,  $\mu_{T_{j'}} \rightarrow \mu_T$  (in the usual sense of Radon measures) in  $U$ , and  $T$  is minimizing in  $N \cap U$ .*

**5.4 Remarks:** (1) Recall (from 5.2) that the hypothesis that  $T_j$  is integer multiplicity is automatic if we assume merely that  $\mathbb{M}_W(T_j) < \infty \forall W \subset\subset U$ .

(2) We make no *a-priori* assumptions on local boundedness of the mass of  $T_j$  (we see in the proof that this is automatic for minimizing currents as in 5.1).

(3) Let  $h(x, t) = x + t(p(x) - x)$ ,  $x \in U$ ,  $0 \leq t \leq 1$ . Using the homotopy formula 2.32 of Ch. 6 (and in particular the inequality 2.34 of Ch. 6) together with the fact that

$N_j \rightarrow N$  in the  $C^2$  sense in  $U$ , it is straightforward to check that

$$T_j - T = \partial R_j, \quad R_j = h_{\#}(\llbracket(0, 1)\rrbracket \times T_j) + p_{\#}(\llbracket E_j \rrbracket - \llbracket E \rrbracket)$$

with

$$\mathbb{M}_W(R_j) \rightarrow 0 \quad \forall W \subset\subset U,$$

provided that  $\chi_{p(E_j)} \rightarrow \chi_E$  as claimed in the theorem. Thus once we establish  $\chi_{p(E_j)} \rightarrow \chi_E$  for some  $E$ , then we can use the argument of 2.4 (with  $S_j = 0$ ) in order to conclude

(i)  $T$  is minimizing in  $U$

(ii)  $\mu_{T_j} \rightarrow \mu_T$  in  $U$ .

(Notice we have not had to use the deformation theorem here.) In the following proof we therefore concentrate on proving  $\chi_{p(E_j)} \rightarrow \chi_E$  in  $L^1_{\text{loc}}(\mathcal{H}^{n+1}, N \cap U)$  for some subsequence  $\{j'\}$  and some  $E$  such that  $\partial \llbracket E \rrbracket$  has locally finite mass in  $U$ . ( $T$  is then automatically integer multiplicity by 5.2.)

**Proof of 5.3:** We first establish a local mass bound for the  $T_j$  in  $U$ : if  $\xi \in N$  and  $B_{\rho_0}(\xi) \subset U$ , then

$$(1) \quad \mathbb{M}(T_j \llcorner B_{\rho}(\xi)) \leq \frac{1}{2} \mathcal{H}^n(\partial B_{\rho}(\xi) \cap N), \quad \mathcal{L}^1\text{-a.e. } \rho \in (0, \rho_0).$$

This is proved by simple area comparison as follows:

With  $r(x) = |x - \xi|$ , by the elementary slicing theory of 4.5(1),(2) of Ch. 6 we have that, for  $\mathcal{L}^1$ -a.e.  $\rho \in (0, \rho_0)$ , the slice  $\langle \llbracket E_j \rrbracket, r, \rho \rangle$  (i.e. the slice of  $\llbracket E_j \rrbracket$  by  $\partial B_{\rho}(\xi)$ ) is integer multiplicity, and (using  $T_j = \partial \llbracket E_j \rrbracket$ ),

$$\partial \llbracket E_j \cap B_{\rho}(\xi) \rrbracket = T_j \llcorner B_{\rho}(\xi) + \langle \llbracket E_j \rrbracket, r, \rho \rangle.$$

Hence (applying  $\partial$  to this identity)

$$\partial(T_j \llcorner B_{\rho}(\xi)) = -\partial \langle \llbracket E_j \rrbracket, r, \rho \rangle, \quad \mathcal{L}^1\text{-a.e. } \rho \in (0, \rho_0),$$

and by Definition 1.1 of minimizing

$$\mathbb{M}(T_j \llcorner B_{\rho}(\xi)) \leq \mathbb{M} \langle \llbracket E_j \rrbracket, r, \rho \rangle.$$

Since  $-\tilde{T}_j$  is also minimizing in  $N \cap U$  we then have

$$(2) \quad \mathbb{M}(T_j \llcorner B_{\rho}(\xi)) \leq \min \left\{ \mathbb{M} \langle \llbracket E_j \rrbracket, r, \rho \rangle, \mathbb{M} \langle \llbracket \tilde{E}_j \rrbracket, r, \rho \rangle \right\}$$

for  $\mathcal{L}^1$ -a.e.  $\rho \in (0, \rho_0)$ , where  $\tilde{E}_j = N \cap U \setminus E_j$ .

Now of course  $[\![\tilde{E}]\!] + [\![E_j]\!] = [\![N \cap U]\!]$ , so that (for a.e.  $\rho \in (0, \rho_0)$ )

$$\langle [\![E_j]\!], r, \rho \rangle + \langle [\![\tilde{E}_j]\!], r, \rho \rangle = \langle N, r, \rho \rangle$$

and hence (2) gives (1) as required (because  $\mathbb{M}(\langle N, r, \rho \rangle) \leq \mathcal{H}^n(N \cap \partial B_\rho(\xi))$  by virtue of the fact that  $|Dr| = 1$ , hence  $|\nabla^N r| \leq 1$ ).

Now by virtue of (1) and 5.2 we deduce from the BV Compactness 2.6 of Ch.2 that some subsequence  $\{\chi_{p(E_j')}\}$  of  $\{\chi_{p(E_j)}\}$  converges in  $L^1_{\text{loc}}(\mathcal{H}^{n+1}, N \cap U)$  to  $\chi_E$ , where  $E \subset N$  is  $\mathcal{H}^{n+1}$ -measurable and such that  $\partial[\![E]\!]$  is integer multiplicity (in  $U$ ). The remainder of the theorem now follows as described in 5.4(3).  $\square$

**5.5 (Existence of tangent cones).** *Suppose  $T = \partial[\![E]\!] \in \mathcal{D}_n(U)$  is integer multiplicity, with  $E \subset N \cap U$ ,  $N \in \mathcal{O}$ , and  $T$  is minimizing in  $U \cap N$ . Then for each  $x \in \text{spt } T$  and each sequence  $\{\lambda_j\} \downarrow 0$  there is a subsequence  $\{\lambda_{j'}\}$  and an integer multiplicity  $C \in \mathcal{D}_n(\mathbb{R}^{n+k})$  with  $C$  minimizing in  $\mathbb{R}^{n+k}$ ,  $0 \in \text{spt } C \subset T_x N$ ,  $\Theta^n(\mu_C, 0) = \Theta^n(\mu_T, x)$ ,  $C = \partial[\![F]\!]$ ,  $F$  an  $\mathcal{H}^{n+1}$ -measurable subset of  $T_x N$ ,*

$$(1) \quad \mu_{\eta_{x, \lambda_{j'} \#} T} \rightarrow \mu_C \text{ in } \mathbb{R}^{n+k}, \quad \chi_{p(\eta_{x, \lambda_{j'}}(E))} \rightarrow \chi_F \text{ in } L^1_{\text{loc}}(\mathcal{H}^{n+1}, T_x N),$$

where  $p$  is the orthogonal projection of  $\mathbb{R}^{n+k}$  onto  $T_x N$ , and

$$(2) \quad \eta_{0, \lambda \#} C = C, \quad \eta_{0, \lambda} F = F \quad \forall \lambda > 0.$$

**5.6 Remark:** The proof given here is independent of the general tangent cone Existence 3.1.

**Proof of 5.5:** As we remarked prior to 5.3,  $\eta_{x, \lambda_j} N$  converges to  $T_x N$  in the  $C^2$  sense in  $W$  for each  $W \subset\subset \mathbb{R}^{n+k}$ . By the Compactness 5.3 we then have a subsequence  $\lambda_j$ , such that all the required conclusions, except possibly for 5.5(2) and the fact that  $0 \in \text{spt } C$ , hold. To check that  $0 \in \text{spt } C$  and that 5.5(2) is valid, we first note by 1.2 that the varifold  $V$  associated with  $T$  is stationary in  $N \cap U$  (and that  $V$  therefore has locally bounded generalized mean curvature  $\underline{H}$  in  $N \cap U$ ). Therefore by the monotonicity formula 4.3 of Ch. 4, and by 4.5 of Ch. 4, we have

$$\Theta^n(\mu_V, x) \text{ exists and is } \geq 1.$$

Since  $\mu_{\eta_{x, \lambda_j \#} T} \rightarrow \mu_C$ , we then have  $\Theta^n(\mu_C, 0) = \Theta^n(\mu_T, x) \geq 1$ , so  $0 \in \text{spt } C$ , and by Theorem ?? of Ch. 4 we deduce that the varifold  $V_C$  associated with  $C$  is a cone. Then in particular  $x \wedge \tilde{C}(x) = 0$  for  $\mu_C$ -a.e.  $x \in \mathbb{R}^{n+k}$  and hence, if we let  $h$  be the homotopy  $h(t, x) = tx + (1-t)\lambda x$ , we have  $h_\#([\![0, 1]\!] \times C) = 0$ , and then by the homotopy formula 2.32 of Ch. 6 (since  $\partial C = 0$ ) we have  $\eta_{0, \lambda \#} C = C$  as required. Finally

since  $\text{spt } C$  has locally finite  $\mathcal{H}^n$ -measure (indeed by 4.5 of Ch. 4  $\text{spt } C$  is the closed set  $\{y \in \mathbb{R}^{n+k} : \Theta^n(\mu_C, y) \geq 1\}$ ), we have

$$[F] = [\tilde{F}],$$

where  $\tilde{F}$  is the (open) set  $\{y \in T_x N \setminus \text{spt } C : \Theta^{n+1}(\mathcal{H}^{n+1}, T_x N, y) = 1\}$ . Evidently  $\eta_{0,\lambda}(\tilde{F}) = \tilde{F}$  (because  $\eta_{0,\lambda}(\text{spt } C) = \text{spt } C$ ). Hence the required result is established with  $\tilde{F}$  in place of  $F$ .  $\square$

**5.7 Corollary.**<sup>4</sup> *Suppose  $T$  is as in 5.5 and in addition suppose there is an  $n$ -dimensional embedded submanifold  $\Sigma$  in  $\mathbb{R}^{n+k}$  with  $x \in \Sigma \subset N \cap U$  for some  $x \in \text{spt } T$ , and suppose  $\text{spt } T \setminus \Sigma$  lies locally, near  $x$ , on one side of  $\Sigma$ . Then  $x \in \text{reg } T$ . (reg  $T$  is as in 4.1)*

**Proof:** Let  $C = \partial[F]$  ( $F \subset T_x N$ ) be any tangent cone for  $T$  at  $x$ . By assumption  $\text{spt}[F] \subset \bar{H}$ , where  $H$  is an open  $\frac{1}{2}$ -space in  $T_x N$  with  $0 \in \partial H$ . Then, by 4.5,  $\text{spt } C \subset \partial H$  and hence the Constancy 2.41 of Ch. 6 since  $C$  is integer multiplicity rectifiable, it follows that  $C = \pm \partial[H]$ . However  $\text{spt}[F] \subset \bar{H}$ , hence  $C = +\partial[H]$ . Then  $\Theta^n(\mu_C, y) \equiv 1$  for  $y \in \partial H$ , and in particular  $\Theta^n(\mu_C, 0) (= \Theta^n(\mu_T, x)) = 1$ , so that  $x \in \text{reg } T$  (by Allard's Theorem 5.2 of Ch. 5) as required.

We next want to prove the main regularity theorem for codimension 1 currents. We continue to define  $\text{sing } T$ ,  $\text{reg } T$  as in 4.1.

**5.8 Theorem.** *Suppose  $T = \partial[E] \in \mathcal{D}_n(U)$  is integer multiplicity, with  $E \subset N \cap U$ ,  $n \in \mathcal{O}$ , and  $T$  minimizing in  $N \cap U$ . Then  $\text{sing } T = \emptyset$  for  $n \leq 6$ ,  $\text{sing } T$  is locally finite in  $U$  for  $n = 7$ , and  $\mathcal{H}^{n-7+\alpha}(\text{sing } T) = 0 \ \forall \alpha > 0$  in case  $n > 7$ .*

**Proof:** We are going to use the abstract dimension reducing argument of Appendix A (Cf. the proof of 4.4).

To begin we note that it is enough (by re-scaling, translation, and restriction) to assume that

$$U = \check{B}_2(0)$$

and to prove that

$$\begin{cases} \text{sing } T \cap B_1(0) = \emptyset \text{ if } n \leq 6, \text{ sing } T \cap B_1(0) \text{ discrete if } n = 7, \\ \mathcal{H}^{n-7+\alpha}(\text{sing } T \cap B_1(0)) = 0 \ \forall \alpha > 0 \text{ if } n > 7. \end{cases}$$

Let  $\mathcal{T}$  be the set of currents as defined in the proof of 4.4<sup>5</sup>, and for each  $S \in \mathcal{T}$  let  $\varphi_S$  be the function  $:\mathbb{R}^{n+k} \rightarrow \mathbb{R}^{n+1}$  associated with  $S$  as in 3.4. Also, let

$$\mathcal{F} = \{\varphi_S : S \in \mathcal{T}\}$$

<sup>4</sup>Cf. Miranda [Mir67]

<sup>5</sup>We still have  $\Theta^n(\mu_S, x) = 1$  for  $\mu_S$ -a.e.  $x \in U_S$ , this time by 5.3 and 5.2 (§)

and define

$$\text{sing } \varphi_S = \text{sing } S.$$

(sing  $S$  as defined in 4.1.)

By A.4 we then have either  $\text{sing } S = \emptyset$  for all  $S \in \mathcal{T}$  (and hence  $\text{sing } T = \emptyset$ ) or

$$\dim B_1(0) \cap \text{sing } S \leq d,$$

where  $d \in [0, n-1]$  is the integer such that

$$\dim B_1(0) \cap \text{sing } S \leq d \text{ for all } S \in \mathcal{T}$$

and such that there is  $S \in \mathcal{T}$  and a  $d$ -dimensional subspace  $L$  of  $\mathbb{R}^{n+k}$  such that

$$\eta_{x, \lambda\#} S = S \quad \forall x \in L, \lambda > 0$$

and

$$5.9 \quad \text{sing } S = L.$$

Supposing without loss of generality that  $L = \mathbb{R}^d \times \{0\}$ , we then (by 3.5) have

$$5.10 \quad S = \llbracket R^d \rrbracket \times S_0$$

where  $\partial S_0 = 0$ ,  $S_0$  is minimizing in  $\mathbb{R}^{n+k-1}$ , and  $\text{sing } S_0 = \{0\}$ . (With  $S$  as in 5.10,  $\text{sing } S_0 = \{0\} \iff 5.9$ .) Also, by definition of  $\mathcal{T}$ ,  $\text{spt } S \subset$  some  $(n+1)$ -dimensional subspace of  $\mathbb{R}^{n+k}$ , hence without loss of generality we have that  $S_0$  is an  $(n-d)$ -dimensional minimizing cone in  $\mathbb{R}^{n-d+1}$  with  $\text{sing } S_0 = \{0\}$ . Then by the result of J. Simons (see B) we have  $n-d > 6$ ; i.e.  $d \leq n-7$ . Notice that this contradicts  $d \geq 0$  in case  $n < 7$ . Thus for  $n < 7$  we must have  $\text{sing } T = \emptyset$  as required. If  $n = 7$ ,  $\text{sing } T$  is discrete by the last part of A.4.

**5.11 Corollary.** *If  $T$  is as in 5.8, and if  $T_1 \in \mathcal{D}_n(U)$  is obtained by equipping a component of  $\text{reg } T$  with multiplicity 1 and with orientation of  $T$ , then  $\partial T_1 = 0$  (in  $U$ ) and  $T_1$  is minimizing in  $U \cap N$ .*

**5.12 Remark:** Notice that this means we can write

$$T = \sum_{j=1}^{\infty} T_j,$$

where each  $T_j$  is obtained by equipping a component  $M_j$  of  $\text{reg } T$  with multiplicity 1 and with the orientation of  $T$ ; then  $M_i \cap M_j = \emptyset \quad \forall i \neq j$ ,  $\partial T_j = 0$ , and  $T_j$  is minimizing in  $U \quad \forall j$ . Furthermore (since  $\mu_{T_j}(B_\rho(x)) \geq c\rho^n$  for  $B_\rho(x) \subset U$  and  $x \in \text{spt } T_j$  by

virtue of 1.2 and the monotonicity 4.3 of Ch. 4) only finitely many  $T_j$  can have support intersecting a given compact subset of  $U$ .

**Proof of 5.11:** The main point is to prove

$$(1) \quad \partial T_1 = 0 \text{ in } U.$$

The fact that  $T_1$  is minimizing in  $U$  will then follow from 1.4 and the fact that  $\mathbb{M}_W(T_1) + \mathbb{M}_W(T - T_1) = \mathbb{M}_W(T) \quad \forall W \subset\subset U$ .

To check (1) let  $\omega \in \mathcal{D}^{n-1}(U)$  be arbitrary and note that if  $\zeta \equiv 0$  in some neighborhood of  $\text{spt } T \setminus M_1$

$$(2) \quad T_1(d(\zeta\omega)) = T(d(\zeta\omega)) = \partial T(\zeta\omega) = 0.$$

Now corresponding to any  $\varepsilon > 0$  we construct  $\zeta$  as follows: since  $\mathcal{H}^{n-1}(\text{sing } T) = 0$  (by 5.8) and since  $\text{sing } T \cap \text{spt } \omega$  is compact, we can find a finite cover of  $\text{sing } T \cap \text{spt } \omega$  by balls  $\{B_{\rho_j}(\xi_j)\}_{j=1,\dots,P}$  with  $\xi_j \in \text{sing } T \cap \text{spt } \omega$  and  $\sum_{j=1}^P \rho_j^{n-1} < \varepsilon$ . For each  $j = 1, \dots, P$  let  $\varphi_j \in C_c^\infty(\mathbb{R}^{n+k})$  be such that  $\varphi_j \equiv 1$  on  $B_{\rho_j}(\xi_j)$ ,  $\varphi_j = 0$  on  $\mathbb{R}^{n+k} \setminus B_{2\rho_j}(\xi_j)$ , and  $0 \leq \varphi_j \leq 1$  everywhere, and  $|D\varphi_j| \leq 2/\rho_j$ . Now choose  $\zeta = \prod_{j=1}^P \varphi_j$  in a neighborhood of  $\text{spt } T_1$  and so that  $\zeta \equiv 0$  in a neighborhood of  $\text{spt } T \setminus \text{spt } T_1$ . Then  $d\zeta = \sum_{i=1}^P \prod_{j \neq i} \varphi_j d\varphi_i$  on  $\text{spt } T_1$ , and hence

$$|T(d(\zeta\omega) - \zeta d\omega)| \leq c|\omega| \sum_{j=1}^P \rho_j^{n-1} \leq c\varepsilon|\omega| \text{ on } \text{spt } T_1.$$

The letting  $\varepsilon \downarrow 0$  in (2), and noting that  $\zeta d\omega \rightarrow d\omega$   $\mathcal{H}^n$ -a.e. in  $\text{spt } T_1 \cap N \cap \text{spt } \omega$  (and using  $|\zeta| \leq 1$ ), we conclude  $T_1(d\omega) = 0$ . That is  $\partial T_1 = 0$  in  $U$  as required.  $\square$

Finally we have the following lemma.

**5.13 Lemma.** *If  $T_1 = \partial \llbracket E_1 \rrbracket$ ,  $T_2 = \partial \llbracket E_2 \rrbracket \in \mathcal{D}_n(U)$ ,  $U$  bounded,  $E_1, E_2 \subset U \cap N$ ,  $N$  of class  $C^4$ ,  $N \in \mathcal{O}$ ,  $T_1, T_2$  minimizing in  $U \cap N$ ,  $\text{reg } T_1, \text{reg } T_2$  are connected, and  $E_1 \cap V \subset E_2 \cap V$  for some neighborhood  $V$  of  $\partial U$ , then  $\text{spt} \llbracket E_1 \rrbracket \subset \text{spt} \llbracket E_2 \rrbracket$  and either  $\llbracket E_1 \rrbracket = \llbracket E_2 \rrbracket$  or  $\text{spt } T_1 \cap \text{spt } T_2 \subset \text{sing } T_1 \cap \text{sing } T_2$ .*

**Proof:** Since  $\mathcal{H}^{n+1}(\text{spt } T_j) = 0$  (in fact  $\text{spt } T_j$  has locally finite  $\mathcal{H}^n$ -measure in  $U$  by virtue of the fact that  $\Theta^n(\mu_{T_j}, x) \geq 1 \quad \forall x \in \text{spt } T_j$ ), we may assume that  $E_1$  and  $E_2$  are open with  $U \cap \partial E_j = U \cap \partial \bar{E}_j = \text{spt } T_j$ ,  $j = 1, 2$ .

Let  $S_1, S_2 \in \mathcal{D}_n(U)$  be the currents defined by

$$S_1 = \partial \llbracket E_1 \cap E_2 \rrbracket, \quad S_2 = \partial \llbracket E_1 \cup E_2 \rrbracket.$$

Using the hypothesis concerning  $V$  we have

$$(3) \quad S_j \llcorner (V \cap U) = T_j \llcorner (V \cap U), \quad j = 1, 2.$$

On the other hand we trivially have

$$[[E_1 \cap E_2]] + [[E_1 \cup E_2]] = [[E_1]] + [[E_2]],$$

so (applying  $\partial$ ) we get

$$(4) \quad S_1 + S_2 = T_1 + T_2.$$

Furthermore  $E_1 \cap E_2 \subset E_1 \cup E_2$ , so

$$(5) \quad \begin{aligned} \mathbb{M}_W(S_1) + \mathbb{M}(S_2) &= \mathbb{M}_W(S_1 + S_2) \\ &= \mathbb{M}_W(T_1 + T_2) \quad (\text{by (4)}) \\ &\leq \mathbb{M}_W(T_1) + \mathbb{M}_W(T_2) \end{aligned}$$

$\forall W \subset\subset U$ . On the other hand, choosing an open  $V_0$  so that  $\partial U \subset V_0 \subset\subset V$ , and using (3) together with the fact that  $T_1$  is minimizing, we have

$$\mathbb{M}_W(S_1) \geq \mathbb{M}_W(T_1), \quad W = U \setminus \bar{V}_0,$$

and hence (combining this with (5))

$$\mathbb{M}_W(S_2) \leq \mathbb{M}_W(T_2)$$

for  $W = U \setminus \bar{V}_0$ . Thus (using (3) with  $j = 2$ )  $S_2$  is minimizing in  $U$ . Likewise  $S_1$  is minimizing in  $U$ .

We next want to prove that either  $T_1 = T_2$  or  $\text{reg } T_1 \cap \text{reg } T_2 = \emptyset$ . Suppose  $\text{reg } T_1 \cap \text{reg } T_2 \neq \emptyset$ . If the tangent spaces of  $\text{reg } T_1$  and  $\text{reg } T_2$  coincide at every point of their intersection, then using suitable local coordinates  $(x, z) \in \mathbb{R}^n \times \mathbb{R}$  for  $N$  near a point  $\xi \in \text{reg } T_1 \cap \text{reg } T_2$ , we can write

$$\text{reg } T_j = \text{graph } u_j, \quad j = 1, 2,$$

where  $Du_1 = Du_2$  at each point where  $u_1 = u_2$ , and where both  $u_1, u_2$  are (weak)  $C^1$  solutions of the equation

$$\frac{\partial}{\partial x_i} \left( \frac{\partial F}{\partial p_i}(x, u, Du) \right) - \frac{\partial F}{\partial z}(x, u, Du) = 0,$$

where  $F = F(x, z, p)$ ,  $(x, z, p) \in \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^n$ , is the area functional for graphs  $z = u(x)$  relative to the local coordinates  $x, z$  for  $N$ . Since  $N$  is  $C^4$  we then deduce (from standard quasilinear elliptic theory—see e.g. [GT01]) that  $u_1, u_2$  are  $C^{3,\alpha}$ . Now the difference  $u_1 - u_2$  of the solutions evidently satisfies an equation of the general form

$$D_j(a_{ij} D_i u) + b_i D_i u + cu = 0,$$

where  $a_{ij}, b_i, c$  are  $C^{2,\alpha}$ . By standard unique continuation results (see e.g. [Pro60]) we then see that  $Du_1 = Du_2$  at each point where  $u_1 = u_2$  is impossible if  $u_1 - u_2$  changes sign. On the other hand the Harnack inequality ([GT01]) tells us that either  $u_1 \equiv u_2$  or  $|u_1 - u_2| > 0$  in case  $u_1 - u_2$  does not change sign. Thus we deduce that either  $T_1 = T_2$  or  $\text{reg } T_1 \cap \text{reg } T_2 = \emptyset$  or there is a point  $\xi \in \text{reg } T_1 \cap \text{reg } T_2$  such that  $\text{reg } T_1$  and  $\text{reg } T_2$  intersect *transversely* at  $\xi$ . But then we would have  $\mathcal{H}^{n-1}(\text{sing } \partial \llbracket E_1 \cap E_2 \rrbracket) > 0$ , which by virtue of 5.8 contradicts the fact (established above) that  $\partial \llbracket E_1 \cap E_2 \rrbracket$  is minimizing in  $U$ .

Thus either  $T_1 = T_2$  or  $\text{reg } T_1 \cap \text{reg } T_2 = \emptyset$ , and it follows in either case that  $E_1 \subset E_2$ . On the other hand we then have  $\text{sing } T_1 \cap \text{reg } T_2 = \emptyset$  and  $\text{sing } T_2 \cap \text{reg } T_1 = \emptyset$  by virtue of 5.7. Thus we conclude that  $E_1 \subset E_2$  and  $\text{spt } T_1 \cap \text{spt } T_2 \subset \text{sing } T_1 \cap \text{sing } T_2$  as required.

# Chapter 8

## Theory of General Varifolds

§ 1	Basics, First Rectifiability Theorem .....	235
§ 2	First Variation .....	238
§ 3	Monotonicity and Consequences .....	240
§ 4	Constancy Theorem .....	243
§ 5	Varifold Tangents and Rectifiability Theorem .....	245

### 1 Basics, First Rectifiability Theorem

We let  $G(n+k, n)$  denote the collection of all  $n$ -dimensional subspaces of  $\mathbb{R}^{n+k}$ , equipped with the metric  $\rho(S, T) = |p_S - p_T| = (\sum_{i,j=1}^{n+k} (p_S^{ij} - p_T^{ij})^2)^{\frac{1}{2}}$  where  $p_S, p_T$  denote the orthogonal projections of  $\mathbb{R}^{n+k}$  onto  $S, T$  respectively, and  $p_S^{ij} = e_i \cdot p_S(e_j)$ ,  $p_T^{ij} = e_i \cdot p_T(e_j)$  are the corresponding matrices with respect to the standard orthonormal basis  $e_1, \dots, e_{n+k}$  for  $\mathbb{R}^{n+k}$ .

For a subset  $A \subset \mathbb{R}^{n+k}$  we define

$$G_n(A) = A \times G(n+k, n),$$

equipped with the product metric. Of course then  $G_n(K)$  is compact for each compact  $K \subset \mathbb{R}^{n+k}$ .  $G_n(\mathbb{R}^{n+k})$  is locally homeomorphic to a Euclidean space of dimension  $n+k+nk$ .

By an  $n$ -varifold we mean simply any Radon measure  $V$  on  $G_n(\mathbb{R}^{n+k})$ . By an  $n$ -varifold on  $U$  ( $U$  open in  $\mathbb{R}^{n+k}$ ) we mean any Radon measure  $V$  on  $G_n(U)$ . Given such an  $n$ -varifold  $V$  on  $U$ , there corresponds a Radon measure  $\mu = \mu_V$  on  $U$  (called the *weight* of

$V$ ) defined by

$$\mu(A) = V(\pi^{-1}(A)), \quad A \subset U,$$

where, here and subsequently,  $\pi$  is the projection  $(x, S) \mapsto x$  of  $G_n(U)$  onto  $U$ . The mass  $\mathbb{M}(V)$  of  $V$  is defined by

$$\mathbb{M}(V) = \mu_V(U) \quad (= V(G_n(U))).$$

for any Borel subset  $A \subset U$  we use the usual terminology  $V \llcorner G_n(A)$  to denote the restriction of  $V$  to  $G_n(A)$ ; thus

$$(V \llcorner G_n(A))(B) = V(B \cap G_n(A)), \quad B \subset G_n(U).$$

Given an  $n$ -rectifiable varifold  $\underline{v}(M, \theta)$  on  $U$  (in the sense of Ch. 4) there is a corresponding  $n$ -varifold  $V$  (also denoted by  $\underline{v}(M, \theta)$ , or simply  $\underline{v}(M)$  in case  $\theta \equiv 1$  on  $M$ ), defined by

$$V(A) = \mu(\pi(TM \cap A)), \quad A \subset G_n(U),$$

where  $\mu = \mathcal{H}^n \llcorner \theta$  and  $TM = \{(x, T_x M) : x \in M_*\}$ , with  $M_*$  the set of  $x \in M$  such that  $M$  has an approximate tangent space  $T_x M$  with respect to  $\theta$  at  $x$  in the sense of 1.7 of Ch. 3. Evidently  $V$ , so defined, has weight measure  $\mu_V = \mathcal{H}^n \llcorner \theta = \mu$ .

The question of when a general  $n$ -varifold actually corresponds to an  $n$ -rectifiable varifold in this way is satisfactorily answered in the next theorem. Before stating this we need a definition:

**1.1 Definition:** Given  $T \in G(n+k, n)$ ,  $x \in U$ , and  $\theta \in (0, \infty)$ , we say that an  $n$ -varifold  $V$  on  $U$  has tangent space  $T$  with multiplicity  $\theta$  at  $x$  if

$$(\ddagger) \quad \lim_{\lambda \downarrow 0} V_{x,\lambda} = \theta \underline{v}(T),$$

where the limit is in the usual sense of Radon measures on  $G_n(\mathbb{R}^{n+k})$ . In 1.1  $(\ddagger)$  we use the notation that  $V_{x,\lambda}$  is the  $n$ -varifold defined by

$$V_{x,\lambda}(A) = \lambda^{-n} V(\{(\lambda y + x, S) : (y, S) \in A\} \cap G_n(U))$$

for  $A \subset G_n(\mathbb{R}^{n+k})$ .

**1.2 (First Rectifiability Theorem.)** Suppose  $V$  is an  $n$ -varifold on  $U$  which has a tangent space  $T_x$  with multiplicity  $\theta(x) \in (0, \infty)$  for  $\mu_V$ -a.e.  $x \in U$ . Then  $V$  is  $n$ -rectifiable; in fact  $M \equiv \{x \in \text{spt } V : T_x \text{ and } \theta(x) \text{ both exist}\}$  is  $\mathcal{H}^n$ -measurable, countably  $n$ -rectifiable,  $\theta$  is locally  $\mathcal{H}^n$ -integrable on  $M$ , and  $V = \underline{v}(M, \theta)$ .

In the proof of 1.2 (and also subsequently) we shall need the following technical lemma:

**1.3 Lemma.** *Let  $V$  be any  $n$ -varifold on  $U$ . Then for  $\mu_V$ -a.e.  $x \in U$  there is a Radon measure  $\eta_V^{(x)}$  on  $G(n+k, n)$  such that, for any continuous  $\beta$  on  $G(n+k, n)$ ,*

$$\int_{G(n+k, n)} \beta(S) d\eta_V^{(x)}(S) = \lim_{\rho \downarrow 0} \frac{\int_{G_n(B_\rho(x))} \beta(S) dV(y, S)}{\mu_V(B_\rho(x))}.$$

Furthermore for any Borel set  $A \subset U$ ,

$$\int_{G_n(A)} \beta(S) dV(x, S) = \int_A \int_{G(n+k, n)} \beta(S) d\eta_V^{(x)}(S) d\mu_V(x)$$

provided  $\beta \geq 0$ .

**Proof:** The proof is a simple consequence of the differentiation theory for Radon measures and the separability of  $\mathcal{K}(X, \mathbb{R})$  (notation as in §5 of Ch. 1) for compact separable metric spaces  $X$ . Specifically, write  $\mathcal{K} = \mathcal{K}(G(n+k, n), \mathbb{R})$ ,  $\mathcal{K}^+ = \{\beta \in \mathcal{K} : \beta \geq 0\}$ , and let  $\beta_1, \beta_2, \dots \in \mathcal{K}^+$  be dense in  $\mathcal{K}^+$ . By the XXX Theorem 4.16 of Ch. 1 we know that (since there is a Radon measure  $\gamma_j$  on  $\mathbb{R}^{n+k}$  characterized by  $\gamma_j(B) = \int_{G_n(B)} \beta_j(S) dV(y, S)$  for Borel sets  $B \subset \mathbb{R}^{n+k}$ )

$$(1) \quad e(x, j) = \lim_{\rho \downarrow 0} \frac{\int_{G_n(B_\rho(x))} \beta_j(S) dV(y, S)}{\mu_V(B_\rho(x))}$$

exists for each  $x \in \mathbb{R}^{n+k} \setminus Z_j$ , where  $Z_j$  is a Borel set with  $\mu_V(Z_j) = 0$  and  $e(x, j)$  is a  $\mu_V$ -measurable function of  $x$ , with

$$(2) \quad \int_A e(x, j) d\mu_V(x) = \int_{G_n(A)} \beta_j(S) dV(y, S)$$

for any Borel set  $A \subset \mathbb{R}^{n+k}$ .

Now let  $\varepsilon > 0$ ,  $\beta \in \mathcal{K}^+$ ,  $x \in \mathbb{R}^{n+k} \setminus (\cup_{j=1}^\infty Z_j)$ , and choose  $\beta_j$  such that  $\sup |\beta - \beta_j| < \varepsilon$ . Then for any  $\rho > 0$

$$(3) \quad \left| \frac{\int_{G_n(B_\rho(x))} \beta(S) dV(y, S)}{\mu_V(B_\rho(x))} - \frac{\int_{G_n(B_\rho(x))} \beta_j(S) dV(y, S)}{\mu_V(B_\rho(x))} \right| \leq \varepsilon \frac{V(G_n(B_\rho(x)))}{\mu_V(B_\rho(x))} = \varepsilon,$$

and hence by (1) we conclude that

$$(4) \quad \tilde{\eta}_V^{(x)}(\beta) \equiv \lim_{\rho \downarrow 0} \frac{\int_{G_n(B_\rho(x))} \beta(S) dV(y, S)}{\mu_V(B_\rho(x))}$$

exists for all  $\beta \in \mathcal{K}^+$  and all  $x \in \mathbb{R}^{n+k} \setminus (\cup_{j=1}^\infty Z_j)$ . Of course, since  $|\tilde{\eta}_V^{(x)}(\beta)| \leq \sup |\beta| \forall \beta \in \mathcal{K}^+$ , by the Riesz Representation Theorem 5.14 of Ch. 1 we have  $\tilde{\eta}_V^{(x)}(\beta) = \int_{G(n+k, n)} \beta(S) d\eta_V^{(x)}(S)$ , where  $\eta_V^{(x)}$  is the total variation measure associated with  $\tilde{\eta}_V^{(x)}$ .

Finally the last part of the lemma follows directly from (2), (3) if we keep in mind that  $e(x, j)$  in (1) is exactly  $\tilde{\eta}_V^{(x)}(\beta_j) \int_{G(n+k, n)} \beta_j(S) d\eta_V^{(x)}(S) \quad \square$

We are now able to give the proof of 1.2.

**Proof of 1.2:** By definition 1.1,  $\mu_V$  has approximate tangent space  $T_x$  with multiplicity  $\theta(x)$  in the sense of 1.7 of Ch.3 for  $\mu_V$ -a.e.  $x \in U$ . Hence by 1.9 of Ch.3 we have that  $M$  is  $\mathcal{H}^n$ -measurable countably  $n$ -rectifiable,  $\theta$  is locally  $\mathcal{H}^n$ -integrable on  $M$  and in fact  $\mu_V = \mathcal{H}^n \llcorner \theta$  in  $U$  (if we set  $\theta \equiv 0$  in  $U \setminus M$ ).

Now if  $x \in M$  is one of the  $\mu_V$ -almost all points such that  $\eta_V^{(x)}$  exists, and if  $\beta$  is a non-negative continuous function on  $G(n+k, n)$ , then we evidently have  $\eta_V^{(x)}(\beta) = \theta(x)\beta(T_x)$  and hence by the second part of 1.3 we have

$$(1) \quad \int_{G_n(A)} \beta(S) dV(x, S) = \int_{M \cap A} \beta(T_x) d\mu_V(x)$$

for any Borel set  $A \subset U$ . From the arbitrariness of  $A$  and  $\beta$  it then easily follows that

$$(2) \quad \int_{G_n(U)} f(x, S) dV(x, S) = \int_M f(x, T_x) d\mu_V(x)$$

for any non-negative  $f \in C_c(G_n(U))$ , and hence we have shown  $V = \underline{v}(M, \theta)$  as required (because  $\mu_V = \mathcal{H}^n \llcorner \theta$  as mentioned above).  $\square$

## 2 First Variation

We can make sense of first variation for a general varifold  $V$  on  $U$ . We first need to discuss *mapping* of such a general  $n$ -varifold. Suppose  $U, \tilde{U}$  open  $\subset \mathbb{R}^{n+k}$  and  $f : U \rightarrow \tilde{U}$  is  $C^1$  with  $f|_{\text{spt } \mu_V \cap U}$  proper. Then we define the image varifold  $f_\# V$  on  $\tilde{U}$  by

$$2.1 \quad f_\# V(A) = \int_{F^{-1}(A)} J_S f(x) dV(x, S), \quad A \text{ Borel}, \quad A \subset G_n(\tilde{U}),$$

where  $F : G_n^+(U) \rightarrow G_n(\tilde{U})$  is defined by  $F(x, S) = (f(x), df_x(S))$  and where

$$J_S f(x) = (\det((df_x|S)^* \circ (df_x|S)))^{\frac{1}{2}}, \quad (x, S) \in G_n(U). \\ G_n^+(U) = \{(x, S) \in G_n(U) : J_S f(x) \neq 0\}.$$

(Notice that this agrees with our previous definition given in §1 of Ch.4 in case  $V = \underline{v}(M, \theta)$ .)

Now given any  $n$ -varifold  $V$  on  $U$  we define the *first variation*  $\delta V$  of  $V$ , which is a linear functional on  $\mathcal{K}(U, \mathbb{R}^{n+k})$  (notation as in §5 of Ch. 1) by

$$2.2 \quad \delta V(X) = \left. \frac{d}{dt} \mathbb{M}(\varphi_{t\#} V \llcorner G_n(K)) \right|_{t=0},$$

where  $\{\varphi_t\}_{-1 < t < 1}$  is any 1-parameter family as in 5.8 of Ch. 2 (and  $K$  compact is as in 5.8 of Ch. 2). Of course we can compute  $\delta V(X)$  explicitly by differentiation under the integral in 2.1. This gives (by *exactly* the computations in §6 of Ch. 2)

$$2.3 \quad \delta V(X) = \int_{G_n(U)} \operatorname{div}_S X(x) dV(x, S),$$

where, for any  $S \in G(n+k, n)$ ,

$$2.4 \quad \operatorname{div}_S X = \sum_{i=1}^{n+k} \nabla_i^S x^i = \sum_{i=1}^n \langle \tau_i, D_{\tau_i} X \rangle,$$

where  $\tau_1, \dots, \tau_n$  is an orthonormal basis for  $S$  and  $\nabla_i^S = e_i \cdot \nabla^S$ , with  $\nabla^S f(x) = p_S(\nabla_{\mathbb{R}^{n+k}} f(x))$ ,  $f \in C^1(U)$ . ( $p_S$  is the orthogonal projection of  $\mathbb{R}^{n+k}$  onto  $S$ .)

By analogy with 2.4 of Ch. 4 we then say that  $V$  is *stationary in  $U$*  if  $\delta V(X) = 0 \ \forall X \in \mathcal{K}(U, \mathbb{R}^{n+k})$ .

More generally  $V$  is said to have *locally bounded first variation in  $U$*  if for each  $W \subset \subset U$  there is a constant  $c < \infty$  such that  $|\delta V(X)| \leq c \sup_U |X| \ \forall X \in \mathcal{K}(U, \mathbb{R}^{n+k})$  with  $\operatorname{spt} |X| \subset W$ . Evidently, by the general Riesz Representation 5.14 of Ch. 1, this is equivalent to the requirement that there is a Radon measure  $\|\delta\|$  (the total variation measure of  $\delta V$ ) on  $U$  characterized by

$$2.5 \quad \|\delta V\|(W) = \sup_{X \in \mathcal{K}(U, \mathbb{R}^{n+k}), |X| \leq 1, \operatorname{spt} |X| \subset W} |\delta V(X)| \quad (< \infty)$$

for any open  $W \subset \subset U$ . Notice that then by 5.14 of Ch. 1 we can write

$$2.6 \quad \delta V(X) = \int_{G_n(U)} \operatorname{div}_S X(x) dV(x, S) \equiv - \int_U v \cdot X d\|\delta V\|,$$

where  $v$  is  $\|\delta V\|$ -measurable with  $|v| = 1$   $\|\delta V\|$ -a.e. in  $U$ . By XXX Theorem 4.16 of Ch. 1 we know furthermore that

$$2.7 \quad D_{\mu_V} \|\delta V\|(x) \equiv \lim_{\rho \downarrow 0} \frac{\|\delta V\|(B_\rho(x))}{\mu_V(B_\rho(x))}$$

exists  $\mu_V$ -a.e. and that (writing  $\underline{H}(x) = D_{\mu_V} \|\delta V\|(x)v(x)$ )

$$2.8 \quad \int_U v \cdot X d\|\delta V\| = \int_U \underline{H} \cdot X d\mu_V + \int_U v \cdot X d\sigma,$$

with

$$2.9 \quad \sigma = \|\delta V\| \llcorner Z, \quad Z = \{x \in U : D_{\mu_V} \|\delta V\|(x) = +\infty\}. \quad (\mu_V(Z) = 0.)$$

Thus we can write

$$2.10 \quad \begin{aligned} \delta V(x) &= \int_{G_n(U)} \operatorname{div}_S X(x) \, dV(x, S) \\ &= - \int_U \underline{H} \cdot X \, d\mu_V - \int_Z \nu \cdot X \, d\sigma \end{aligned}$$

for  $X \in \mathcal{K}(U, \mathbb{R}^{n+k})$ .

By analogy with the classical identity 5.7 of Ch. 2 we call  $\underline{H}$  *the generalized mean curvature of  $V$* ,  $Z$  *the generalized boundary of  $V$* ,  $\sigma$  *the generalized boundary measure of  $V$* , and  $\nu|_Z$  *the generalized unit co-normal of  $V$* .

### 3 Monotonicity and Consequences

In this section we assume that  $V$  is an  $n$ -varifold in  $U$  with locally bounded first variation in  $U$  (as in 2.5).

Choose (as in 2.3 of Ch. 4)  $X|_y = h(y)\gamma(r)(x - \xi)$  where  $\gamma : \mathbb{R} \rightarrow \mathbb{R}$  is  $C^1$  and  $h \in C^1(U)$  are such that  $h\gamma$  has compact support in  $U$  (as in §3 of Ch. 4). Note that (by essentially the same computation as in §3 of Ch. 4)

$$3.1 \quad \operatorname{div}_S X = n\gamma(r) + r\gamma'(r) \sum_{i,j=1}^{n+k} e_S^{ij} \frac{x^i - \xi^i}{r} \frac{x^j - \xi^j}{r},$$

where  $(e_S^{ij})$  is the matrix of the orthogonal projection  $p_S$  of  $\mathbb{R}^{n+k}$  onto the  $n$ -dimensional subspace  $S$ . Thus the first variation identity

$$3.2 \quad \int_{G_n(U)} \operatorname{div}_S X(x) \, dV(x, S) = \delta V(X)$$

with  $X|_x = h(x)\gamma(r)(x - \xi)$  implies the following natural generalization of the identity 3.4 of Ch. 4:

$$3.3 \quad \int (\gamma(r)h + r\gamma'(r)|\nabla^S r|^2 h + \gamma(r)(x - \xi) \cdot \nabla^S h) \, dV(x, S) = \delta V(h\gamma(r)(x - \xi)).$$

Now consider a ball  $B_{\rho_0}(\xi) \subset U$  and  $\Lambda \geq 0$  with

$$3.4 \quad \|\delta V\|(B_\rho(\xi)) \leq \Lambda \mu_V(B_\rho(\xi)), \quad 0 < \rho < \rho_0.$$

Subject to 3.4 we can then take  $h = 1$  and  $\gamma(r) = \varphi(r/\rho)$  (again as in §4 of Ch.4) and, noting that  $\sum_{i,j=1}^{n+k} e_S^{ij} \frac{x^i - \xi^i}{r} \cdot \frac{x^j - \xi^j}{r} = 1 - |p_{S^\perp}(\frac{x-\xi}{r})|^2$ , conclude (Cf. 4.3 of Ch.4 with  $\alpha = 1$ ) that  $e^{\Lambda\rho} \rho^{-n} \mu_V(B_\rho(\xi))$  is increasing in  $\rho$ ,  $0 < \rho < \rho_0$ , and, for  $0 < \sigma \leq \rho < \rho_0$ ,

$$3.5 \quad \Theta^n(\mu_v, \xi) \leq e^{\Lambda\sigma} \omega_n^{-1} \sigma^{-n} \mu_V(B_\sigma(\xi)) \leq e^{\Lambda\rho} \omega_n^{-1} \rho^{-n} \mu_V(B_\rho(\xi)) \\ - \omega_n^{-1} \int_{G_n(B_\rho(\xi) \setminus B_\sigma(\xi))} r^{-n-2} |p_{S^\perp}(x - \xi)|^2 dV(x, S).$$

In fact if  $\Lambda = 0$  (so that  $V$  is stationary in  $B_{\rho_0}(\xi)$ ) we get the precise identity

$$3.6 \quad \omega_n^{-1} \rho^{-n} \mu_V(B_\rho(\xi)) - \Theta^n(\mu_v, \xi) = \omega_n^{-1} \int_{G_n(B_\rho(\xi))} r^{-n-2} |p_{S^\perp}(x - \xi)|^2 dV(x, S).$$

By a similar argument (using 3.3 with an arbitrary  $h \in C^1(U)$  rather than the special choice  $h = 1$  used above) we also deduce that the following analogue of 6.1 of Ch.4:

$$3.7 \quad \frac{d}{d\rho}(\rho^{-n} \tilde{I}(\rho)) = \rho^{-n} \frac{d}{d\rho} \int |p_{S^\perp}(x - \xi)/r|^2 \varphi(r/\rho) h(y) dV(x, S) \\ + \rho^{-n-1} (\delta V(X) + \int (x - \xi) \cdot \nabla^S h(y) \varphi(r/\rho) dV(y, S)),$$

where  $\tilde{I}(\rho) = \int \varphi(r/\rho) h d\mu_V$  and  $X|_x = h\gamma(r)(x - \xi)$ .

**3.8 Lemma.** *Suppose  $V$  has locally bounded first variation in  $U$ . Then for  $\mu_V$ -a.e.  $x \in U$ ,  $\Theta^n(\mu_V, x)$  exists and is real-valued; in fact  $\Theta^n(\mu_V, x)$  exists whenever there is a constant  $\Lambda(x) < \infty$  such that*

$$\|\delta V\|(B_\rho(x)) \leq \Lambda(x) \mu_V(B_\rho(x)), \quad 0 < \rho < \frac{1}{2} \text{dist}(x, \partial U).$$

(Such a constant  $\Lambda(x)$  exists for  $\mu_V$ -a.e.  $x \in U$  by virtue of Theorem 4.16 of Ch.1.)

Furthermore  $\Theta^n(\mu_V, x)$  is a  $\mu_V$ -measurable function of  $x$ .

**Proof:** The first part of the lemma follows directly from the monotonicity formula 3.5.

The  $\mu_V$ -measurability of  $\Theta^n(\mu_V, \cdot)$  follows from the fact that  $\mu_V(B_\rho(x)) \geq \limsup_{y \rightarrow x} \mu_V(B_\rho(y))$  which guarantees that  $\mu_V(B_\rho(x))/(\omega_n \rho^n)$  is Borel measurable and hence  $\mu_V$ -measurable for each fixed  $\rho$ . Since

$$\Theta^n(\mu_V, x) = \lim_{\rho \downarrow 0} (\omega_n \rho^n)^{-1} \mu_V(B_\rho(x))$$

for  $\mu_V$ -a.e.  $x \in U$ , we then have  $\mu_V$ -measurability of  $\Theta^n(\mu_V, \cdot)$  as claimed.

**3.9 Theorem. (Semi-continuity of  $\Theta^n$  under varifold convergence.)** *Suppose  $V_i \rightarrow V$  (as Radon measures in  $G_n(U)$ ) and  $\Theta^n(V_i, y) \geq 1$  except on a set  $B_i \subset U$  with  $\mu_{V_i}(B_i) \cap$*

$W) \rightarrow 0$  for each  $W \subset\subset U$ , and suppose that each  $V_i$  has locally bounded first variation in  $U$  with  $\liminf \|\delta V_i\|(W) < \infty$  for each  $W \subset\subset U$ . Then  $\|\delta V\|(W) \leq \liminf \|\delta V_i\|(W) \quad \forall W \subset\subset U$  and  $\Theta^n(\mu_V, y) \geq 1$   $\mu_V$ -a.e. in  $U$ .

**3.10 Remarks:** (1) The fact that  $\|\delta V\|(W) \leq \liminf \|\delta V_i\|(W)$  is a trivial consequence of the definitions of  $\|\delta V\|$ ,  $\|\delta V_i\|$  and the fact that  $V_i \rightarrow V$ , so we have only to prove the last conclusion that  $\Theta^n(\mu_V, y) \geq 1$   $\mu_V$ -a.e.

(2) The proof that  $\Theta^n(\mu_V, y) \geq 1$   $\mu_V$ -a.e. to be given below is slightly complicated; the reader should note that if  $\|\delta V\| \leq \Lambda \mu_V$  in  $U$  (i.e. if  $V$  has generalized boundary measure  $\sigma = 0$  and bounded  $\underline{H}$ —see 2.10 above—then the result is a very easy consequence of the monotonicity formula 3.5.

**Proof of 3.9:** Set  $\mu_i = \mu_{V_i}$ ,  $\mu = \mu_V$ , and take any  $W \subset\subset U$  and  $\rho_0 \in (0, \text{dist}(W, \partial U))$ . For  $i, j \geq 1$ , consider the set  $A_{i,j}$  consisting of all points  $y \in W \setminus B_i$  such that

$$(1) \quad \|\delta V_i\|(B_\rho(y)) \leq j\mu_i(B_\rho(y)), \quad 0 < \rho < \rho_0,$$

and let  $B_{i,j} = W \setminus A_{i,j}$ . Then if  $x \in B_{i,j}$  we have *either*  $x \in B_i \cap W$  or

$$(2) \quad \mu_i(B_\sigma(x)) \leq j^{-1}\|\delta V_i\|(B_\sigma(x)) \text{ for some } \sigma \in (0, \rho_0).$$

Let  $\mathcal{B}$  be the collection of balls  $B_\sigma(x)$  with  $x \in B_{i,j}$ ,  $\sigma \in (0, \rho_0)$ , and with (2) holding. By the Besicovitch Covering Lemma (§4.5 of Ch. 1) there are families  $\mathcal{B}_1, \dots, \mathcal{B}_N \subset \mathcal{B}$  with  $N = N(n+k)$ , with  $B_{i,j} \setminus B_i \subset \cup_{\ell=1}^N (\cup_{B \in \mathcal{B}_\ell} B)$  and with each  $\mathcal{B}_\ell$  a pairwise disjoint family. Hence if we sum in (2) over balls  $B \in \cup_{\ell=1}^N \mathcal{B}_\ell$ , we get

$$\mu_i(B_{i,j}) \leq Nj^{-1}\|\delta V_i\|(\widetilde{W}) + \mu_i(B_i \cap W)$$

( $\widetilde{W} = \{x \in U : \text{dist}(x, W) < \rho_0\}$ ), so

$$\mu_i(B_{i,j}) \leq cj^{-1} + \mu_i(B_i \cap W),$$

with  $c$  independent of  $i, j$ . In particular for each  $i, j \geq 1$

$$(3) \quad \mu(\text{interior}(\cap_{\ell=i}^\infty B_{\ell,j})) \leq \liminf_{q \rightarrow \infty} \mu_q(\text{interior}(\cap_{\ell=i}^\infty B_{\ell,j})) \leq cj^{-1},$$

since  $\mu_q(B_q \cap W) \rightarrow 0$  as  $q \rightarrow \infty$ .

Now let  $j \in \{1, 2, \dots\}$  and consider the possibility that there is a point  $x \in W$  such that  $x \in W \setminus \text{interior}(\cap_{q=i}^\infty B_{q,j})$  for *each*  $i = 1, 2, \dots$ . Then we could select, for each  $i = 1, 2, \dots$ ,  $y_i \in W \setminus \cap_{q=i}^\infty B_{q,j}$  with  $|y_i - x| < 1/i$ . Thus there are sequences  $y_i \rightarrow x$  and  $q_i \rightarrow \infty$  such that  $y_i \notin B_{q_i,j}$  for each  $i = 1, 2, \dots$ . Then  $y_i \in A_{q_i,j}$  and hence (by (1))

$$\|\delta V_{q_i}\|(B_\rho(y_i)) \leq j\mu_{q_i}(B_\rho(y_i)), \quad 0 < \rho < \rho_0,$$

for all  $i = 1, 2, \dots$ . Then by the monotonicity formula 3.5 (with  $\Lambda = j$ ) together with the fact that  $\Theta^n(\mu_{q_i}, y_i) \geq 1$  we have

$$\mu_{q_i}(B_\rho(y_i)) \geq e^{-j\rho} \omega_n \rho^n, \quad 0 < \rho < \rho_0,$$

so that  $\Theta^n(\mu, x) \geq 1$  for such an  $x$ . Thus we have proved  $\Theta^n(\mu, x) \geq 1$  for each  $x$  with  $x \in W \setminus (\cup_{i=1}^\infty \text{interior}(\cap_{\ell=i}^\infty B_{\ell,j}))$  for some  $j \in \{1, 2, \dots\}$ . That is

$$(4) \quad \Theta^n(\mu, x) \geq 1 \quad \forall x \in W \setminus (\cap_{j=1}^\infty \cup_{i=1}^\infty \text{interior}(\cap_{\ell=i}^\infty B_{\ell,j})).$$

However

$$\begin{aligned} (5) \quad \mu(\cap_{j=1}^\infty \cup_{i=1}^\infty \text{interior}(\cap_{\ell=i}^\infty B_{\ell,j})) &\leq \mu(\cup_{i=1}^\infty \text{interior}(\cap_{\ell=i}^\infty B_{\ell,j})) \quad \forall j \geq 1 \\ &= \lim_{i \rightarrow \infty} \mu(\text{interior}(\cap_{\ell=i}^\infty B_{\ell,j})) \\ &\leq c j^{-1} \text{ by (3),} \end{aligned}$$

so  $\mu(\cap_{j=1}^\infty \cup_{i=1}^\infty \text{interior}(\cap_{\ell=i}^\infty B_{\ell,j})) = 0$  and the theorem is established (by (4)).  $\square$

## 4 Constancy Theorem

**4.1 (Constancy Theorem.)** *Suppose  $V$  is an  $n$ -varifold in  $U$ ,  $V$  is stationary in  $U$ , and  $U \cap \text{spt } \mu_V \subset M$ , where  $M$  is a connected  $n$ -dimensional  $C^2$  embedded submanifold of  $\mathbb{R}^{n+k}$ . Then  $V = \theta_0 \underline{v}(M)$  for some constant  $\theta_0$ .*

**4.2 Remarks:** (1) Notice in particular this implies  $(\overline{M} \setminus M) \cap U = \emptyset$  (if  $V \neq 0$ ); this is not *a-priori* obvious from the assumptions of the theorem.

(2) J. Duggan in his PhD thesis [Dug86] has extended 4.1 to the case when  $M$  is merely Lipschitz.

(3) The reader will see that, with only minor modifications to the proof to be given below, the theorem continues to hold if  $N$  is an embedded  $(n+k)$ -dimensional  $C^2$  submanifold of  $\mathbb{R}^{n+k}$  and if  $V$  is stationary in  $U \cap N$  in the sense that  $\delta V(X) = 0 \quad \forall X \in \mathcal{K}(U; \mathbb{R}^{n+k})$  with  $X_x \in T_x N \quad \forall x \in N$ , *provided we are given*  $\text{spt } V \subset \{(x, S) : x \in N \text{ and } S \subset T_x N\}$ . (This last is equivalent to  $\text{spt } \mu_V \subset N$  and  $p_\# V = V$ , where  $p : U \rightarrow U \cap N$  coincides with the nearest point projection onto  $U \cap N$  in some neighborhood of  $U \cap N$ .)

**Proof of 4.1:** We first want to argue that  $V = \underline{v}(M, \theta)$  for some positive locally  $\mathcal{H}^n$ -integrable function  $\theta$  on  $M$ .

To do this first take any  $f \in C_c^2(U)$  with  $M \subset \{x \in U : f(x) = 0\}$  and note that by 2.3

$$(1) \quad \delta V(f \nabla f) = \int |p_S(\nabla f)|^2 dV(x, S),$$

because (using notation as in 2.3)

$$\begin{aligned} \operatorname{div}_S(f \nabla f) &= \nabla^S f \cdot \nabla f + f \operatorname{div}_S \nabla f \\ &= |p_S(\nabla f)|^2 \text{ on } M, \end{aligned}$$

where we used  $f \equiv 0$  on  $M$ . Since  $\delta V = 0$ , we conclude from (1) that

$$(2) \quad p_S(\nabla f(x)) = 0 \quad \text{for all } (x, S) \in \operatorname{spt} V.$$

Now let  $\xi \in M$  be arbitrary. We can find an open  $W \subset U$  with  $\xi \in W$  and such that there are  $C_c^2(U)$  functions  $f_1, \dots, f_k$  with  $M \subset \bigcap_{j=1}^k \{x : f_j(x) = 0\}$  and with  $(T_x M)^\perp$  being exactly the space spanned by  $\nabla f_1(x), \dots, \nabla f_k(x)$  for each  $x \in M \cap W$ . (One easily checks that such  $W$  and  $f_1, \dots, f_k$  exists.) Then (2) implies that

$$(3) \quad p_S((T_x M)^\perp) = 0 \quad \text{for all } (x, S) \in G_n(W) \cap \operatorname{spt} V.$$

But (3) says exactly that  $S = T_x M$  for all  $(x, S) \in G_n(W) \cap \operatorname{spt} V$ , so that (since  $\xi$  was an arbitrary point of  $M$ ), we have

$$(4) \quad \int f(x, S) dV(x, S) = \int_{M \cap U} f(x, T_x M) d\mu(x), \quad f \in C_c(G_n(U)),$$

On the other hand we know from monotonicity 3.5 that  $\theta(x) \equiv \Theta^n(\mu_V, x)$  exists for all  $x \in M \cap U$ , and hence (since  $\Theta^n(\mathcal{H}^n \llcorner M, x) = 1$  for each  $x \in M$ , by smoothness of  $M$ ), we can use the XXX Theorem 4.16 of Ch. 1 to conclude from (4) that in fact

$$(5) \quad \int f(x, S) dV(x, S) = \int_{M \cap U} f(x, T_x M) \theta(x) d\mathcal{H}^n(x), \quad f \in C_c(G_n(U)),$$

(so that  $V = \underline{v}(M, \theta)$  as required).

It thus remains only to prove that  $\theta = \text{const.}$  on  $M \cap U$ . Since  $M$  is  $C^2$  we can take  $X \in \mathcal{K}(U, \mathbb{R}^{n+k})$  such that  $X_x \in T_x M \quad \forall x \in M \cap U$ . Then by (5) and 2.3  $\delta V(X) = 0$  is just the statement that  $\int_{M \cap U} \operatorname{div} X \theta d\mathcal{H}^n = 0$ , where  $\operatorname{div} X$  is the classical divergence of  $X|_M$  in the usual sense of differential geometry. Using local coordinates (in some neighborhood  $\tilde{U} \subset \mathbb{R}^n$ ) this tells us that

$$\int_{\tilde{U}} \sum_{i=1}^n \frac{\partial X_i}{\partial x_i} \tilde{\theta} d\mathcal{L}^n = 0 \quad \text{if } X_i \in C_c^1(\tilde{U}), \quad i = 1, \dots, n,$$

where  $\tilde{\theta}$  is  $\theta$  expressed in terms of the local coordinates. In particular

$$\int_{\tilde{U}} \frac{\partial \zeta}{\partial x_i} \tilde{\theta} d\mathcal{L}^n = 0 \quad \forall \zeta \in C_c(U), \quad i = 1, \dots, n$$

and it is then standard that  $\tilde{\theta} = \text{constant}$  in  $\tilde{U}$ . Hence (since  $M$  is connected)  $\theta$  is constant in  $M$ .  $\square$

## 5 Varifold Tangents and Rectifiability Theorem

Let  $V$  be a  $n$ -varifold in  $U$  and let  $x$  be any point of  $U$  such that

$$5.1 \quad \Theta^n(\mu_V, x) = \theta_0 \in (0, \infty) \quad \text{and} \quad \lim_{\rho \downarrow 0} \rho^{1-n} \|\delta V\|(B_\rho(x)) = 0.$$

By definition of  $\delta V$  (in 2) and the Compactness Theorem 5.15 of Ch. 1 for Radon measures, we can select a sequence  $\lambda_j \downarrow 0$  such that  $\eta_{x, \lambda_j \#} V$  converges (in the sense of Radon measures) to a varifold  $C$  such that

$$C \text{ is stationary in } \mathbb{R}^{n+k}$$

and

$$5.2 \quad \frac{\mu_C(B_\rho(x))}{\omega_n \rho^n} \equiv \theta_0 \quad \forall \rho > 0.$$

Since  $\delta C = 0$  we can use 5.2 together with the monotonicity formula 3.6 to conclude

$$\int_{G_n(B_\rho(0))} \frac{|p_{S^\perp}(x)|^2}{|x|^{n+2}} dC(x, S) = 0 \quad \forall \rho > 0,$$

so that  $p_{S^\perp}(x) = 0$  for  $C$ -a.e.  $(x, S) \in G_n(\mathbb{R}^{n+k})$ , and hence  $p_{S^\perp}(x) = 0$  for *all*  $(x, S) \in \text{spt } C$  by continuity of  $p_{S^\perp}(x)$  in  $(x, S)$ . We can apply the same argument as in the proof of Theorem 5.1 of Ch. 4 with  $\delta = 0$ , except that we now use 3.7 in place of 6.1 of Ch. 4, so  $\mu_C$  satisfies

$$5.3 \quad \Theta^n(\mu_C, y) = \Theta^n(\mu_C, \lambda y) \quad \forall \lambda > 0$$

We would *like* to prove the stronger result  $\eta_{0, \lambda \#} C = C$  (which of course implies 5.3), but we are only able to do this in case  $\Theta^n(\mu_C, x) > 0$  for  $\mu_C$ -a.e.  $x$  (see 5.7 below). Whether or not  $\eta_{0, \lambda \#} C = C$  without the additional hypothesis on  $\Theta^n(\mu_C, \cdot)$  seems to be an open question.

**5.4 Definition:** Given  $V$  and  $x$  as in 5.1 we let  $\text{Var Tan}(V, x)$  (“the varifold tangent of  $V$  at  $x$ ”) be the collection of all  $C = \lim \eta_{x, \lambda_j \#} V$  obtained as described above.

Notice that by the above discussion any  $C \in \text{VarTan}(V, x)$  is stationary in  $\mathbb{R}^{n+k}$  and satisfies 5.3.

The following rectifiability theorem for  $n$ -varifolds is a central part of the theory of  $n$ -varifolds with locally bounded first variation.

**5.5 Theorem (Rectifiability Theorem.)** *Suppose  $V$  has locally bounded first variation in  $U$  and  $\Theta^n(\mu_V, x) > 0$  for  $\mu_V$ -a.e.  $x \in U$ . Then  $V$  is an  $n$ -rectifiable varifold. (Thus  $V = \underline{v}(M, \theta)$ , with  $M$  a  $\mathcal{H}^n$ -measurable countably  $n$ -rectifiable subset of  $U$  and  $\theta$  a non-negative locally  $\mathcal{H}^n$ -integrable function on  $U$ .)*

**5.6 Remark:** We are going to use 1.2. In fact we show that  $V$  has a tangent plane (in the sense of 1.1) at the point  $x$  where (i)  $\Theta^n(\mu_V, x) > 0$ , (ii)  $\eta_V^{(x)}$  (as in 1.3) exists, (iii)  $\Theta^n(\mu_V, \cdot)$  is  $\mu_V$ -approximately continuous at  $x$ , and (iv)  $\|\delta V\|(B_\rho(x)) \leq \Lambda(x)\mu_V(B_\rho(x))$  for  $0 < \rho < \rho_0 = \min\{1, \text{dist}(x, \partial U)\}$ . Since conditions (i)–(iv) all hold  $\mu_V$ -a.e. in  $U$  (notice that (iii) holds  $\mu_V$ -a.e. by virtue of the  $\mu_V$ -measurability of  $\Theta^n(\mu_V, \cdot)$  proved in 3.8), the required rectifiability of  $V$  will then follow from 1.2

Before beginning the proof of 5.5 we give the following important corollary.

**5.7 Corollary.** *Suppose  $x \in U$ , 5.1 holds, and  $\lim_{\rho \downarrow 0} \rho^{-n} \mu_V(\{y \in B_\rho(x) : \Theta^n(\mu_V, y) < 1\}) = 0$ . If  $C \in \text{VarTan}(V, x)$ , then  $C$  is rectifiable and*

$$\eta_{0, \lambda \#} C = C \quad \forall \lambda > 0.$$

**Proof:** From the hypothesis  $\rho^{-n} \mu_V(\{y \in B_\rho(x) : \Theta^n(\mu_V, y) < 1\}) \rightarrow 0$  and the Semi-continuity 3.9, we have  $\Theta^n(\mu_C, y) \geq 1$  for  $\mu_C$ -a.e.  $y \in \mathbb{R}^{n+k}$ . Hence by 5.5 we have that  $C$  is  $n$ -rectifiable. On the other hand, since  $\Theta^n(\mu_C, y) = \Theta^n(\mu_C, \lambda y) \quad \forall \lambda > 0$  (by 5.3), we can write  $C = \underline{v}(M, \theta)$  with  $\eta_{0, \lambda}(M) = M \quad \forall \lambda > 0$  and  $\theta(\lambda y) = \theta(y) \quad \forall \lambda > 0, y \in \mathbb{R}^{n+k}$ . (Viz. simply set  $\theta(y) = \Theta^n(\mu_C, y)$  and  $M = \{y \in \mathbb{R}^{n+k} : \theta(y) > 0\}$ .) It then trivially follows that  $y \in T_y M$  whenever the approximate tangent space  $T_y M$  exists, and hence  $\eta_{0, \lambda \#} C = C$  as required.  $\square$

**Proof of 5.5:** Let  $x$  be as in 5.6(i)–(iv) and take  $C \in \text{VarTan}(V, x)$ . (We know  $\text{VarTan}(V, x) \neq \emptyset$  because 5.6(i), (iv) imply 5.1.) Then  $C$  is stationary in  $\mathbb{R}^{n+k}$  and

$$(1) \quad \frac{\mu_C(B_\rho(0))}{\omega_n \rho^n} \equiv \theta_0 \quad \forall \rho > 0 \quad (\theta_0 = \Theta^n(\mu_V, x)).$$

Also for any  $y \in \mathbb{R}^{n+k}$  (using (1) and the monotonicity formula 3.5)

$$\begin{aligned} \frac{\mu_C(B_\rho(y))}{\omega_n \rho^n} &\leq \frac{\mu_C(B_R(y))}{\omega_n R^n} \leq \frac{\mu_C(B_{R+|y|}(0))}{\omega_n (R+|y|)^n} (1 + |y|/R)^n \\ &= \theta_0 (1 + |y|/R)^n \rightarrow \theta_0 \text{ as } R \uparrow \infty. \end{aligned}$$

That is (again using the monotonicity formula 3.5),

$$(2) \quad \Theta^n(\mu_C, y) \leq \frac{\mu_C(B_\rho(y))}{\omega_n \rho^n} \leq \theta_0 \quad \forall y \in \mathbb{R}^{n+k}, \quad \rho > 0.$$

Now let  $V_j = \eta_{x, \lambda_j} V$ , where  $\lambda_j \downarrow 0$  is such that  $\lim \eta_{x, \lambda_j} V = C$  and where we are still assuming  $x$  is as in 5.6(i)–(iv).

From 5.6(iii) we have (with  $\varepsilon(\rho) \downarrow 0$  as  $\rho \downarrow 0$ )

$$(3) \quad \Theta^n(\mu_V, y) \geq \theta_0 - \varepsilon(\rho), \quad y \in G \cap B_\rho(x),$$

where  $G \subset U$  is such that

$$(4) \quad \mu_V(B_\rho(x) \setminus G) \leq \varepsilon(\rho) \rho^n, \quad \rho \text{ sufficiently small.}$$

Taking  $\rho = \lambda_j$  we see that (3), (4) imply

$$(5) \quad \Theta^n(\mu_{V_j}, y) \leq \theta_0 - \varepsilon_j, \quad y \in G_j \cap B_1(0)$$

with  $G_j$  such that

$$(6) \quad \mu_{V_j}(B_1(0) \setminus G_j) \leq \varepsilon_j,$$

where  $\varepsilon_j \rightarrow 0$  as  $j \rightarrow \infty$ . Thus, using (5), (6) and the semi-continuity result of 3.9, we obtain

$$(7) \quad \Theta^n(\mu_C, y) \geq \theta_0 \text{ for } \mu_C\text{-a.e. } y \in \mathbb{R}^{n+k}$$

(and hence for every  $y \in \text{spt } \mu_C$  by 3.6). Then by combining (2) and (7) we have

$$\Theta^n(\mu_C, y) \equiv \theta_0 \equiv \frac{\mu_C(B_\rho(y))}{\omega_n \rho^n} \quad \forall y \in \text{spt } \mu_C, \quad \rho > 0.$$

Then by the monotonicity formula 3.6 (with  $V = C$ ), we have

$$p_{S^\perp}(x - y) = 0 \text{ for } C\text{-a.e. } (x, S) \in G_n(\mathbb{R}^{n+k}).$$

Thus (using the continuity of  $p_{S^\perp}(x - y)$  in  $(x, S)$ ) we have

$$(8) \quad x - y \in S \quad \forall y \in \text{spt } \mu_C \text{ and } \forall (x, S) \in \text{spt } C.$$

In particular, choosing  $T$  such that  $(0, T) \in \text{spt } C$  (such  $T$  exists because  $0 \in \text{spt } \mu_C = \pi(\text{spt } C)$ ), (8) implies  $y \in T \quad \forall y \in \text{spt } \mu_C$ . Thus  $\text{spt } \mu_C \subset T$ , and hence  $C = \theta_0 \underline{v}(T)$  by Constancy 4.1.

Thus we have shown that, for  $x \in U$  such that 5.6(i), (iii), (iv) hold, each element of  $\text{Var Tan}(V, x)$  has the form  $\theta_0 \underline{v}(T)$ , where  $T$  is an  $n$ -dimensional subspace of  $\mathbb{R}^{n+k}$ . On the other hand, since we are assuming (5.6(ii)) that  $\eta_V^{(x)}$  exists, it follows that for continuous  $\beta$  on  $G(n+k, n)$

$$(9) \quad \lim_{\rho \downarrow 0} \frac{\int_{G_n(B_\rho(x))} \beta(S) dV(y, S)}{\mu_V(B_\rho(x))} = \int_{G(n+k, n)} \beta(S) d\eta_V^{(x)}(S).$$

Now let  $\theta_0 \underline{v}(T)$  be any such element of  $\text{Var Tan}(V, x)$  and select  $\lambda_j \downarrow 0$  so that  $\lim \eta_{x, \lambda_j \#} V = \theta_0 \underline{v}(T)$ . Then in particular

$$\lim_{j \rightarrow \infty} \frac{\int_{G_n(B_1(0))} \beta(S) dV_j(y, S)}{\mu_{V_j}(B_1(0))} = \beta(T),$$

and hence (9) gives

$$\beta(T) = \int_{G(n+k, n)} \beta(S) d\eta_V^{(x)}(S),$$

thus showing that  $\theta_0 \underline{v}(T)$  is the *unique* element of  $\text{Var Tan}(V, x)$ . Thus

$$\lim_{\lambda \downarrow 0} \eta_{x, \lambda \#} V = \theta_0 \underline{v}(T),$$

so that  $T$  is the tangent space for  $V$  at  $x$  in the sense of 1.1. This completes the proof.  $\square$

The following *compactness theorem* for rectifiable varifolds is now a direct consequence of the Rectifiability 5.5, the Semi-continuity 3.9, and the Compactness Theorem 5.15 of Ch. 1 for Radon measures, and its proof is left to the reader.

**5.8 Theorem (Compactness theorem for  $n$ -varifolds.)** *Suppose  $\{V_j\}$  is a sequence of rectifiable  $n$ -varifolds in  $U$  which are locally bounded first variation in  $U$ ,*

$$\sup_{j \geq 1} (\mu_{V_j}(W) + \|\delta V_j\|(W)) < \infty \quad \forall W \subset\subset U,$$

*and  $\Theta^n(\mu_{V_j}, x) \geq 1$  on  $U \setminus A_j$ , where  $\mu_{V_j}(A_j \cap W) \rightarrow 0$  as  $j \rightarrow \infty$   $\forall W \subset\subset U$ .*

*Then there is a subsequence  $\{V_{j'}\}$  and a rectifiable varifold  $V$  of locally bounded first variation in  $U$ , such that  $V_{j'} \rightarrow V$  (in the sense of Radon measures on  $G_n(U)$ ),  $\Theta^n(\mu_V, x) \geq 1$  for  $\mu_V$ -a.e.  $x \in U$ , and  $\|\delta V\|(W) \leq \liminf_{j \rightarrow \infty} \|\delta V_j\|(W)$  for each  $W \subset\subset U$ .*

**5.9 Remark:** An important additional result (also due to Allard [All72]) is the Integral Compactness Theorem, which asserts that if all the  $V_j$  in the above theorem are integer multiplicity, then  $V$  is also integer multiplicity. (Notice that in this case the hypothesis  $\Theta^n(\mu_{V_j}, x) \geq 1$  on  $U \setminus A_j$  is automatically satisfied with an  $A_j$  such that  $\mu_{V_j}(A_j) = 0$ .)

**Proof that  $V$  is integer multiplicity if the  $V_i$  are:** Let  $W \subset\subset U$ . We first assert that for  $\mu_V$ -a.e.  $x \in W$  there exists  $c$  (depending on  $x$ ) such that

$$(1) \quad \liminf \|\delta V_i\|(B_\rho(x)) \leq c \mu_V(B_\rho(x)), \quad \rho < \min\{1, \text{dist}(x, \partial U)\}.$$

Indeed otherwise  $\exists$  a set  $A \subset W$  with  $\mu_V(A) > 0$  such that for each  $j \geq 1$  and each  $x \in A$  there are  $\rho_x > 0$ ,  $i_x \geq 1$  such that  $B_{\rho_x}(x) \subset W$  and

$$\mu_V(B_{\rho_x}(x)) \leq j^{-1} \|\delta V_{i_x}\|(B_{\rho_x}(x)), \quad i_x \geq i.$$

By the Besicovitch Covering Lemma (§4.5 of Ch. 1) we then have

$$\mu_V(A_i) \leq c j^{-1} \|\delta V_\ell\|(W), \quad \ell \geq i,$$

where  $A_i = \{x \in A : i_x \leq i\}$ . Thus

$$\mu_V(A_i) \leq c j^{-1} \limsup_{\ell \rightarrow \infty} \|\delta V_\ell\|(W),$$

and hence  $A_i \uparrow A$  as  $i \uparrow \infty$  we have

$$\mu_V(A) \leq c j^{-1}$$

for some  $c (< \infty)$  independent of  $j$ . That is,  $\mu_V(A) = 0$ , a contradiction, and hence (1) holds. Since  $\Theta^n(\mu_V, x)$  exists  $\mu_V$ -a.e.  $x \in U$ , we in fact have from (1) that for  $\mu_V$ -a.e.  $x \in U$  there is a  $c = c(x)$  such that

$$(2) \quad \liminf \|\delta V_i\|(B_\rho(x)) \leq c \rho^n, \quad 0 < \rho < \min\{1, \text{dist}(x, \partial U)\}.$$

Now since  $V = \underline{v}(M, \theta)$ , it is also true that for  $\mu_V$ -a.e.  $\xi \in \text{spt } \mu_V$  we have  $\eta_{\xi, \lambda\#} V \rightarrow \theta_0 \underline{v}(P)$  as  $\lambda \downarrow 0$ , where  $P = T_\xi M$  and  $\theta_0 = \theta(\xi)$ . Then (because  $V_i \rightarrow V$ , and hence  $\eta_{\xi, \lambda\#} V_i \rightarrow \eta_{\xi, \lambda\#} V$  for each fixed  $\lambda > 0$ ), it follows that for  $\mu_V$ -a.e.  $\xi \in U$  we can select a sequence  $\lambda_i \downarrow 0$  such that, with  $W_i = \eta_{\xi, \lambda_i\#} V_i$ ,

$$(3) \quad W_i \rightarrow \theta_0 \underline{v}(P)$$

and (by (2)) for each  $R > 0$

$$(4) \quad \|\delta W_i\|(B_R(0)) \rightarrow 0.$$

We claim that  $\theta_0$  must be an integer for any such  $\xi$ ; in fact for an arbitrary sequence  $\{W_i\}$  of integer multiplicity varifolds in  $\mathbb{R}^{n+k}$  satisfying (3), (4), we claim that  $\theta_0$  always has to be an integer.

To see this, take (without loss of generality)  $P = \mathbb{R}^n \times \{0\}$ , let  $q$  be orthogonal projection onto  $(\mathbb{R}^n \times \{0\})^\perp$ , and note first that (3) implies

$$(5) \quad p_{\mathbb{R}^n\#}(W_i \llcorner G_n\{x \in \mathbb{R}^{n+k} : |q(x)| < \varepsilon\}) \rightarrow \theta_0 \underline{v}(\mathbb{R}^n)$$

for each fixed  $\varepsilon > 0$ . However by the mapping formula for varifolds (§1 of Ch.4), we know that (5) says

$$(6) \quad \underline{v}(\mathbb{R}^n, \psi_i) \rightarrow \theta_0 \underline{v}(\mathbb{R}^n),$$

where

$$\psi_i(x) = \sum_{y \in p_{\mathbb{R}^n \times \{0\}}^{-1}(x) \cap \{z \in \mathbb{R}^{n+k} : |q(z)| < \varepsilon\}} \theta_i(y)$$

( $\theta_i$  = multiplicity function of  $W_i$ , so that  $\psi_i$  has values in  $\mathbb{Z} \cup \{\infty\}$ ). Notice that (6) implies in particular that

$$(7) \quad \int_{\mathbb{R}^n} f \psi_i d\mathcal{L}^n \rightarrow \theta_0 \int_{\mathbb{R}^n} f d\mathcal{L}^n \quad \forall f \in C_c^0(\mathbb{R}^n).$$

(i.e. measure-theoretic convergence of  $\psi_i$  to  $\theta_0$ .)

Now we claim that there are sets  $A_i \subset B_1(0)$  such that

$$(8) \quad \psi_i(x) \leq \theta_0 + \varepsilon_i \quad \forall x \in B_1(0) \setminus A_i, \quad \mathcal{L}^n(A_i) \rightarrow 0, \quad \varepsilon_i \downarrow 0;$$

this will of course (when used in combination with (7)) imply that for any integer  $N > \theta_0$ ,  $\max\{\psi_i, N\}$  converges in  $L^1(B_1(0))$  to  $\theta_0$ , and, since  $\max\{\psi_i, N\}$  is integer-valued, it then follows that  $\theta_0$  is an integer.

On the other hand (8) evidently follows by setting  $W = W_i$  in the following lemma, so the proof is complete.  $\square$

In this lemma,  $p, q$  denote orthogonal projection of  $\mathbb{R}^{n+k}$  onto  $\mathbb{R}^n \times \{0\} \subset \mathbb{R}^{n+k}$  and  $\{0\} \times \mathbb{R}^k \subset \mathbb{R}^{n+k}$  respectively.

**5.10 Lemma.** *For each  $\delta \in (0, 1)$ ,  $\Lambda \geq 1$ , there is  $\varepsilon = \varepsilon(\delta, \Lambda, n) \in (0, \delta^2)$  such that if  $W$  is an integer multiplicity varifold in  $B_3(0)$  with*

$$(\ddagger) \quad \mu_W(B_3(0)) \leq \Lambda, \quad \|\delta W\|(B_3(0)) < \varepsilon^2, \quad \int_{B_3(0)} \|p_S - p\| dW(y, S) < \varepsilon^2$$

*there there is a set  $A \subset B_1^n(0)$  such that  $\mathcal{L}^n(A) < \delta$  and,  $\forall x \in B_1(0) \setminus A$ ,*

$$\sum_{y \in p^{-1}(x) \cap \text{spt } \mu_W \cap \{z : |q(z)| < \varepsilon\}} \Theta^n(\mu_W, y) \leq (1 + \delta) \frac{\mu_W(B_2(x))}{\omega_n 2^n} + \delta.$$

**5.11 Remark:** It suffices to prove that for each fixed  $N$  there is a  $\delta_0 = \delta_0(N) \in (0, 1)$  such that if  $\delta \in (0, \delta_0)$  then  $\exists \varepsilon = \varepsilon(n, \Lambda, N, \delta) \in (0, \delta^2)$  such that 5.10( $\ddagger$ ) implies

the existence of  $A \subset B_1^n(0)$  with  $\mathcal{L}^n(A) < \delta$  and, for  $x \in B_1^n(0) \setminus A$  and distinct  $y_1, \dots, y_N \in p^{-1}(x) \cap \text{spt } \mu_W \cap \{z : |q(z)| < \varepsilon\}$ ,

$$(\ddagger) \quad \sum_{j=1}^N \Theta^n(\mu_W, y_j) \leq (1 + \delta) \frac{\mu_W(B_2(x))}{\omega_n 2^n} + \delta.$$

Because this firstly implies an *a-priori* bound, depending only on  $n, k, \Lambda$ , on the number  $N$  of possible points  $y_j$ , and hence the lemma, as originally stated, then follows. (Notice that of course the validity of the lemma for small  $\delta$  implies its validity for any larger  $\delta$ .)

**Proof of 5.10:** By virtue of the above Remark, we need only to prove 5.11  $(\ddagger)$ . Let  $\mu = \mu_W$ , and consider the possibility that  $y \in B_1(0)$  satisfies the inequalities

$$(1) \quad \delta \|W\|(B_\rho(y)) \leq \varepsilon \mu(B_\rho(y)), \quad \rho \in (0, 1),$$

$$(2) \quad \int_{B_{\rho(y)}} \|p_S - p\| dW(z, S) \leq \varepsilon \rho^n, \quad \rho \in (0, 1).$$

Let

$$A_1 = \{y \in B_2(0) \cap \text{spt } W : (1) \text{ fails for some } \rho \in (0, 1)\}$$

$$A_2 = \{y \in B_2(0) \cap \text{spt } W : (2) \text{ fails for some } \rho \in (0, 1)\}.$$

Evidently  $y \in \text{spt } \mu_W \cap B_2(0) \setminus A_1 \Rightarrow$  (by the monotonicity formula 3.5)

$$(3) \quad \frac{\mu(B_\rho(y))}{\omega_n \rho^n} \leq e^\varepsilon \frac{\mu(B_1(y))}{\omega_n} \leq c, \quad 0 < \rho < 1,$$

( $c = c(\Lambda, n)$ ), while if  $y \in A_2 \setminus A_1$  we have (using (3))

$$(4) \quad \int_{B_{\rho(y)}} \|p_S - p\| dW(z, S) \geq \varepsilon \rho_y^n \geq c \varepsilon \mu(B_{\rho_y}(y))$$

for some  $\rho_y \in (0, 1)$ . If  $y \in A_1$  then

$$(5) \quad \mu(B_{\rho_y}(y)) \leq \varepsilon^{-1} \|\delta W\|(B_{\rho_y}(y))$$

for some  $\rho_y \in (0, 1)$ .

Since then  $\{B_{\rho_y}(y)\}_{y \in A_1 \cup A_2}$  covers  $A_1 \cup A_2$  we deduce from (4), (5) and the Besicovitch Covering (§4.5 of Ch. 1) that

$$(6) \quad \begin{aligned} \mu(A_1 \cup A_2) &\leq c \varepsilon^{-1} \left( \int_{B_3(0)} \|p_S - p\| dW(a, S) + \|\delta W\|(B_3(0)) \right) \\ &\leq c \varepsilon \end{aligned}$$

by the hypotheses on  $W$ .

Our aim now is to show 5.11 ( $\ddagger$ ) whenever  $x \in B_1^n(0) \setminus p(A_1 \cup A_2)$ . In view of (6) this will establish the required result (with  $A = p(A_1 \cup A_2)$ ). So let  $x \in B_1^n(0) \setminus p(A_1 \cup A_2)$ . In view of the monotonicity formula 3.6 it evidently suffices (by translating and changing scale by a factor of  $3/2$ ) to assume that  $x = 0 \in B_1^n(0) \setminus p(A_1 \cup A_2)$ . We shall subsequently assume this.

We first want to establish the two inequalities, that, for  $y \in B_1^n(0) \setminus p(A_1 \cup A_2)$  and  $\tau > 0$ ,

$$(7) \quad \Theta^n(\mu, y) \leq e^{\varepsilon\sigma} \frac{\mu(U_\sigma^{2\tau}(y))}{\omega_n \sigma^n} + c\varepsilon\sigma/\tau, \quad 0 < \sigma < 1,$$

$$(8) \quad \frac{\mu(U_\sigma^\tau(y))}{\omega_n \sigma^n} \leq e^{\varepsilon\sigma} \frac{\mu(U_\sigma^{2\tau}(y))}{\omega_n \sigma^n} + c\varepsilon\sigma/\tau, \quad 0 < \sigma < \rho \leq 1,$$

where

$$U_\sigma^\tau(y) = B_\sigma(y) \cap \{z \in \mathbb{R}^{n+k} : |q(z - y)| < \tau\}.$$

Indeed these two inequalities follow directly from 3.5 and 3.7. For example to establish (7) we note first that 3.5 gives (7) directly if  $\tau \geq \sigma$ , while if  $\tau < \sigma$  then we first use 3.5 to give  $\Theta^n(\mu, y) \leq e^{\varepsilon\tau} \frac{\mu(B_\tau(y))}{\omega_n \tau^n}$  and then use 3.7 with  $h$  of the form  $h(z) = f(|q(z - y)|)$ ,  $f(t) \equiv 1$  for  $t < \tau$  and  $f(t) \equiv 0$  for  $t > 2\tau$ .

Since  $|\nabla^S f(|q(z - y)|)| \leq f'(|q(z - y)|)|p_S - p|$  (Cf. the computation in ?? of Ch. 4) we then deduce (by integrating in 3.7 from  $\tau$  to  $\sigma$  and using (3))

$$\frac{\mu(B_\tau(y))}{\omega_n \tau^n} \leq \frac{\mu(U_\sigma^{2\tau}(y))}{\omega_n \sigma^n} + c\varepsilon\sigma/\tau.$$

(8) is proved by simply integrating 3.7 from  $\sigma$  to  $\rho$  (and using (3)).

Our aim now is to use (7) and (8) to establish

$$(9) \quad \sum_{j=1}^N \frac{\mu(U_\sigma^\tau(y_j))}{\omega_n \sigma^n} \leq (1 + c\delta^2) \frac{\mu(B_2(0))}{\omega_n 2^n} + c\delta^2$$

with  $c = c(n, k, N, \Lambda)$ , provided  $2\delta^2\sigma \leq \tau \leq \frac{1}{4} \min_{j \neq \ell} |y_j - y_\ell|$ ,  $y_j \in \text{spt } \mu \cap p^{-1}(0) \cap \{z : |q(z)| < \varepsilon\}$ ,  $0 \notin p(A_1 \cup A_2)$ . (In view of (7) this will prove the required result 5.11 ( $\ddagger$ ) for suitable  $\delta_0(N)$ .)

We proceed by induction on  $N$ .  $N = 1$  trivially follows from (8) by noting that  $U_\rho^{2\tau}(y_1) \subset B_\rho(y_1)$  (by definition of  $U_\rho^{2\tau}(y_1)$ ) and then using the monotonicity 3.5 together with the fact that  $|y_1| < \varepsilon$ . Thus assume  $N \geq 2$  and that (9) has been established with any  $M < N$  in place of  $N$ .

Let  $y_1, \dots, y_N$  be as in (9), and choose  $\rho \in [\sigma, 1)$  such that  $\min_{j \neq \ell} |q(y_j) - q(y_\ell)|$  ( $= \min_{j \neq \ell} |y_j - y_\ell|$ )  $= 4\delta^2\rho$ , and set  $\tilde{\tau} = 2\delta^2\rho$  ( $\geq 2\tau$ ). Then

$$\begin{aligned} \frac{\mu(U_\sigma^\tau(y_j))}{\sigma^n} &\leq \frac{\mu(U_\sigma^{\frac{1}{2}\tilde{\tau}}(y_j))}{\sigma^n} \\ &\leq e^{\varepsilon\rho} \frac{\mu(U_\rho^{\tilde{\tau}}(y_j))}{\rho^n} + c\varepsilon \quad (\text{by (8)}), \end{aligned}$$

$c = c(n, k, \delta)$ . Now since  $\tilde{\tau} = \frac{1}{2} \min_{j \neq \ell} |q(y_j) - q(y_\ell)|$  we can select  $\{z_1, \dots, z_Q\} \subset \{y_1, \dots, y_N\}$  ( $Q \leq N - 1$ ) and  $\tilde{\tau} \leq c\tilde{\tau}$  such that  $\hat{\tau} \geq 3\delta^2\rho$  and

$$\cup_{j=1}^N U_\rho^{\tilde{\tau}}(y_j) \subset \cup_{\ell=1}^Q U_{\rho(1+c\delta^2)}^{\hat{\tau}}(z_\ell),$$

where  $c = c(N)$ , and such that  $\hat{\tau} \leq \frac{1}{4} \min_{i \neq j} |z_i - z_j|$ . Since  $c\delta^2 < 1/2$  for  $\delta < \delta_0(N)$  (if  $\delta_0(N)$  is chosen suitably) we then  $\hat{\tau} \geq 2\delta^2\tilde{\rho}$  and

$$\sum_{j=1}^N \frac{\mu(U_\rho^{\tilde{\tau}}(y_j))}{\rho^n} \leq (1 + c\delta^2) \sum_{j=1}^Q \frac{\mu(U_{\tilde{\rho}}^{\hat{\tau}}(z_j))}{\tilde{\rho}^n},$$

where  $\tilde{\rho} = (1 + c\delta^2)\rho$  and  $c = c(N)$ . Since  $Q \leq N - 1$ , the required result then follows by induction (choosing  $\varepsilon$  appropriately).  $\square$



# Appendix A

## A General Regularity Theorem

We here prove a useful general regularity theorem, which is essentially an abstraction of the “dimension reducing” argument of [Fed70]. There are a number of important applications of this general theorem in the text.

Let  $P \geq n \geq 2$  and let  $\mathcal{F}$  be a collection of functions  $\varphi = (\varphi^1, \dots, \varphi^Q) : \mathbb{R}^P \rightarrow \mathbb{R}^Q$  ( $Q = 1$  is an important case) such that each  $\varphi^j$  is locally  $\mathcal{H}^n$ -integrable on  $\mathbb{R}^P$ . For  $\varphi \in \mathcal{F}$ ,  $y \in \mathbb{R}^P$  and  $\lambda > 0$  we let  $\varphi_{y,\lambda}$  be defined by

$$\varphi_{y,\lambda}(x) = \varphi(y + \lambda x), \quad x \in \mathbb{R}^P.$$

Also, for  $\varphi \in \mathcal{F}$  and a given sequence  $\{\varphi_k\} \subset \mathcal{F}$  we write  $\varphi_k \rightarrow \varphi$  if  $\int \varphi_k f \, d\mathcal{H}^n \rightarrow \int \varphi f \, d\mathcal{H}^n$  (in  $\mathbb{R}^Q$ ) for each given  $f \in C_c^0(\mathbb{R}^P)$ .

We subsequently make the following 3 special assumptions concerning  $\mathcal{F}$ :

**A.1 (Closure under appropriate scaling and translation):** If  $|y| \leq 1 - \lambda$ ,  $0 < \lambda < 1$ , and if  $\varphi \in \mathcal{F}$ , then  $\varphi_{y,\lambda} \in \mathcal{F}$ .

**A.2 (Existence of homogeneous degree zero “tangent functions”):** If  $|y| < 1$ , if  $\{\lambda_k\} \downarrow 0$  and if  $\varphi \in \mathcal{F}$ , then there is a subsequence  $\{\lambda_{k'}\}$  and  $\psi \in \mathcal{F}$  such that  $\varphi_{y,\lambda_{k'}} \rightarrow \psi$  and  $\psi_{0,\lambda} = \psi$  for each  $\lambda > 0$ .

**A.3 (“Singular set” hypotheses):** We assume there is a map

$$\text{sing} : \mathcal{F} \rightarrow \mathcal{C} \quad (= \text{ set of closed subsets of } \mathbb{R}^P)$$

such that:

- (1)  $\text{sing } \varphi = \emptyset$  if  $\varphi \in \mathcal{F}$  is a constant multiple of the indicator function of an  $n$ -dimensional subspace of  $\mathbb{R}^P$ ,
- (2) If  $|y| \leq 1 - \lambda$ ,  $0 < \lambda < 1$ , then  $\text{sing } \varphi_{y,\lambda} = \lambda^{-1}(\text{sing } \varphi - y)$ ,

(3) If  $\varphi, \varphi_k \in \mathcal{F}$  with  $\varphi_k \rightharpoonup \varphi$ , then for each  $\varepsilon > 0$  there is a  $k(\varepsilon)$  such that

$$B_1(0) \cap \text{sing } \varphi_k \subset \{x \in \mathbb{R}^P : \text{dist}(\text{sing } \varphi, x) < \varepsilon\} \quad \forall k \geq k(\varepsilon).$$

We can now state the main result of this section:

**A.4 Theorem.** *Subject to the notation and assumptions A.1, A.2, A.3 above we have*

$$(\ddagger) \quad \dim(B_1(0) \cap \text{sing } \varphi) \leq n - 1 \quad \forall \varphi \in \mathcal{F}.$$

(Here “dim” is Hausdorff dimension, i.e.  $(\ddagger)$  means  $\mathcal{H}^{n-1+\alpha}(\text{sing } \varphi) = 0 \quad \forall \alpha > 0$ .)

*In fact either  $\text{sing } \varphi \cap B_1(0) = \emptyset$  for every  $\varphi \in \mathcal{F}$  or else there is an integer  $d \in [0, n - 1]$  such that*

$$\dim \text{sing } \varphi \cap B_1(0) \leq d \quad \forall \varphi \in \mathcal{F}$$

*and such that there is some  $\psi \in \mathcal{F}$  and a  $d$ -dimensional subspace  $L \subset \mathbb{R}^P$  with*

$$(\ddagger\ddagger) \quad \text{sing } \psi = L \text{ and } \psi_{y,\lambda} = \psi \quad \forall y \in L, \lambda > 0.$$

*If  $d = 0$  then  $\text{sing } \varphi \cap B_\rho(0)$  is finite for each  $\varphi \in \mathcal{F}$  and each  $\rho < 1$ .*

**A.5 Remark:** One readily checks that if  $L$  is an  $n$ -dimensional subspace of  $\mathbb{R}^P$  and  $\psi \in \mathcal{F}$  satisfies A.4  $(\ddagger\ddagger)$ , then  $\psi$  is exactly a constant multiple of the indicator function of  $L$  (hence  $\text{sing } \psi = \emptyset$  by A.3(1)); otherwise we would have  $P > n$  and  $\psi \equiv \text{const.} \neq 0$  on some  $(n + 1)$ -dimensional half-space, thus contradicting the fact that  $\psi$  is locally  $\mathcal{H}^n$ -integrable on  $\mathbb{R}^P$ .

**Proof of A.4:** Assume  $\text{sing } \varphi \cap B_1(0) \neq \emptyset$  for some  $\varphi \in \mathcal{F}$ , and let  $d = \sup\{\dim L : L \text{ is a } d\text{-dimensional subspace of } \mathbb{R}^P \text{ and there is } \varphi \in \mathcal{F} \text{ with } \text{sing } \varphi \neq \emptyset \text{ and } \varphi_{y,\lambda} = \varphi \quad \forall y \in L, \lambda > 0\}$ . Then by A.5 we have  $d \leq n - 1$ .

For a given  $\varphi \in \mathcal{F}$  and  $y \in B_1(0)$  we let  $T(\varphi, y)$  be the set of  $\psi \in \mathcal{F}$  with  $\psi_{0,\lambda} = \psi \quad \forall \lambda > 0$  and with  $\lim \varphi_{y,\lambda_k} = \psi$  for some sequence  $\lambda_k \downarrow 0$ . ( $T(\varphi, y) \neq \emptyset$  by assumption A.2.)

Let  $\ell \geq 0$  and let

$$\mathcal{F}^\ell = \left\{ \varphi \in \mathcal{F} : \mathcal{H}^\ell(\text{sing } \varphi \cap B_1(0)) > 0 \right\}.$$

Our first task is to prove the implication

$$(1) \quad \varphi \in \mathcal{F}^\ell \Rightarrow \exists \psi \in T(\varphi, x) \cap \mathcal{F}^\ell$$

for  $\mathcal{H}^\ell$ -a.e.  $x \in \text{sing } \varphi \cap B_1(0)$ .

To see this, let  $\mathcal{H}_\delta^\ell$  be the “size  $\delta$  approximation” of  $\mathcal{H}^\ell$  as described in §2 of Ch. 1 and recall  $\mathcal{H}^\ell(A) > 0 \iff \mathcal{H}_\infty^\ell(A) > 0$ , so that

$$\mathcal{F}^\ell = \left\{ \varphi \in \mathcal{F} : \mathcal{H}_\infty^\ell(\text{sing } \varphi \cap B_1(0)) > 0 \right\}.$$

Also note that (by 3.8(2) of Ch. 1), for any bounded subset  $A$  of  $\mathbb{R}^P$ ,

$$\mathcal{H}_\infty^\ell(A) > 0 \Rightarrow \Theta^{*n}(\mathcal{H}_\infty^\ell \llcorner A, x) > 0 \quad \text{for } \mathcal{H}^\ell\text{-a.e. } x \in A.$$

Thus we see that if  $\varphi \in \mathcal{F}^\ell$  then for  $\mathcal{H}^\ell$ -a.e.  $x \in \text{sing } \varphi \cap B_1(0)$  we have

$$\Theta^{*\ell}(\mathcal{H}_\infty^\ell \llcorner \text{sing } \varphi, x) > 0.$$

For such  $x$  we thus have a sequence  $\lambda_k \downarrow 0$  such that

$$(2) \quad \lim_{k \rightarrow \infty} \frac{\mathcal{H}_\infty^\ell(\text{sing } \varphi \cap B_{\lambda_k}(x))}{\lambda_k^\ell} > 0,$$

and by assumption A.2 there is a subsequence  $\{\lambda_{k'}\}$  such that  $\varphi_{x, \lambda_{k'}} \rightarrow \psi \in T(\varphi, x)$ . If now  $\mathcal{H}_\infty^\ell(\text{sing } \psi) = 0$ , then for any  $\varepsilon > 0$  we could find open balls  $\{B_{\rho_j}(x_j)\}$  such that

$$\text{sing } \psi \subset \cup_j B_{\rho_j}(x_j)$$

and

$$(3) \quad \sum_j \omega_\ell \rho_j^\ell < \varepsilon$$

(be definition of  $\mathcal{H}_\infty^\ell$ ). Now (2) in particular implies that  $K \equiv B_1(0) \setminus \cup_j B_{\rho_j}(x_j)$  is a compact set with positive distance from  $\text{sing } \psi$ . Hence by assumption A.3(3) we have

$$\text{sing } \varphi_{x, \lambda_{k'}} \cap B_1(0) \subset \cup_j B_{\rho_j}(x_j)$$

for all sufficiently large  $k$ , and hence by (3)

$$\mathcal{H}_\infty^\ell(\text{sing } \varphi_{x, \lambda_{k'}} \cap B_1(0)) < \varepsilon, \quad k \geq k(\varepsilon).$$

Thus since  $\lambda_k^{-1}(\text{sing } \varphi - x) = \text{sing } \varphi_{x, \lambda_k}$  (by A.3(2)) we have

$$\lambda_k^{-\ell} \mathcal{H}_\infty^\ell(\text{sing } \varphi \cap B_{\lambda_k}(x)) < \varepsilon$$

for all sufficiently large  $k$ , thus a contradiction for  $\varepsilon < \lim_{k \rightarrow \infty} \lambda_k^{-\ell} \mathcal{H}_\infty^\ell(\text{sing } \varphi \cap B_{\lambda_k}(x))$ . (Such  $\varepsilon$  can be chosen by (2).)

We have therefore established the general implication (1). From now on take  $\ell > d - 1$  so that  $\mathcal{F}^\ell \neq \emptyset$  (which is automatic for  $\ell \leq d$  by definition of  $d$ ). By (1) there is  $\varphi \in \mathcal{F}^\ell$  with  $\varphi_{0, \lambda} = \varphi \quad \forall \lambda > 0$ . Suppose also that there is a  $k$ -dimensional subspace ( $k \geq 0$ )  $S$  of  $\mathbb{R}^P$  such that  $\varphi_{y, \lambda} = \varphi \quad \forall y \in S, \lambda > 0$ . (Notice of course this is no additional restriction for  $\varphi$  in case  $k = 0$ .) Now if  $k \geq d + 1$  then, by definition of  $d$ , we can assert  $\text{sing } \varphi = \emptyset$ ,

thus contradicting the fact that  $\varphi \in \mathcal{F}^\ell$ . Therefore  $0 \leq k \leq d$ , and if  $k \leq d - 1$  ( $< \ell$ ), then  $\mathcal{H}^\ell(S) = 0$  and in particular

$$(4) \quad \exists x \in B_1(0) \cap \text{sing } \varphi \setminus S.$$

But by A.2 we can choose  $\psi \in T(\varphi, x)$ . Since  $\psi = \lim \varphi_{x, \lambda_j}$  for some sequence  $\lambda_j \downarrow 0$ , we evidently have (since  $\varphi_{y+x, \lambda} = \varphi_{x, \lambda} \ \forall y \in S, \lambda > 0$ )

$$\psi_{y,1} = \lim \varphi_{y+x, \lambda_j} = \lim \varphi_{x, \lambda_j} = \psi \ \forall y \in S$$

and

$$\psi_{\beta x,1} = \lim \varphi_{x+\lambda_j \beta x, \lambda_j} = \psi \ \forall \beta \in \mathbb{R}.$$

(All limits in the weak sense described at the beginning of the section.) Thus  $\psi_{z,\lambda} = \psi$  for each  $\lambda > 0$  and each  $z$  in the  $(k+1)$ -dimensional subspace  $T$  of  $\mathbb{R}^P$  spanned by  $S$  and  $x$ .  $\text{sing } \psi \neq \emptyset$  (by A.3(3)), hence by induction on  $k$  we can take  $k = d - 1$ ; i.e.  $\dim T = d$ , and hence  $\text{sing } \psi \subset T$  by A.3(2). On the other hand if  $\exists \tilde{x} \in \text{sing } \psi \setminus T$  then we can repeat the above argument (beginning at (4)) with  $T$  in place of  $S$  and  $\psi$  in place of  $\varphi$ . This would then give a  $(d+1)$ -dimensional subspace  $\tilde{T}$  and a  $\tilde{\psi} \in \mathcal{F}$  with  $\text{sing } \tilde{\psi} \supset \tilde{T}$ , thus contradicting the definition of  $d$ . Therefore  $\text{sing } \varphi = T$ . Furthermore if  $\ell > d$  then the above induction works up to  $k = d$  and again therefore we would have a contradiction. Thus  $\dim(B_1(0) \cap \text{sing } \varphi) \leq d \ \forall \varphi \in \mathcal{F}$ .

Finally to prove the last claim of the theorem, we suppose that  $d = 0$ . Then we have already established that

$$(5) \quad \mathcal{H}^\alpha(\text{sing } \varphi \cap B_1(0)) = 0 \ \forall \alpha > 0, \ \varphi \in \mathcal{F}.$$

If  $\text{sing } \varphi \cap B_\rho(0)$  is not finite, then we select  $x \in B_\rho(0)$  such that  $x = \lim x_k$  for some sequence  $x_k \in \text{sing } \varphi \cap B_1(0) \setminus \{x\}$ . Then letting  $\lambda_k = 2|x_k - x|$  we see from A.3(2) that there is a subsequence  $\{\lambda_{k'}\}$  with  $\varphi_{x, \lambda_{k'}} \rightarrow \psi \in T(\varphi, x)$  and  $(x_{k'} - x)/|x_{k'} - x| \rightarrow \xi \in \partial B_1(0)$ . Now by A.3(2), (3) we know that  $\{\xi/2\} \cap \{0\} \subset \text{sing } \psi$  and, since  $\psi_{0,\lambda} = \psi$ , this (together with A.3(2)) gives  $L_\xi \subset \text{sing } \psi$  where  $L_\xi$  is the ray determined by 0 and  $\xi$ . Then  $\mathcal{H}^1(\text{sing } \psi \cap B_1(0)) > 0$ , thus contradicting (5), because  $\psi \in \mathcal{F}$ .  $\square$

## Appendix B

# Non-existence of Stable Minimal Hypercones, $n \leq 6$

Here we describe J. Simons [Sim68] result on non-existence in  $\mathbb{R}^{n+1}$  of  $n$ -dimensional stable minimal cones (previously established in case  $n = 2, 3$  by Fleming [Fle62] and Almgren [Alm66] respectively). The proof here follows essential Schoen-Simon-Yau [SSY75], which is a slight variant of the original proof in [Sim68].

Suppose to begin that  $C \in \mathcal{D}_n(\mathbb{R}^{n+1})$  is a cone ( $\eta_{0,\lambda\#}C = C$ ) and  $C$  is integer multiplicity with  $\partial C = 0$ . If  $\text{sing } C \subset \{0\}$  and if  $C$  is minimizing in  $\mathbb{R}^{n+1}$  then, writing  $M = \text{spt } C \setminus \{0\}$  and taking  $M_t$  as in §6 of Ch.2, we have  $\frac{d}{dt}\mathcal{H}^n(M_t)|_{t=0} = 0$  and  $\frac{d^2}{dt^2}\mathcal{H}^n(M_t)|_{t=0} \geq 0$ . (This is clear because in fact  $\mathcal{H}^n(M_t)$  takes its minimum value at  $t = 0$ , by virtue of our assumption that  $C$  is minimizing.) Notice that  $M$  is orientable, with orientation induced from  $C$ , and hence in particular we can deduce from 6.5 of Ch. 2 that

$$B.1 \quad \int_M (|\nabla^M \zeta|^2 - \zeta^2 |A|^2) d\mathcal{H}^n \geq 0$$

for every  $\zeta \in C_c^1(M)$  (notice  $0 \notin M$ , so such  $\zeta$  vanish in a neighborhood of 0). Here  $A$  is the second fundamental form of  $M$  and  $|A|$  is its length, as described in §4 of Ch.2 and 6.5 of Ch.2.

The main result we need is given in the following theorem.

**B.2 Theorem.** *Suppose  $2 \leq n \leq 6$  and  $M$  is an  $n$ -dimensional cone embedded in  $\mathbb{R}^{n+1}$  with zero mean curvature (see §4 of Ch.2) and with  $\overline{M} \setminus M = \{0\}$ , and suppose that  $M$  is stable in the sense that B.1 holds. Then  $\overline{M}$  is a hyperplane.*

As explained above, the hypotheses are in particular satisfied if  $M = \text{spt } C \setminus \{0\}$ , with  $C \in \mathcal{D}_n(\mathbb{R}^{n+1})$  a minimizing cone with  $\partial C = 0$  and  $\text{sing } C \subset \{0\}$ .)

**B.3 Remark:** B.2 is false for  $n = 7$ ; J. Simons [Sim68] was the first to point out that the cone  $M = \left\{ (x^1, \dots, x^8) \in \mathbb{R}^8 : \sum_{i=1}^4 (x^i)^2 = \sum_{i=5}^8 (x^i)^2 \right\}$  is a stable minimal cone. (Notice that  $M$  is the cone over the compact manifold  $(\frac{1}{\sqrt{2}}\mathbb{S}^3) \times (\frac{1}{\sqrt{2}}\mathbb{S}^3) \subset \mathbb{S}^7 \subset \mathbb{R}^8$ .) The fact that the mean curvature of  $M$  is zero is checked by direct computation. The fact that  $M$  is actually *stable* is checked as follows. First, by direct computation one checks that the second fundamental form  $A$  of  $M$  satisfies  $|A|^2 = 6/|x|^2$ .

On the other hand for a stationary hypersurface  $M \subset \mathbb{R}^{n+1}$  the first variation formula ?? of Ch.2 says  $\int \text{div}_M X \, d\mathcal{H}^n = 0$  if  $\text{spt } |X|$  is a compact subset of  $M$ . Taking  $X_x = (\zeta^2/r^2)x$ ,  $\zeta \in C_c^\infty(M)$ ,  $r = |x|$ , and computing as in §4 of Ch.4, we get

$$(n-2) \int_M (\zeta^2/r^2) \, d\mathcal{H}^n = -2 \int_M \zeta r^{-2} x \cdot \nabla^M \zeta \, d\mathcal{H}^n.$$

Using the Schwarz inequality on the right we get

$$\frac{(n-2)^2}{4} \int_M (\zeta^2/r^2) \, d\mathcal{H}^n \leq \int_M |\nabla^M \zeta|^2 \, d\mathcal{H}^n.$$

Thus we have stability for  $M$  (in the sense of B.1) whenever  $A$  satisfies  $|x|^2 |A|^2 \leq (n-2)^2/4$ .

For the example above we have  $n = 7$  and  $|x|^2 |A|^2 = 6$ , so that this inequality is satisfied, and the cone over  $\mathbb{S}^3 \times \mathbb{S}^3$  is stable as claimed. (Similarly the cone over  $\mathbb{S}^q \times \mathbb{S}^q$  is stable for  $q \geq 3$ ; i.e. when the dimension of the cone is  $\geq 7$ .)

Before giving the proof of B.2 we need to derive the identity of J. Simons for the Laplacian of the length of the second fundamental form of a hypersurface (B.8 below).

The simple derivation here assumes the reader's familiarity with basic Riemannian geometry. (A completely elementary derivation, assuming no such background, is described in [Giu84].)

For the moment let  $M$  be an arbitrary hypersurface in  $\mathbb{R}^{n+1}$  ( $M$  not necessarily a cone, and not necessarily having zero mean curvature).

Let  $\tau_1, \dots, \tau_n$  be a locally defined family of smooth vector fields which, together with the unit normal  $\nu$  of  $M$ , define an orthonormal basis for  $\mathbb{R}^{n+1}$  at all points in some region of  $M$ .

The second fundamental form of  $M$  relative to the unit normal  $\nu$  is the tensor  $A = h_{ij} \tau_i \otimes \tau_j$ , where  $h_{ij} = \langle D_{\tau_j} \nu, \tau_i \rangle$ . (Cf. §4 of Ch.2.) Recall (see 4.32 of Ch.2) that

$$B.4 \quad h_{ij} = h_{ji},$$

and, since the Riemann tensor of the ambient space  $\mathbb{R}^{n+1}$  is zero, we have the Codazzi equations

$$B.5 \quad h_{ij,k} = h_{ik,j}, \quad i, j, k \in \{1, \dots, n\}.$$

Here  $h_{ij,k}$  denotes the covariant derivative of  $A$  with respect to  $\tau_k$ ; that is,  $h_{ij,k}$  are such that  $\nabla_{\tau_k} A = h_{ij,k} \tau_i \otimes \tau_j$ .

We also have the Gauss curvature equations

$$B.6 \quad R_{ijk\ell} = h_{i\ell} h_{jk} - h_{ik} h_{j\ell},$$

where  $R = R_{ijk\ell} \tau_i \otimes \tau_j \otimes \tau_k \otimes \tau_\ell$  is the Riemann curvature tensor of  $M$ , and where we use the sign convention such that  $R_{ijji}$  ( $i \neq j$ ) are sectional curvatures of  $M$  ( $= +1$ , if  $M = \mathbb{S}^n$ ).

From the properties of  $R$  (in fact essentially by definition of  $R$ ) we also have, for any 2-tensor  $a_{ij} \tau_i \otimes \tau_j$ ,

$$a_{ij,k\ell} = a_{ij,\ell k} + a_{im} R_{mj\ell k} + a_{mj} R_{mi\ell k}$$

(where  $a_{ij,k\ell}$  means  $a_{ij,k,\ell}$ —i.e. the covariant derivative with respect to  $\tau_\ell$  of the tensor  $a_{ij,k} \tau_i \otimes \tau_j \otimes \tau_k$ ). In particular

$$B.7 \quad \begin{aligned} h_{ij,k\ell} &= h_{ij,\ell k} + h_{im} R_{mj\ell k} + h_{mj} R_{mi\ell k} \\ &= h_{ij,\ell k} + h_{im} [h_{m\ell} h_{jk} - h_{mk} h_{j\ell}] - h_{mj} [h_{i\ell} h_{mk} - h_{ik} h_{m\ell}] \end{aligned}$$

by B.6, where, here and subsequently, repeated indices are summed from 1 to  $n$ .

**B.8 Lemma.** *In the notation above,*

$$\Delta_M \left( \frac{1}{2} |A|^2 \right) = \sum_{i,j,k} h_{ij,k}^2 - |A|^4 + h_{ij} H_{,ij} + H h_{mi} h_{mj} h_{ij},$$

where  $H = h_{kk} = \text{trace } A$ .

**Proof:** We first compute  $h_{ij,kk}$ :

$$\begin{aligned} h_{ij,kk} &= h_{ik,jk} \quad (\text{by B.5}) \\ &= h_{ki,jk} \quad (\text{by B.4}) \\ &= h_{ki,kj} + h_{km} [h_{mj} h_{ik} - h_{mk} h_{ij}] \\ &\quad - h_{mi} [h_{kj} h_{mk} - h_{kk} h_{mj}] \quad (\text{by B.7}) \\ &= h_{ki,kj} - (\sum_{m,k} h_{mk}^2) h_{ij} + h_{kk} h_{mi} h_{mj} \\ &= h_{kk,ij} - (\sum_{m,k} h_{mk}^2) h_{ij} + h_{kk} h_{mi} h_{mj} \quad (\text{by B.5}) \end{aligned}$$

Now multiplying by  $h_{ij}$  we then get (since  $h_{ij}h_{ij,kk} = \frac{1}{2}(\sum_{i,j} h_{ij}^2)_{,kk} - \sum_{i,j,k} h_{ij,k}^2$ )

$$B.9 \quad \frac{1}{2}(\sum_{i,j} h_{ij}^2)_{,kk} = \sum_{i,j,k} h_{ij,k}^2 - (\sum_{i,j} h_{ij}^2)^2 + h_{ij}H_{,ij} + Hh_{mi}h_{mj}h_{ij},$$

which is the required identity.

We now want to examine carefully the term  $\sum_{i,j,k} h_{ij,k}^2$  appearing in the identity of B.8 in case  $M$  is a cone with vertex at 0 (i.e.  $\eta_{0,\lambda}M = M \ \forall \lambda > 0$ ). In particular we want to compare  $\sum_{i,j,k} h_{ij,k}^2$  with  $|\nabla^M A|^2$  in this case. Since  $|\nabla^M A|^2 = \sum_{k=1}^n |A|^{-2}(h_{ij}h_{ij,k})^2$ , we look at the difference

$$B.10 \quad D \equiv \sum_{i,j,k} h_{ij,k}^2 - \sum_{k=1}^n |A|^{-2}(h_{ij}h_{ij,k})^2.$$

**B.11 Lemma.** *If  $M$  is a cone (not necessarily minimal) the quantity  $D$  defined in B.10 satisfies*

$$D(x) \geq 2|x|^{-2}|A(x)|^2, \quad x \in M.$$

**Proof:** Let  $x \in M$  and select the frame  $\tau_1, \dots, \tau_n$  so that  $\tau_n$  is radial ( $x/|x|$ ) along the ray  $\ell_x$  through  $x$ , and so (as vectors in  $\mathbb{R}^{n+1}$ )  $\tau_1, \dots, \tau_n$  are constant along  $\ell_x$ . Then

$$(1) \quad h_{nj} = h_{jn} = 0 \quad \text{on } \ell_x, \quad j = 1, \dots, n,$$

and (since  $h_{ij}(\lambda x) = \lambda^{-1}h_{ij}(x)$ ,  $\lambda > 0$ )

$$(2) \quad h_{ij,n} = -r^{-1}h_{ij} \quad \text{on } \ell_x.$$

Rearranging the expression for  $D$ , we have

$$D = \frac{1}{2} \sum_{k=1}^n \sum_{i,j,r,s=1}^n |A|^{-2} (h_{rs}h_{ij,k} - h_{ij}h_{rs,k})^2,$$

as one easily checks by expanding the square on the right. Now since

$$\sum_{i,j,r,s=1}^n (h_{rs}h_{ij,k} - h_{ij}h_{rs,k})^2 \geq 4 \sum_{\substack{i,j,r=1 \\ s=n}}^{n-1} (h_{rs}h_{ij,k} - h_{ij}h_{rs,k})^2,$$

we thus have

$$D \geq 2|A|^{-2} \sum_{k=1}^n \sum_{i,j,r=1}^{n-1} (h_{ij}h_{rn,k})^2.$$

By the Codazzi equations B.5 and (2) this gives

$$\begin{aligned} D &\geq 2r^{-2}|A|^{-2} \sum_{k=1}^n \sum_{i,j,r=1}^{n-1} h_{ij}^2 h_{rk}^2 \\ &= 2r^{-2}|A|^{-2}|A|^4 \quad (\text{by (1)}) \\ &= 2r^{-2}|A|^2, \end{aligned}$$

as required.  $\square$

**Proof of B.2:** Notice that so far we have not used the minimality of  $M$  (i.e. we have not used  $H(=h_{kk})=0$ ). We now do set  $H=0$  in the above computations, thus giving (by B.8, B.11)

$$(1) \quad \Delta_M(\tfrac{1}{2}|A|^2) + |A|^4 \geq 2r^{-2}|A|^2 + |\nabla|A||^2$$

for the minimal cone  $M$ . (Notice that  $|A|$  is Lipschitz, and hence  $|\nabla|A||$  makes sense  $\mathcal{H}^n$ -a.e. in  $M$ .)

Our aim now is to use (1) in combination with the stability inequality B.1 to get a contradiction in case  $2 \leq n \leq 6$ .

Specifically, replace  $\zeta$  by  $\zeta|A|$  in B.1. This gives

$$(2) \quad \begin{aligned} \int_M \zeta^2 |A|^4 &\leq \int_M |\nabla(\zeta|A|)|^2 \\ &= \int_M (|\nabla\zeta|^2 |A|^2 + \zeta^2 |\nabla|A||^2) + 2 \int_M \zeta |A| \nabla\zeta \cdot \nabla|A|. \end{aligned}$$

Now

$$\begin{aligned} 2 \int_M \zeta |A| \nabla\zeta \cdot \nabla|A| &= 2 \int_M \zeta \nabla\zeta \cdot \nabla(\tfrac{1}{2}|A|^2) \\ &= \int_M (\nabla\zeta^2) \cdot \nabla(\tfrac{1}{2}|A|^2) \\ &= - \int_M \zeta^2 \Delta_M(\tfrac{1}{2}|A|^2) \\ &\leq \int_M (|A|^5 \zeta^2 - 2r^{-2} \zeta^2 |A|^2 + \zeta^2 |\nabla|A||^2) \quad \text{by (1),} \end{aligned}$$

and hence (2) gives

$$(3) \quad 2 \int_M r^{-2} \zeta^2 |A|^2 \leq \int_M |A|^2 |\nabla\zeta|^2 \quad \forall \zeta \in C_c^1(M).$$

Now we claim that (3) is valid even if  $\zeta$  does not have compact support on  $M$ , provided that  $\zeta$  is locally Lipschitz and

$$(4) \quad \int_M r^{-2} \zeta^2 |A|^2 < \infty.$$

(This is proved by applying (3) with  $\zeta\gamma_\varepsilon$  in place of  $\zeta$ , where  $\gamma_\varepsilon$  is such that  $\gamma_\varepsilon(x) \equiv 1$  for  $|x| \in (\varepsilon, \varepsilon^{-1})$ ,  $|\nabla\gamma_\varepsilon(x)| \leq 3/|x|$  for all  $x$ ,  $\gamma_\varepsilon(x) = 0$  for  $|x| < \varepsilon/2$  or  $|x| > 2\varepsilon^{-1}$ , and  $0 \leq \gamma_\varepsilon \leq 1$  everywhere, then letting  $\varepsilon \downarrow 0$  and using (4).)

Since  $M$  is a cone we can write

$$(5) \quad \int_M \varphi(x) d\mathcal{H}^n(x) = \int_0^\infty r^{n-1} \int_\Sigma \varphi(r\omega) d\mathcal{H}^{n-1}(\omega) dr$$

for any non-negative continuous  $\varphi$  on  $M$ , where  $\Sigma = M \cap \mathbb{S}^n$  is a compact  $(n-1)$ -dimensional embedded submanifold. Since  $|A(x)|^2 = r^{-2}|A(x/|x|)|^2$ , we can now use (5) to check that  $\zeta = r^{1+\varepsilon}r_1^{1-n/2-2\varepsilon}$ ,  $r_1 = \max\{1, r\}$ , is a valid choice to ensure (4), hence we may use this choice in (3). This is easily seen to give

$$(6) \quad 2 \int_M r^{2\varepsilon} r_1^{2-n-4\varepsilon} |A|^2 \leq \left(\frac{n}{2} - 2 + \varepsilon\right)^2 \int_{M \cap \{r>1\}} |A|^2 r^{2-n-2\varepsilon} \\ + (1 + \varepsilon)^2 \int_{M \cap \{r<1\}} |A|^2 r^{2\varepsilon} < \infty.$$

For  $2 \leq n \leq 6$  we can choose  $\varepsilon$  such that  $(\frac{n}{2} - 2 + \varepsilon)^2 < 2$  and  $(1 + \varepsilon)^2 < 2$ , hence (6) gives  $|A|^2 \equiv 0$  on  $M$  as required.  $\square$

# Bibliography

- [All72] William K. Allard, *On the first variation of a varifold*, Ann. of Math. (2) **95** (1972), 417–491. MR MR0307015 (46 #6136)
- [Alm66] F. J. Almgren, Jr., *Some interior regularity theorems for minimal surfaces and an extension of Bernstein's theorem*, Ann. of Math. (2) **84** (1966), 277–292. MR MR0200816 (34 #702)
- [Alm68] ———, *Existence and regularity almost everywhere of solutions to elliptic variational problems among surfaces of varying topological type and singularity structure*, Ann. of Math. (2) **87** (1968), 321–391. MR MR0225243 (37 #837)
- [Alm84] ———,  *$Q$ -valued functions minimizing dirichlet's integral and the regularity of area minimizing rectifiable currents up to codimension 2*, Preprint, Princeton University, 1984.
- [Bes28] A. S. Besicovitch, *On the fundamental geometrical properties of linearly measurable plane sets of points*, Math. Ann. **98** (1928), no. 1, 422–464. MR MR1512414
- [Bes38] ———, *On the fundamental geometrical properties of linearly measurable plane sets of points (II)*, Math. Ann. **115** (1938), no. 1, 296–329. MR MR1513189
- [Bes39] ———, *On the fundamental geometrical properties of linearly measurable plane sets of points (III)*, Math. Ann. **116** (1939), no. 1, 349–357. MR MR1513231
- [DG61] Ennio De Giorgi, *Frontiere orientate di misura minima*, Seminario di Matematica della Scuola Normale Superiore di Pisa, 1960-61, Editrice Tecnico Scientifica, Pisa, 1961. MR MR0179651 (31 #3897)
- [Dug86] J. P. Duggan, *Regularity theorems for varifolds with mean curvature*, Indiana Univ. Math. J. **35** (1986), no. 1, 117–144. MR MR825631 (87e:49065)
- [EG92] C. Evans and R. Gariepy, *Measure theory and fine properties of functions*, Studies in Advanced Mathematics, CRC Press, 1992. MR MR0257325 (41 #1976)
- [Fed69] Herbert Federer, *Geometric measure theory*, Die Grundlehren der mathematis-

- chen Wissenschaften, Band 153, Springer-Verlag New York Inc., New York, 1969. MR MR0257325 (41 #1976)
- [Fed70] ———, *The singular sets of area minimizing rectifiable currents with codimension one and of area minimizing flat chains modulo two with arbitrary codimension*, Bull. Amer. Math. Soc. **76** (1970), 767–771. MR MR0260981 (41 #5601)
- [FF60] Herbert Federer and Wendell H. Fleming, *Normal and integral currents*, Ann. of Math. (2) **72** (1960), 458–520. MR MR0123260 (23 #A588)
- [Fle62] Wendell H. Fleming, *On the oriented Plateau problem*, Rend. Circ. Mat. Palermo (2) **11** (1962), 69–90. MR MR0157263 (28 #499)
- [Giu84] Enrico Giusti, *Minimal surfaces and functions of bounded variation*, Monographs in Mathematics, vol. 80, Birkhäuser Verlag, Basel, 1984. MR MR775682 (87a:58041)
- [GT01] David Gilbarg and Neil S. Trudinger, *Elliptic partial differential equations of second order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001, REPRINT of the 1998 edition. MR MR1814364 (2001k:35004)
- [Mir67] M. Miranda, *Sulle singolarità delle frontiere minimali*, Rend. Sem. Mat. Univ. Padova **38** (1967), 180–188. MR MR0221351 (36 #4403)
- [Pro60] M. H. Protter, *Unique continuation for elliptic equations*, Trans. Amer. Math. Soc. **95** (1960), 81–91. MR MR0113030 (22 #3871)
- [Rei60] E. R. Reifenberg, *Solution of the Plateau Problem for  $m$ -dimensional surfaces of varying topological type*, Acta Math. **104** (1960), 1–92. MR MR0114145 (22 #4972)
- [Rei64] ———, *An isoperimetric inequality related to the analyticity of minimal surfaces*, Ann. of Math. (2) **80** (1964), 1–14. MR MR0171197 (30 #1428)
- [Ros84] M. Ross, *Federer's structure theorem*, Research Report, Centre for Mathematical Analysis, Australian National University, 1984.
- [Roy88] H. L. Royden, *Real analysis*, third ed., Macmillan Publishing Company, New York, 1988. MR MR1013117 (90g:00004)
- [Sim68] James Simons, *Minimal varieties in riemannian manifolds*, Ann. of Math. (2) **88** (1968), 62–105. MR MR0233295 (38 #1617)
- [Sim83] Leon Simon, *Asymptotics for a class of nonlinear evolution equations, with applications to geometric problems*, Ann. of Math. (2) **118** (1983), no. 3, 525–571. MR MR727703 (85b:58121)
- [Sol82] B. Solomon, Thesis, Princeton University, 1982.
- [SSY75] R. Schoen, L. Simon, and S. T. Yau, *Curvature estimates for minimal hypersur-*

- faces*, Acta Math. **134** (1975), no. 3-4, 275–288. MR MR0423263 (54 #11243)
- [Whi82] B. White, Thesis, Princeton University, 1982.